

UNIVERSIDADE DE SÃO PAULO

INSTITUTO DE FÍSICA  
CAIXA POSTAL 20516  
01498-970 SÃO PAULO - SP  
BRASIL

# PUBLICAÇÕES

IFUSP/P-1039

**PRECISION TEST OF REACTION THEORY WITH  
EPITHERMAL NEUTRONS AND THE LONGITUDINAL  
ASYMMETRY ANOMALY**

**M.S. Hussein**

Instituto de Física, Universidade de São Paulo

**A.K. Kerman**

Center for Theoretical Physics, Laboratory for  
Nuclear Science and Department of Physics  
Massachusetts Institute of Technology  
Cambridge, Massachusetts, 02139, USA

Fevereiro/1993

# Precision Test of Reaction Theory With Epithermal Neutrons and the Longitudinal Asymmetry Anomaly

M.S. Hussein\*

Nuclear Theory and Elementary Particle Phenomenology Group  
Instituto de Física, Universidade de São Paulo  
C.P. 20516, 01498-970 São Paulo, SP, Brazil

and

A.K. Kerman†

Center for Theoretical Physics, Laboratory for Nuclear Science and  
Department of Physics, Massachusetts Institute of Technology,  
Cambridge, Massachusetts, 02139, USA

February 19, 1993

## Abstract

We analyse the longitudinal asymmetry of epithermal neutron scattering. We show, using the Optical Background Representation, that the energy average of the parity non-conserving matrix element is the optical model one. We also derive an expression for the average longitudinal asymmetry which involves the off-energy-shell PNC matrix element.

\*Supported in part by the CNPq - Brazil, FAPESP - Brazil

†Supported in part by funds provided by the U.S. Department of Energy (D.O.E.) under contract DE-ACO2-76ER03069

## 1 Introduction

The recent discovery of sign correlations in parity non-conserving (PNC) epithermal neutron-induced compound-nucleus reactions involving the heavy nucleus  $^{232}\text{Th}$ , has prompted intensive theoretical discussion of their origin. The great interest in the TRIPLE data[1] stems from the fact that the statistical theory (ST) of these reactions, supposedly quite appropriate at these low energies, though predicts large PNC for individual resonances as the data also show, it rules out any sign correlations in the longitudinal asymmetry contrary to what the data reveals. In fact, according to the discussion of Ref.2), the above mentioned longitudinal asymmetry, defined for the  $M$ th resonance to be

$$P_R = \frac{\sigma_R^{(+)} - \sigma_R^{(-)}}{\sigma_R^{(+)} + \sigma_R^{(-)}} \quad (1)$$

where  $\sigma_R^{(\pm)}$  is the  $R$ -th P-resonance total cross-section for positive (negative) neutron helicity, can be represented as

$$P_R = P_R^{II} + \bar{P} \quad (2)$$

In the Eq.(2),  $P_R^{II}$  averages to zero over many resonances, whereas  $\bar{P}$  exhibits the behaviour

$$\bar{P} = \sqrt{\frac{1\text{eV}}{E_n}} B \quad (3)$$

The TRIPLE data gives for  $B$  the value 0.08.

Several models have been proposed to discuss the large value of  $B$  reported by the TRIPLE collaboration. We mention first the work of Auerbach[3] and Bowman and Auerbach[4], based on the doorway mechanism. The authors assume that the  $n+^{232}\text{Th}$  system is modulated by a  $0^-$  collective state (doorway) that leads to the PNC enhancement of  $B$ . However, it was found that such a mechanism leads to a serious discrepancy when compared with the single particle model. Another proposal for the large value of  $B$  was advanced by Lewenkopf and Weidenmüller[5]. These authors propose a more detailed reaction mechanism based on conventional concepts. They

give arguments that the external mixing, involving compound - direct - compound processes, may be responsible for the enhanced value of B.

Optical model (OM) analysis of the data has also been performed by Koonin, Johnson and Vogel[6] and Carlson and Hussein[7, 8]. In such an analysis quantitative results can be obtained concerning the value of B. The result obtained by Refs.6,7) which is based on the use of a strong parity conserving (PC) complex interaction plus a weak PNC, indicated that the latter interaction is more than two orders of magnitude larger than estimates based on standard meson exchange models. However, as pointed out by Carlson and Hussein[8], care must be taken when confronting the data on the longitudinal asymmetry with the optical model. See also Ref.5.

The reason for this is that the optical model describes the quantity

$$P_{OM} \equiv \frac{\langle \sigma_R^{(+)} - \sigma_R^{(-)} \rangle_E}{\langle \sigma_R^{(+)} + \sigma_R^{(-)} \rangle_E} \quad (4)$$

In contrast the average asymmetry  $\bar{P}$  is defined by

$$\bar{P} \equiv \langle \frac{\sigma_R^{(+)} - \sigma_R^{(-)}}{\sigma_R^{(+)} + \sigma_R^{(-)}} \rangle_{Resonance} \quad (5)$$

Therefore two potentially important differences between  $P_{OM}$  and  $\bar{P}$  can be indicated by writing

$$\bar{P} = P_{OM} + \Delta P_{corr.} + \Delta P_{Erg.} \quad (6)$$

where the term  $\Delta P_{corr.}$  arises from the numerator-denominator correlations and  $\Delta P_{Erg.}$  is due to the replacement of resonance average by energy average (ergotic theorem). From the raw data supplied to us by J.D. Bowman we found out that  $\Delta P_{Erg.}$  is in fact very small. This we obtained by constructing from the data the quantity

$$\bar{P} \equiv \frac{\langle \sigma_R^{(+)} - \sigma_R^{(-)} \rangle_{Resonance}}{\langle \sigma_R^{(+)} + \sigma_R^{(-)} \rangle_{Resonance}} \quad (7)$$

and by writing

$$\bar{P} \cong P_{OM} + \Delta P_{Erg.} \quad (8)$$

Since  $P_{OM}$ , found by reference to be  $6.7 \times 10^{-4}$  (for  $E_n = 1eV$  and the PNC parameter  $\epsilon_7 = 1$ ) and this is about 400 times smaller than  $\bar{P}$ , we reach the conclusion that

$$\bar{P} \cong \Delta P_{corr.} \quad (9)$$

Namely, the optical model supplies the background contribution to  $\bar{P}$ , which is very small as expected. This is so since if no p-resonances were present the fluctuations in  $\sigma_R$  disappear and accordingly  $\Delta P_{corr.}$  becomes zero.

In this contribution to the Weidenmüller Festschrift we analyse the nature of  $\Delta P_{corr.}$ . We do this with the optical background representation of Kawai, Kerman and McVoy[9]. This method allows writing all quantities in terms of optical of the different contributions to  $\bar{P}$  less model dependent. Further, the OBR of Ref.9 supplies a mean through which energy averages can be straight forwardly performed, since the thrust of this method is the separation of the total open channel (elastic in this case) wave function into an optical piece, and a fluctuating whose average is zero.

In the following we present the formal decomposition of  $\bar{P}$  into several physically well defined terms, as done in Ref.5. In contrast to Ref.5, however, these terms are calculable with optical model generated wave and Green functions. We leave the numerical analysis for a future publication.

In Section 2 we give a brief description of the optical background representation of Kawai, Kerman and McVoy. In the same Section the average longitudinal asymmetry is analysed. In Section 3 the different contributions to  $\bar{P}$  are discussed and the term which contains, what Ref.5 calls, the "Barrier Penetration Enhancement" is thoroughly analysed. We show that this term can be written in terms of an off-energy-shell version of the optical matrix element of Refs.6,7,8.

## 2 The Optical Background Representation (OBR) for Epithermal Neutron Scattering

At the very low energies of the TRIPLE experiment, one expects the population of widely spaced, isolated, resonances in the compound nucleus  $^{233}Th$ . To describe the

scattering problem, we use the optical background representation method developed by Kawai, Kerman and McVoy[9]. We also allow for the existence of a single doorway to test its influence.

Before we describe the (OBR) we give in the following, the theoretical ingredients needed to describe  $\bar{P}$ . The interaction between the neutron and the target is described by the Hamiltonian

$$H = H_{PC} + V_{PNC} \quad (10)$$

where  $H_{PC}$  is the strong parity conserving many body Hamiltonian usually employed in reaction theory and  $H_{PNC}$  is the weak parity nonconserving interaction given usually by (in its one-body version).

$$V_{one-body}^{PNC} = \frac{1}{2} \epsilon \gamma \{ f(r), \vec{\sigma} \cdot \vec{p} \} \quad (11)$$

where  $f(r)$  is a form factor that follows the shape of the nuclear density.

The longitudinal asymmetry can be calculated with first order perturbation theory, which gives

$$P_{p1/2} \approx \frac{8\pi^2}{k^2} \frac{Im T^{PNC}(E)}{\frac{1}{2}(\sigma_{p1/2}^+ + \sigma_{p1/2}^-)} = \frac{8\pi^2}{k^2} \frac{Im T^{PNC}(E)}{\sigma_{p1/2}^0(E)} \quad (12)$$

where  $T^{PNC}(E)$  is the distorted wave matrix element

$$T^{PNC}(E) = \langle \Psi^{p(-)}(E) | V_{PNC} | \Psi^{p(+)}(E) \rangle \quad (13)$$

where  $\Psi^{p(+)}(E)$  is the scattering solution for  $s_{1/2}$  wave neutrons, and  $\Psi^{p(-)}(E)$  that of the  $p_-$  wave. In the following we perform the optical background decomposition on  $|\Psi^{p(+)}\rangle$  and  $\langle\Psi^{p(-)}|$ , and thus analyse, in details,  $T^{PNC}(E)$ .

The many-body Schrodinger equation describing the  $n +$  Target system can be written as usual, within the Feshbach formalism

$$\begin{aligned} (E - PHP) P\Psi &= PHQ Q\Psi \\ (E - QHQ) Q\Psi &= QHP P\Psi \end{aligned} \quad (14)$$

where  $P$  projects onto the elastic channel ( $p$  and  $s$  waves), and  $Q$  onto the compound states. We consider all the subspace Hamiltonians to contain a  $PC$  and a  $PNC$  terms. The second equation in (14) can be formally solved for  $Q\Psi$ , and when the solution is inserted into Eq. 14a, we obtain

$$\begin{aligned} (E - PHP - PHQG_Q QHP) P\Psi &= 0 \\ G_Q &= \frac{1}{E - QHQ} \end{aligned} \quad (15)$$

We now perform the OBR on Eq.(15). We do this by writing

$$G_Q = \bar{G}_Q + (G_Q - \bar{G}_Q) \quad (16)$$

where  $\bar{G}_Q$  is the energy-averaged compound propagator which is given by

$$\bar{G}_Q = \frac{1}{E - QHQ + iI/2} \quad (17)$$

with  $I$  denoting the energy interval that contains many compound resonances but still smaller than the width of the doorway.

Clearly, Eq.16 can be rewritten as

$$G_Q = \bar{G}_Q + \bar{G}_Q^{1/2} (iI/2) G_Q \bar{G}_Q^{1/2} \quad (18)$$

and thus we can write formally

$$\begin{aligned} (E - PHP - PHQ \bar{G}_Q QHP) P\Psi &= PHQ \bar{G}_Q^{1/2} \left(\frac{iI}{2}\right)^{1/2} G_Q \left(\frac{iI}{2}\right)^{1/2} \bar{G}_Q^{1/2} QHP P\Psi \\ &\equiv PVQ G_Q QVPP\Psi; \\ PVQ &\equiv PHQ \bar{G}_Q^{1/2} \left(\frac{iI}{2}\right)^{1/2} \end{aligned} \quad (19)$$

We then obtain the desired solution

$$\begin{aligned} P\Psi &= \bar{P}\Psi + G_{opt}^{(+)} PVQ \frac{1}{E - QHQ - QVPG_{opt}^{(+)} PVQ} QVPP\Psi \\ &\equiv \bar{P}\Psi + P\Psi^f \end{aligned} \quad (20)$$

The energy average of the second term on the RHS of Eq.(7) is identically zero by construction. The solution  $\bar{P}\Psi$  is the optical model wave function and  $G_{Opt}^{(+)}$  is

$$G_{Opt}^{(+)} = (E - PHP - PHQ\bar{G}_Q QHP + i\epsilon)^{-1} \quad (21)$$

We are now in a position to analyse the PNC matrix element,  $T^{PNC}$ , of Eq. (13). We generalize the decomposition of Lewenkopf and Weidenmüller for  $T_{PNC}$  to write, within the OBR,

$$T_{PNC} = T_{PP} + T_{PQ} + T_{QP} + T_{QQ} \quad (22)$$

where  $T_{ij} = \langle \Psi^{(-)}(E) | i V j | \Psi^{(+)}(E) \rangle$ . Note that  $T_{PP} \equiv T_{PP}^{Opt} + T_{PP}^{fl}$  where  $T_{PP}^{fl} \equiv \langle P\Psi_{jl}^{(-)} | V_{PNC} | P\Psi_{jl}^{(+)} \rangle$  (see Eq. 22). The quantity  $T_{PP}^{Opt}$  is the optical model generated matrix element.

We now show that the energy average of  $T_{PNC}$  is just the optical model matrix element,  $T_{PP}^{Opt}$ . To do this we write

$$T_{PNC} = T_{PNC}^{Opt} + T_{PNC}^{fluc} \quad (23)$$

Here  $T_{PNC}^{fluc}$  contains eight terms

$$\begin{aligned} T_{PNC}^{fl} = & \langle \bar{P}\Psi_p^{(-)} | V_{PNC} | P\Psi_s^{(+)} \rangle + \langle P\Psi_p^{fl(-)} | V_{PNC} | \bar{P}\Psi_s^{(+)} \rangle \\ & + \langle \bar{P}\Psi_p^{(-)} | V_{PNC} | Q\Psi_s^{(+)} \rangle + \langle Q\Psi_p^{(-)} | V_{PNC} | \bar{P}\Psi_s^{(+)} \rangle \\ & + \langle P\Psi_p^{(-)fl} | V_{PNC} | P\Psi_s^{(+)} \rangle + \langle P\Psi_p^{(-)fl} | V_{PNC} | Q\Psi_s^{(+)} \rangle \\ & + \langle Q\Psi_p^{(-)} | V_{PNC} | P\Psi_s^{(+)} \rangle + \langle Q\Psi_p^{(-)} | V_{PNC} | Q\Psi_s^{(+)} \rangle \end{aligned} \quad (24)$$

On the average, the first four terms in Eq.(27) vanish by construction. Since  $|P\Psi_s^{(+)} \rangle$  and  $|P\Psi_p^{(-)fl} \rangle$  can be written as a sum over isolated s1/2 and p1/2 resonances, respectively (Eq.20).

$$\begin{aligned} |P\Psi_s^{(+)} \rangle &= G_{Opt}^{(+)} \sum_{q'} PVQ | q' \rangle \frac{1}{E - E_{q'} + \frac{i\Gamma_{q'}}{2}} \langle q' | QVP | \bar{P}\Psi_s^{(+)} \rangle \\ \langle P\Psi_p^{(-)fl} | &= \sum_q \langle \bar{P}\Psi_p^{(-)} | PVQ | q \rangle \frac{1}{E - E_q - \frac{i\Gamma_q}{2}} \langle q | QVPG_{Opt}^{(+)} \end{aligned} \quad (25)$$

and  $Q | \Psi \rangle$  as

$$\begin{aligned} Q | \Psi \rangle &= \frac{1}{E - QHQ - QVPG_{Opt}^{(+)} PVQ} QVP | \bar{P}\Psi \rangle \\ &= \sum_{q'} \frac{\langle q' | QVP | \bar{P}\Psi \rangle}{E - E_{q'} + \frac{i\Gamma_{q'}}{2}} \end{aligned} \quad (26)$$

we can, to first order in  $V_{PNC}$ , write the following compact form for  $\langle T_{PNC}^{fl} \rangle$ ,

$$\begin{aligned} \langle T_{PNC}^{fl} \rangle &= \frac{1}{2\pi} \langle \sum_{q'q} \gamma_q^p \langle \frac{1}{E - E_q - i\Gamma_q/2} \langle q | \tilde{V}_{PNC} | q' \rangle \cdot \\ &\quad \cdot \frac{1}{E - E_{q'} + i\Gamma_{q'}/2} \gamma_{q'}^i \rangle \end{aligned} \quad (27)$$

$q' : s1/2, q : p1/2$

In Eq. (27) we have introduced the effective PNC interaction,  $\tilde{V}_{PNC}$ , given by

$$\begin{aligned} \tilde{V}_{PNC} \equiv & V_{PNC} + QVPG_{Opt}^{(+)} V_{NFC} + V_{NFC} G_{Opt}^{(+)} PVQ \\ & + QVPG_{Opt}^{(+)} V_{PNC} G_{Opt}^{(+)} PVQ \end{aligned} \quad (28)$$

Further,

$$\begin{aligned} \gamma_q^i \equiv \langle P\Psi_s^{(-)} | PVQ | q \rangle \sqrt{2\pi} = \langle q | QVP | \bar{P}\Psi_s^{(+)} \rangle \sqrt{2\pi}; \\ \langle \gamma_q^i \rangle_{q=0} = 0 \end{aligned} \quad (29)$$

In the evaluating  $\langle T_{PNC}^{fl} \rangle$  we observe that only the non-diagonal terms in the double sum contribute, and accordingly, using the usual statistical arguments,  $\langle \gamma_q^p \gamma_{q'}^s \rangle = 0$  the energy average vanishes. Accordingly

$$\langle T_{PNC} \rangle = T_{PNC}^{Opt} \quad (30)$$

Eq.(33) is an important result as it allows, as we show in the following, to write a rather compact form for the quantity of interest here namely  $\langle \frac{T_{PNC}}{G_{Opt}^{(+)}} \rangle_{p1/2}$ ,

From the discussion following Eq. (9), we may thus write (See Eq.(26))

$$\begin{aligned} \bar{P} &\cong P_{OM} + \Delta P_{corr.} \\ \Delta P_{corr.} &= \frac{8\pi^2}{k^2} \text{Im} < \frac{T_{PNC}^{f1}}{\sigma_{p1/2}^{(0)}} > , \end{aligned} \quad (31)$$

where we have used  $< \frac{1}{\sigma_{p1/2}^{(0)}} >^{-1} \approx (< \sigma_{p1/2}^{(0)} >)^{-1}$ .

Eq.(31) is the principle result of this section. It expresses  $\Delta P_{corr.}$  as a sum of eight well defined terms (Eq.27), that correspond to different mechanisms for parity non-conservation in the compound nucleus reaction. We call the first four contributions in Eq.27, the direct-compound (DC) mixing, while the last four contributions. The compound-compound (CC) mixing. We remind the reader that room is left for doorway effects to be included. Said differently, one can, if required, take into account the possible effect of a collective O- state by considering the projector, P, in Eq.15, to be composed of the s1/2 and p1/2 channels plus this doorway state. Clearly, the presence of this doorway will also be felt by the optical piece of  $\bar{P}$ ,  $P_{OM}$ , which may become appreciably larger.

The first and third terms in (27) involves s1/2 resonances the second and fourth ones, p1/2 resonances, while the last four terms (the CC mixing) contain both types of resonances. The sum of these four terms leads to the unaveraged version of the RHS of Eq.30, written in terms of  $\bar{V}_{PNC}$  (Eq.28).

### 3 General Discussion and Conclusions

In Ref.(5), the term proportional to  $T_{QQ}$  is said to contain the "dynamical enhancement" of  $P_M$ , whose energy average is purported to be zero. Here we generalize this statement by suggesting that the whole CC mixing, contained in the last four terms in Eq.(24) (which sum to the unaveraged version of Eq.(27). average to zero. Further, these authors indicated that the second DC term in Eq. (24) contains what they called the "barrier penetration enhancement". We write the contribution of this term explicitly

$$\Delta P_{corr}^{BBP} = \frac{8\pi^2}{k^2} < \frac{\text{Im} < P\Psi^{f(-)} | V_{PNC} | \bar{P}\Psi_s^{(+)} >}{\sigma_{p1/2}^{(0)}} > \quad (32)$$

And thus

$$\Delta P_{corr}^{BBP} = \frac{8\pi^2}{k^2} \frac{1}{\sqrt{2\pi}} < \frac{\text{Im} \sum_q \gamma_q^p \frac{1}{E - E_q - i\Gamma_q/2} < q | QVPG_{Op,p}^{(+)}(E) V_{PNC} | \bar{P}\Psi_s^{(+)} >}{\sigma_{p1/2}^{(0)}} > \quad (33)$$

Eq. (33) corresponds to the following process. The population of resonance q from the p1/2 channel, followed by the decay back to the p1/2 channels, which then weakly couples, via  $V_{PNC}$ , to the s1/2 channel.

Note, that in contrast to Ref.5 our channel wave and Green functions are optical ones. This is advantageous as these can be easily generated from convenient optical model codes. The argument given in Ref.(5) to show that  $\Delta P_{corr}^{BBP}$  could potentially be responsible for the sign correlation resides in the observation that the term

$$\begin{aligned} &< q | QVPG_{Op,p}^{(+)} V_{PNC} | \bar{P}\Psi_s^{(+)} > \\ &\equiv \sum_P \int \frac{< q | QVP | \bar{P}\Psi_p^{(+)}(E') > < \bar{P}\Psi_p^{(+)}(E') | V_{PNC} | \bar{P}\Psi_s^{(+)}(E) >}{E - E' + i\epsilon} dE', \end{aligned} \quad (34)$$

contains contributions from all  $E'$ . In fact, the dominant ones would come from  $E' \gg E$  (off-shell) since at these energies the P-wave barrier penetration reduction is absent.

We should emphasize again at this point that within our OBR, the matrix element given in Eq.(34) is represented with optical model wave functions of the type used in Refs.(6) and (7). The dual wave  $< \bar{P}\Psi_p^{(+)}(E') |$  is related to the DWBA wave function  $< \bar{P}\Psi_p^{(-)}(E') |$ , through[10]

$$\begin{aligned} < \bar{P}\Psi_p^{(+)}(E) | &= \sum_p \int dE' < \bar{P}\Psi_p^{(+)}(E) | \bar{P}\Psi_p^{(-)}(E') > < \bar{P}\Psi_p^{(-)}(E') | \\ &\equiv S_p^{-1}(E) < \bar{P}\Psi_p^{(-)}(E) | \end{aligned} \quad (35)$$

Thus, we can write Eq. (37) as

$$\begin{aligned} & \langle q | QVPG_{Opt,p}^{(+)} V_{PNC} | P\Psi_p^{(+)} \rangle = \\ & = \sum_p \int dE' \frac{\langle q | QVP | P\Psi_p^{(+)}(E') \rangle S_p^{-1}(E') \langle P\Psi_p^{(-)}(E') | V_{PNC} | P\Psi_p^{(+)}(E) \rangle}{E - E' + i\epsilon} \end{aligned} \quad (36)$$

In Eq. (36), the weak matrix element  $\langle P\Psi_p^{(-)}(E') | V_{PNC} | P\Psi_p^{(+)} \rangle$  is the off-shell version of the one calculated in Refs. (6) and (7), with appropriate optical potentials. We see clearly that enhancement due to absorption of p1/2 neutrons is manifestly present through the inverse elastic S-matrix element  $\bar{S}_p^{-1}(E')$ .

We turn now to the evaluation of  $\Delta P_{corr}^{BP}$ , Eq.(33). The average can be easily performed by first recognizing that  $\sigma_p^{(0)1/2}$  is just a sum over Breit-Wigner terms

$$\sigma_{p1/2}^{(0)} = \frac{\pi}{k^2} \sum_q \frac{|\gamma_q^p|^2 \Gamma_q}{(E - E_q)^2 + \Gamma_q^2/4} \quad q:p1/2 \quad (37)$$

Thus, at a resonance, q, we have the contribution  $\sigma_p^{(0)1/2}(E_q) = \frac{\pi}{k^2} |\gamma_q^p|^2 / \Gamma_q$ . When this is inserted into Eq.(33) with  $E = E_q$ , we obtain,

$$\Delta P_{corr}^{BP} \simeq 2\sqrt{2\pi} \langle Im \frac{i}{\gamma_q^p} \langle q | QVPG_{Opt,p}^{(+)}(E_q) V_{PNC} | P\Psi_p^{(+)} \rangle \rangle_q, \quad (38)$$

where we have replaced E-average by resonance average, and, owing to the isolated nature of these p1/2 resonances, have taken the phases of  $\gamma_q^p$  to be those of the OM wave function. To proceed further, we recognize from Eq.(34), that the matrix element  $\langle q | QVP | P\Psi_p^{(+)}(E') \rangle$  is just  $\frac{1}{\sqrt{2\pi}} \gamma_q^p(E')$ , where the energy variation arises from the p-wave optical wave function  $| P\Psi_p^{(+)}(E') \rangle$ . The variation of this latter is smooth, and thus we can write, as Ref.5,  $\gamma_q^p(E')/\gamma_q^p(E_q) \equiv C(E')$ , which does not depend on q. With this, the q-average in Eq.(41) is done.

The  $\delta$ -function contribution to (38) is expected to be small since  $\langle P\Psi_p^{(-)}(E_q) |$  is very small at  $E_q \sim 10$  eV. Thus we maintain only the principal part of  $(E_q - E' + i\epsilon)^{-1}$  and ignore  $E_q$  compared to E. We thus find

$$\Delta P_{corr}^{BP} \simeq -2 \int \frac{dE'}{E'} | C(E') | \bar{S}_p^{-1}(E') | Re \langle P\Psi_p^{(-)}(E') | V_{PNC} | P\Psi_p^{(+)}(E) \rangle \quad (39)$$

In Eq.(39), the modulus of  $\bar{S}_p^{-1}(E')$  appears because the phase is canceled by that of  $\gamma_q^p(E')$  and  $\langle P\Psi_p^{(-)}(E') |$ . In Refs. 6,7,8, the optical model matrix element,  $Im \langle P\Psi_p^{(-)}(E) | V_{PNC} | P\Psi_p^{(+)}(E) \rangle$ , was calculated. We see clearly in Eq.(39) that  $\Delta P_{corr}^{BP}$  depends on the off-energy-shell real part.

The contributions of the other DC mixing terms are expected to be small. For the purpose of completeness we give below the contribution of the CC mixing ("dynamical enhancement") to  $\Delta P_{corr}$ . Using the same procedure as above in performing the average, we can write for the contribution of  $\langle Q\Psi_p^{(-)} | V_{PNC} | Q\Psi_p^{(+)} \rangle$ , the following

$$\Delta P_{corr}^{CC} = \langle 2 \sum_{q'} \frac{\langle q' | V_{PNC} | q \rangle \gamma_{q'}^q}{E_q - E_q'} \frac{\gamma_{q'}^q}{\gamma_q^p} \rangle_q, \quad (40)$$

The average above is zero since,  $\gamma_{q'}^q$  and  $\gamma_q^p$  refer to different compound states and channels, thus according to our statistical hypothesis, these partial widths are uncorrelated. From  $\langle \gamma_{q'}^q \rangle_q = 0$ , we thus obtain  $\Delta P_{corr}^{CC} = 0$ . The other CC terms are also expected to average to zero for the same reasons as above.

Finally the contributions of the DC terms,  $\langle P\Psi_p^{(*)} | V_{PNC} | P\Psi_p^{(+)f} \rangle$  and  $\langle P\Psi_p^{(*)} | V_{PNC} | Q\Psi_p^{(+)} \rangle$  are very small since their contribution to  $\Delta P_{corr}$  would involve  $\langle \frac{1}{|\gamma_q^p|^2} \gamma_{q'}^q \rangle_q$ . The last DC term,  $\langle Q\Psi_p^{(-)} | V_{PNC} | P\Psi_p^{(+)} \rangle$  does not contain the barrier penetration enhancement.

It would be certainly interesting to extend the optical model calculation of Ref.(7) to off-shell energies, in order to assess the over all enhancement in  $\Delta P_{corr}^{BP}$ . Detailed investigation of this, as well as the possible rôle of a doorway state, will be published later.

## Acknowledgements

We thank C.H. Lewenkopf for useful comments and discussion.

## References

- [1] C.M. Frankle et al., *Phys. Rev. Lett.* 67 (1991) 564.
- [2] J.D. Bowman et al., *Phys. Rev. Lett* 68 (1992) 780.
- [3] N. Auerbach, *Phys. Rev.* C45 (1992) R514.
- [4] N. Auerbach and J.D. Bowman, *Phys. Rev* C46 (1992) 2582.
- [5] C. Lewenkopf and H.A. Weidenmüller, *Phys. Rev.* C46 (1992) 2601.
- [6] S.E. Koonin, C.W. Johnson and P. Vogel, *Phys. Rev. Lett.* 69 (1992) 1163.
- [7] B.V. Carlson and M.S. Hussein, *Phys. Rev.* C47 (1993) 376.
- [8] B.V. Carlson and M.S. Hussein, to be published.
- [9] M. Kawai, A.K. Kerman and K.W. McVoy, *Ann. Phys. (NY)* 75 (1973) 156.
- [10] M.S. Hussein, *Ann. Phys. (NY)* 175 (1987) 197.