

nyuno: 874222

CERN-TH.7508/94
IFUSP-P 1129

Bounds on Scalar Leptoquarks from Z Physics

J. K. Mizukoshi ^{*},

*Instituto de Física, Universidade de São Paulo,
Caixa Postal 20516, 01452-990 São Paulo, Brazil.*

O. J. P. Éboli [†], and M. C. Gonzalez-Garcia [‡]

*Theory Division, CERN,
CH-1211 Geneva 23, Switzerland.*

Abstract

We analyse the constraints on scalar leptoquarks coming from radiative corrections to Z physics. We perform a global fitting to the LEP data including the contributions of the most general effective Lagrangian for scalar leptoquarks, which exhibits the $SU(2)_L \times U(1)_Y$ gauge invariance. We show that the bounds on leptoquarks that couple to the top quark are much stronger than the ones obtained from low energy experiments.

CERN-TH.7508/94

November 1994

^{*}E-mail: mizukoshi@uspif.if.usp.br (internet) or 47602::MIZUKOSHI (decnet).

[†]Permanent address: Instituto de Física da USP, C. P. 20.516, 01452-990 São Paulo, SP, Brazil.

E-mail: eboli@snfmail.if.usp.br (internet) or 47602::EBOLI (decnet).

[‡]E-mail: concha@vxcern.cern.ch (internet) or VXCERN::CONCHA (decnet).

I. INTRODUCTION

The Standard Model (SM) of electroweak interactions is extremely successful in explaining all the available experimental data [1], which is a striking confirmation of the $SU(2)_L \times U(1)_Y$ invariant interactions involving fermions and gauge bosons. However, not only has the SM some unpleasant features, like the large number of free parameters, but it also leaves some questions answered, such as the reason for the proliferation of fermion generations and their complex pattern of masses and mixing angles. These problems must be addressed by models that are intended to be more fundamental than the SM. A large number of such extensions of the SM predict the existence of colour triplet particles carrying simultaneously leptonic and baryonic number, the so-called leptoquarks. Leptoquarks are present in models that treat quarks and leptons on the same footing, and consequently quark-lepton transitions are allowed. This class of models includes composite models [2], grand unified theories [3], technicolour models [4], and superstring-inspired models [5].

Since leptoquarks are an undeniable signal for physics beyond the SM, there have been several direct searches for them in accelerators. At the CERN Large Electron-Positron Collider (LEP), the experiments established a lower bound $M_{LQ} \gtrsim 45-73$ GeV for scalar leptoquarks [6]. On the other hand, the search for scalar leptoquarks decaying into an electron-jet pair in $p\bar{p}$ colliders constrained their masses to be $M_{LQ} \gtrsim 113$ GeV [7]. Furthermore, the experiments at the DESY ep collider HERA [8] place limits on their masses and couplings, leading to $M_{LQ} \gtrsim 92 - 184$ GeV depending on the leptoquark type and couplings. There have also been many studies of the possibility of observing leptoquarks in the future $p\bar{p}$ [9], ep [10,11], e^+e^- [12], $e\gamma$ [13], and $\gamma\gamma$ [14] colliders.

One can also constrain the masses and couplings of leptoquarks using the low-energy experiments [15,16]. For energies much smaller than the leptoquarks mass, they induce two-lepton-two-quark effective interactions that can give rise to atomic parity violation,

contributions to meson decay, flavour-changing neutral currents (FCNC), and to meson-anti-meson mixing. However, only leptoquarks that couple to first and second generation quarks and leptons are strongly constrained.

In the present work we study the constraints on scalar leptoquarks that can be obtained from their contributions to the radiative corrections to the Z physics. We evaluated the one-loop contribution due to leptoquarks to all LEP observables and made a global fit in order to extract the 95% confidence level limits on the leptoquarks masses and couplings. The more stringent limits are for leptoquarks that couple to the top quarks. Therefore, our results turn out to be complementary to the low energy limits bounds [15,16] since these constrain more strongly first and second generation leptoquarks.

The outline of this paper is as follows. In Sec. II we introduce the effective $SU(2)_L \times U(1)_Y$ invariant effective Lagrangians that we analysed and we also discuss the existing low energy constraints. Section III contains the relevant analytical expressions for the one-loop corrections induced by leptoquarks. Our results and respective discussion are shown in Sec. IV, and we summarize our conclusions in Sec. V. This paper is supplemented with an appendix that contains the relevant Feynman rules.

II. EFFECTIVE INTERACTION AND THEIR LOW ENERGY CONSTRAINTS

A natural hypothesis for theories beyond the SM is that they exhibit the gauge symmetry $SU(2)_L \times U(1)_Y$ above the symmetry breaking scale v . Therefore, we imposed this symmetry on the leptoquark interactions. In order to avoid strong bounds coming from the proton lifetime experiments, we required baryon (B) and lepton (L) number conservation, which forbids the leptoquarks to couple to diquarks. The most general effective Lagrangian for leptoquarks satisfying the above requirements and electric charge and colour conservation is given by [10]

$$\begin{aligned} \mathcal{L}_{\text{eff}} &= \mathcal{L}_{F=2} + \mathcal{L}_{F=0}, \\ \mathcal{L}_{F=2} &= (g_{1L} \bar{q}_L^i i\tau_2 \ell_L + g_{1R} \bar{u}_R^c e_R) S_1 + \bar{g}_{1R} \bar{d}_R^c e_R \tilde{S}_1 + g_{3L} \bar{q}_L^i i\tau_2 \vec{\tau} \ell_L \cdot \vec{S}_3, \end{aligned} \quad (1)$$

$$\mathcal{L}_{F=0} = h_{2L} R_2^T \bar{u}_R i\tau_2 \ell_L + h_{2R} \bar{q}_L e_R R_2 + \bar{h}_{2L} \bar{R}_2^T \bar{d}_R i\tau_2 \ell_L,$$

where $F = 3B + L$, q (ℓ) stands for the left-handed quark (lepton) doublet, and u_R , d_R , and e_R are the singlet components of the fermions. We denote the charge conjugated fermion fields by $\psi^c = C\bar{\psi}^T$ and we omitted in (1) the flavour indices of the couplings to fermions and leptoquarks. The leptoquarks S_1 and \tilde{S}_1 are singlets under $SU(2)_L$ while R_2 and \tilde{R}_2 are doublets, and S_3 is a triplet. Furthermore, we assumed in this work that the leptoquarks belonging to a given $SU(2)_L$ multiplet are degenerate in mass, with their mass denoted by M .

Local invariance under $SU(2)_L \times U(1)_Y$ implies that leptoquarks also couple to the electroweak gauge bosons. To obtain the couplings to W^\pm , Z , and γ , we substituted ∂_μ by the electroweak covariant derivative in the leptoquark kinetic Lagrangian:

$$\mathcal{L}_{\text{kin}} = (D_\mu \Phi)^\dagger D^\mu \Phi, \quad (2)$$

with

$$D_\mu \Phi = \left[\partial_\mu - i \frac{e}{\sqrt{2}s_W} (W_\mu^+ T^+ + W_\mu^- T^-) - ieQ_Z Z_\mu + ieQ^\gamma A_\mu \right] \Phi, \quad (3)$$

where Φ stands for the leptoquarks interpolating fields, Q^γ is the electric charge matrix of the leptoquarks, s_W is the sine of the weak mixing angle, and the T 's are the generators of $SU(2)_L$ for the representation of the leptoquarks. The weak neutral charge Q_Z is given by

$$Q_Z = \frac{T_3 - s_W^2 Q^\gamma}{s_W c_W}. \quad (4)$$

In the last reference of [12], for instance, there is a table with all the quantum numbers for all scalar leptoquarks. In the Appendix we present the Feynman rules for the interactions defined by Eqs. (1) and (2).

Low-energy experiments can lead to strong bounds on the couplings and masses of leptoquarks, due to induced four fermion interactions:

- Leptoquarks can give rise to FCNC processes if they couple to more than one family of quarks or leptons [17,18]. In order to avoid strong bounds from FCNCs, we assumed

that the leptoquarks couple to a single generation of quarks and a single one of leptons. However, due to mixing effects on the quark sector, there is still some amount of FCNC [15] and, therefore, leptoquarks that couple to the first two generations of quarks must comply with some low-energy bounds [15].

- The analyses of the decays of pseudoscalar mesons, like the pions, put stringent bounds on leptoquarks unless their coupling is chiral – that is, it is either left-handed or right-handed [17].

- Leptoquarks that couple to the first family of quarks and leptons are strongly constrained by atomic parity violation [19]. In this case, there is no choice of couplings that avoids the strong limits.

It is interesting to keep in mind that the low-energy data constrain the masses of the first generation leptoquarks to be bigger than 0.5–1 TeV when the coupling constants are equal to the electromagnetic coupling e [15].

III. ANALYTICAL EXPRESSIONS

In this work we employed the on-shell-renormalization scheme, adopting the conventions of Ref. [20]. We used as inputs the fermion masses, G_F , α_{em} , and the Z mass, and the electroweak mixing angle being a derived quantity that is defined through $\sin^2\theta_w = s_w^2 \equiv 1 - M_Z^2/M_W^2$. We evaluated the loops integrals using dimensional regularization [21], since it is a gauge-invariant regularization procedure, and we adopted the Feynman gauge to perform the calculations.

Close to the Z resonance, the physics can be summarized by the effective neutral current

$$J_\mu = (\sqrt{2}G_F M_Z^2 \rho_f)^{1/2} [(I_3^f - 2Q^f s_w^2 \kappa_f) \gamma_\mu - I_3^f \gamma_\mu \gamma_5], \quad (5)$$

where Q^f (I_3^f) is the fermion electric charge (third component of weak isospin). The form factors ρ_f and κ_f have universal contributions, *i.e.* independent of the fermion species, as well as non-universal parts:

$$\rho_f = 1 + \Delta\rho_{\text{univ}} + \Delta\rho_{\text{non}}, \quad (6)$$

$$\kappa_f = 1 + \Delta\kappa_{\text{univ}} + \Delta\kappa_{\text{non}}. \quad (7)$$

Leptoquarks can affect the physics at the Z pole through their contributions to both universal and non-universal corrections. The universal contributions can be expressed in terms of the unrenormalized vector boson self-energy (Σ) as

$$\Delta\rho_{\text{univ}}^{LQ}(s) = -\frac{\Sigma_{LQ}^Z(s) - \Sigma_{LQ}^Z(M_Z^2)}{s - M_Z^2} + \frac{\Sigma_{LQ}^Z(M_Z^2)}{M_Z^2} - \frac{\Sigma_{LQ}^W(0)}{M_W^2} - 2\frac{s_w}{c_w} \frac{\Sigma_{LQ}^Z(0)}{M_Z^2} - \chi_e - \chi_\mu, \quad (8)$$

$$\Delta\kappa_{\text{univ}}^{LQ} = -\frac{c_w}{s_w} \frac{\Sigma_{LQ}^Z(M_Z^2)}{M_Z^2} - \frac{c_w}{s_w} \frac{\Sigma_{LQ}^Z(0)}{M_Z^2} + \frac{c_w^2}{s_w^2} \left[\frac{\Sigma_{LQ}^Z(M_Z^2)}{M_Z^2} - \frac{\Sigma_{LQ}^W(M_W^2)}{M_W^2} \right], \quad (9)$$

where the factors χ_ℓ are defined below. The diagrams with leptoquark contributions to the self-energies are shown in Fig. 1. They can be easily evaluated, yielding

$$\Sigma_{LQ}^V(k^2) = -\frac{\alpha_{\text{em}}}{4\pi} N_c \sum_j (Q_j^V)^2 \mathcal{H}(k^2, M^2), \quad (10)$$

$$\Sigma_{LQ}^Z(k^2) = -\frac{\alpha_{\text{em}}}{4\pi} N_c \sum_j (Q_j^Z)^2 \mathcal{H}(k^2, M^2), \quad (11)$$

$$\Sigma_{LQ}^{\gamma Z}(k^2) = \frac{\alpha_{\text{em}}}{4\pi} N_c \sum_j Q_j^V Q_j^Z \mathcal{H}(k^2, M^2), \quad (12)$$

$$\Sigma_{LQ}^W(k^2) = -\frac{\alpha_{\text{em}}}{4\pi s_w^2} N_c \sum_j (T_3^j)^2 \mathcal{H}(k^2, M^2), \quad (13)$$

where $N_c = 3$ is the number of colours and the sum is over all members of the leptoquark multiplet. The function \mathcal{H} is defined according to:

$$\mathcal{H}(k^2, M^2) = -\frac{k^2}{3} \Delta_M - \frac{2}{9} k^2 - \frac{4M^2 - k^2}{3} \int_0^1 dx \ln \left[\frac{x^2 k^2 - x k^2 + M^2 - i\epsilon}{M^2} \right], \quad (14)$$

with

$$\Delta_M = \frac{2}{4-d} - \gamma_E + \ln(4\pi) - \ln \left(\frac{M^2}{\mu^2} \right), \quad (15)$$

and d being the number of dimensions. In this section we denote the leptoquark masses by capital M with no subindex.

The factors χ_ℓ ($\ell = e, \mu$) stem from corrections to the effective coupling between the W and fermions at low energy. Leptoquarks modify this coupling through the diagrams shown in Fig. 2, inducing a contribution that we parametrize as

$$i \frac{e}{\sqrt{2}s_W} \chi_\ell \gamma_\mu P_L, \quad (16)$$

where P_L (P_R) is the left-handed (right-handed) projector and ℓ stands for the lepton flavour. Since this correction modifies the muon decay, it contributes to Δr , and consequently, to $\Delta\rho_{\text{univ}}$. Leptoquarks with right-handed couplings, as well as the $F = 0$ ones, do not contribute to χ_ℓ , while the evaluation of the diagrams of Fig. 2, for left-handed leptoquarks in the $F = 2$ sector, leads to

$$\begin{aligned} \chi_\ell = & \lim_{p_W^2 \rightarrow 0} \frac{g_{LQ}^2}{16\pi^2} N_c \left\{ \frac{1}{2} \sum_{j,q} M_{\ell q}^{j\dagger} M_{q\ell}^j B_1(0, m_q^2, M^2) + \frac{1}{2} \sum_{j,q'} M_{\nu q'}^{j\dagger} M_{q'\nu}^j B_1(0, m_{q'}^2, M^2) \right. \\ & \sum_{q, q', j} M_{\ell q}^{j\dagger} M_{q'\nu}^j V_{qq'} \left[(2-d) C_{00}(0, p_W^2, 0, M^2, m_q^2, m_{q'}^2) + p_W^2 C_{12}(0, p_W^2, 0, M^2, m_q^2, m_{q'}^2) \right] \\ & \left. - 2 \sum_{q, j, j'} T_{jj'} M_{\ell q}^{j\dagger} M_{q\nu}^j C_{00}(0, p_W^2, 0, m_q^2, M^2, M^2) \right\}, \quad (17) \end{aligned}$$

where j stands for the leptoquarks belonging to a given multiplet and $V_{qq'}$ is the Cabibbo-Kobayashi-Maskawa mixing matrix for the quarks, which, for simplicity, we will take to be the unit matrix¹. The functions B_1 , C_0 , C_{00} , and C_{12} are the Passarino-Veltman functions [22]. It is interesting to notice that for degenerate massless quarks the above expression vanishes, and none of the leptoquarks contribute to χ_ℓ . On the other hand, for the leptoquarks coupling to the third family (neglecting the bottom quark mass), we have that

$$\chi_\ell = \frac{g_{1L}^2}{64\pi^2} N_c \left\{ \frac{m_{\text{top}}^2}{M^2 - m_{\text{top}}^2} + \frac{m_{\text{top}}^4 - 2m_{\text{top}}^2 M^2}{(M^2 - m_{\text{top}}^2)^2} \ln \left(\frac{M^2}{m_{\text{top}}^2} \right) \right\}, \quad (18)$$

for the S_{1L} leptoquark, while the S_{3L} contribution can be obtained from (18) by the substitution $g_{1L}^2 \Rightarrow -g_{3L}^2$.

¹In general, this expression is divergent and requires a renormalization of the elements of the CKM matrix.

Corrections to the vertex Zff give rise to non-universal contributions to ρ_f and κ_f . Leptoquarks affect these couplings of the Z through the diagrams given in Fig. 3 whose results we parametrize as

$$i \frac{e}{2s_W c_W} \left[\gamma_\mu F_{VLQ}^{Zf} - \gamma_\mu \gamma_5 F_{ALQ}^{Zf} + I_3^f \gamma_\mu (1 - \gamma_5) \frac{c_W}{s_W} \frac{\Sigma_{LQ}^{Zf}(0)}{M_2^2} \right], \quad (19)$$

where for leptons (ℓ) and leptoquarks with $F = 2$

$$\begin{aligned} F_{VLQ}^{\ell} = & \pm F_{ALQ}^{\ell} = \frac{g_{LQ}^2}{32\pi^2} N_c \sum_{j,q} M_{\ell q}^{j\dagger} M_{q\ell}^j \\ & \left\{ \frac{g_X^2}{2} - s_W c_W Q_Z^j - (g_X^2 + 2s_W c_W Q_Z^j) \frac{M^2 - m_q^2}{M_2^2} \left[-\frac{1}{2} \ln \left(\frac{M^2}{m_q^2} \right) + \bar{B}_0(0, m_q^2, M^2) \right] \right. \\ & + 2s_W c_W Q_Z^j \frac{M^2 - m_q^2 - \frac{1}{2} M_2^2}{M_2^2} \left[-\ln \left(\frac{M^2}{m_q^2} \right) + \bar{B}_0(M_2^2, M^2, M^2) \right] \\ & + g_X^2 \frac{M^2 - m_q^2 - \frac{1}{2} M_2^2}{M_2^2} \bar{B}_0(M_2^2, m_q^2, m_q^2) + g_X^2 \bar{B}_1(0, m_q^2, M^2) \\ & + \left[g_{\pm X}^2 m_q^2 + g_X^2 \frac{(M^2 - m_q^2)^2}{M_2^2} \right] C_0(0, M_2^2, 0, M^2, m_q^2, m_q^2) \\ & \left. - 2s_W c_W Q_Z^j \frac{(M^2 - m_q^2)^2 + m_q^2 M_2^2}{M_2^2} C_0(0, M_2^2, 0, m_q^2, M^2, M^2) \right\}, \quad (20) \end{aligned}$$

where the $+$ ($-$) corresponds to left- (right-) handed leptoquarks and $g_{L/R}^f = v^f \mp a^f$, the neutral current couplings being $a_f = I_3^f$ and $v_f = I_3^f - 2Q^f s_W^2$. We used the convention $X = L, R$ and $-L = R$ ($-R = L$). We also defined

$$B_0(k^2, M^2, M^2) \equiv \frac{1}{2} \Delta_M + \frac{1}{2} \Delta_{M'} + \bar{B}_0(k^2, M^2, M^2), \quad (21)$$

$$B_1(k^2, M^2, M^2) \equiv -\frac{1}{2} \Delta_M + \bar{B}_1(k^2, M^2, M^2), \quad (22)$$

with Δ_M given by Eq. (15). From this last expression we can obtain the effect of $F = 2$ leptoquarks on the vertex $Zq\bar{q}$ simply by the change $\ell \Leftrightarrow q$. Moreover, we can also employ the expression (20) to $F = 0$ leptoquarks provided we substitute $g_{LQ,X} \Rightarrow h_{LQ,X}$ and $g_{\pm X}^2 \Rightarrow -g_{\pm X}^2$.

With all this we have

$$\Delta\rho_{\text{non}}^{LQ} = \frac{F_{ALQ}^{Zf}}{a_f} (M_2^2), \quad (23)$$

$$\Delta\kappa_{\text{non}}^{LQ} = -\frac{1}{2s_W^2 Q^f} \left[F_{VLQ}^{Zf}(M_2^2) - \frac{v_f}{a_f} F_{ALQ}^{Zf}(M_2^2) \right]. \quad (24)$$

One very interesting property of the general leptoquark interactions that we are analysing is that all the physical observables are rendered finite by using the same counter-terms as appear in the SM calculations [20]. For instance, starting from the unrenormalized self-energies (10-13) and the mass and wave-function counter-terms we obtain finite expression for the two-point functions of vector bosons. Moreover, the contributions to the vertex functions $Zf\bar{f}$ and $Wf\bar{f}'$ are finite, as can be seen from the explicit expressions (17) and (20).

In order to check the consistency of our calculations, we analysed the effect of leptoquarks to the $\gamma f\bar{f}$ vertex at zero momentum, which is used as one of the renormalization conditions in the on-shell scheme. This vertex function can be obtained from (20) by the substitutions $Q_Z^j \Rightarrow -Q_j^f$, $g_X^j = Q_j^f$, $e/2s_W c_W \Rightarrow -e$, and $M_Z^2 \Rightarrow k^2$, with k^2 being the squared four momentum of the photon. It turns out that the leptoquark contribution to the vertex function $\gamma f\bar{f}$ not only is finite but also vanishes at $k^2 = 0$ for all fermion species. Therefore, our expressions for the different leptoquark contributions satisfy the appropriate QED Ward identities [23], and leave the fermion electric charges unchanged. Moreover, we also verified explicitly that the leptoquarks decouple in the limit of large M .

IV. RESULTS AND DISCUSSION

The above expressions for the radiative corrections to Z physics due to leptoquarks are valid for couplings and masses of any leptoquark. For the sake of simplicity, we assumed that the leptoquarks couple to leptons and quarks of the same family. In the conclusions, we comment on how we can extend our results to other cases.

In order to gain some insight on which corrections are the most relevant, let us begin our analyses by studying just the oblique corrections [24], which can be parametrized in terms of the variables ϵ_1 , ϵ_2 , and ϵ_3 . The parameter ϵ_1 vanishes since it is proportional to $\Delta\rho(0)$ and we assumed the multiplets to be degenerate. The parameter ϵ_3 also vanishes, while

$$\epsilon_2 = \frac{2N_c}{\pi} \sum_j (T_3^j)^2 \left\{ -\left(\frac{1}{6} - \frac{2}{3} \frac{M^2}{M_W^2}\right) \bar{B}_0(M_W^2, M^2, M^2) + \left(\frac{1}{6} - \frac{2}{3} \frac{M^2}{M_Z^2}\right) \bar{B}_0(M_Z^2, M^2, M^2) \right\}, \quad (25)$$

where the sum is over all members of a given multiplet. Leptoquarks that are singlets under $SU(2)_L$ also lead to $\epsilon_2 = 0$. Notice that the above results depend only upon the interaction of leptoquarks with the gauge bosons. Imposing that the leptoquark contribution to ϵ_2 must be within the limits allowed by the LEP data [25], we find out that the constraints coming from oblique corrections are less restrictive than the available experimental limits [6-8]. Therefore, the contribution of the leptoquarks to the oblique parameters are very small and do not lead to new bounds.

We then performed a global fit to all LEP data including both universal and non-universal contributions. In Table I we show the most recent combined results of the four LEP experiments. These results are not statistically independent and the correlation matrix can be found in [1]. We can express the theoretical predictions to these observables in terms of κ^f , ρ^f , and Δr , with the SM contributions being obtained from the program ZFITTER [26]. In order to perform the global fit we constructed the χ^2 function and minimized it using the package MINUIT. In our fit we used five parameters, three from the SM (m_{top} , M_H , and $\alpha_s(M_Z^2)$) and two new ones: M and g_{LQ} . We present our results as 95% CL lower limits in the leptoquark mass and study the dependence of these limits upon all the other parameters.

The parameter of the SM that most strongly affects our results is the top mass (m_{top}), as expected. For this reason, Figs. 4 exhibit the 95% CL limits obtained for a third generation leptoquark as a function of m_{top} for several values of the coupling constants g_{LQ} ($= \sqrt{4\pi}$, 1, and e/s_W). In these figures, we fixed the value of $M_H = 300$ GeV and $\alpha_s(M_Z^2) = 0.126$, which are the best values obtained from a fit in the framework of the SM [1]. We can see from these figures that the limits on the leptoquarks S_1 , S_3 , and R_2 become stronger as m_{top} increases. This result is analogous to the SM result for the radiative corrections to $Z \rightarrow b\bar{b}$

[27], where there is an enhancement by powers of the top quark mass, as can be seen, for instance, in Eq. (18). We can also learn from these figures that the limits are better for left-handed leptoquarks than for right-handed ones, given a leptoquark type.

The contributions from \tilde{R}_2 and \tilde{S}_1 are not enhanced by powers of the top quark mass since these leptoquarks do not couple directly to up-type quarks. Therefore, the limits are much weaker, depending on m_{top} only through the SM contribution, and the bounds for these leptoquarks are worse than the present discovery limits unless they are strongly coupled ($g_{LQ}^2 = 4\pi$). Therefore, we do not plot the bounds for these cases. Moreover, the limits on first and second generation leptoquarks are also uninteresting for the same reason. Nevertheless, if we allow leptoquarks to mix the third generation of quarks with leptons of another generation the bounds obtained are basically the same as the ones discussed above², since the main contribution to the constraints comes from the Z widths.

Let us finally comment on the dependence of our bounds on other SM parameters besides m_{top} . For a fixed m_{top} , the dependence on $\alpha_s(M_Z^2)$ and M_H is rather weak, but it shows the pattern that the limits are more stringent as $\alpha_s(M_Z^2)$ increases and M_H decreases. For instance, the limits vary by about 10% when $\alpha_s(M_Z^2)$ is in the interval $0.12 < \alpha_s(M_Z^2) < 0.13$ and by about 25% for $60 < M_H < 1000$ GeV.

V. SUMMARY AND CONCLUSIONS

We summarize our results in Table II, where we assumed a top mass of 175 GeV³. Our analyses show that the LEP data constrain more strongly the leptoquarks that couple to the top quark. Since the most important ingredients in the bounds are the widths of the Z , we can conclude that our results should also give a good estimate for the cases where

²In the case of first generation leptons, we must also add a tree level t -channel leptoquark exchange to some observables.

³Our results for R_{2L} are in qualitative agreement with those of Ref. [28].

the leptoquarks couple to quarks and leptons from different families. It is interesting to notice that the constraints on leptoquarks coming from LEP data are complementary to the low-energy ones, since these are more stringent for leptoquarks that couple to the first two families, while the former are stronger when the coupling is to the top quark.








The bounds on scalars leptoquarks coming from low-energy and Z physics exclude large regions of the parameter space where the new collider experiments could search for these particles, however, not all of it [9,11–14]. Nevertheless, we should keep in mind that nothing substitutes the direct observation.

ACKNOWLEDGEMENTS

We want to thank D. Bardine and A. Okevsy for providing us with the latest version of ZFITTER. O.J.P.E. is grateful to the CERN Theory Division for its kind hospitality. This work was supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq-Brazil) and by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP).


APPENDIX: FEYNMAN RULES FOR SCALAR LEPTOQUARKS

In this section we denote by Φ the scalar leptoquark multiplet that is a vector in the weak isospin space. The Feynman rules for the gauge couplings of scalar leptoquarks are




The couplings to fermions are different for $F = 2$ and $F = 0$ leptoquarks. They can be parametrized as:

For $F=2$




$$ig_X M_{qt}^j CP_X$$




$$-ig_X M_{lq}^{j\dagger} P_{-X} C^\dagger$$

For $F=0$



$$ig_X M_{qt}^j P_X$$



$$ig_X M_{lq}^{j\dagger} P_{-X}$$

where for S_1

$$M_L = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad M_R = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad (\text{A1})$$

while for \bar{S}_1

$$M_R = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}. \quad (\text{A2})$$

The triplet S_3 in the diagonal basis $S_3 = (S_3^+, S_3^0, S_3^-)$ leads to

$$M_L^+ = \sqrt{2} \begin{pmatrix} 0 & 0 \\ 0 & -1 \end{pmatrix}, \quad M_L^0 = \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}, \quad M_L^- = \sqrt{2} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad (\text{A3})$$

and the doublet $R_2 = (R_2^{1/2}, R_2^{-1/2})$ is associated to

$$M_L^+ = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad M_L^- = \begin{pmatrix} -1 & 0 \\ 0 & 0 \end{pmatrix}, \quad (\text{A4})$$

$$M_R^+ = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad M_R^- = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}.$$

For \tilde{R}_2 we have

$$M_L^+ = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \quad M_L^- = \begin{pmatrix} 0 & 0 \\ -1 & 0 \end{pmatrix}. \quad (\text{A5})$$

REFERENCES

- [1] See, for instance, D. Schaile, *Precision Tests of the Electroweak Interaction*, plenary talk (Pl-2) presented at the 27th International Conference on High Energy Physics, Glasgow, Scotland, July 1994; "Combined Preliminary data on Z^0 Parameters from the LEP experiments and Constraints on the Standard Model", The LEP Collaborations and The LEP Electroweak Working Group, report in preparation.
- [2] See, for instance, W. Buchmüller, *Acta Phys. Austriaca Suppl.* **XXVII** (1985) 517.
- [3] See, for instance, P. Langacker, *Phys. Rep.* **72** (1981) 185.
- [4] See, for instance, E. Farhi and L. Susskind, *Phys. Rep.* **74** (1981) 277.
- [5] See, for instance, J. L. Hewett and T. G. Rizzo, *Phys. Rep.* **183** (1989) 193.
- [6] L3 Collaboration, B. Adeva *et al.*, *Phys. Lett.* **B261** (1992) 169; OPAL Collaboration, G. Alexander *et al.*, *Phys. Lett.* **B263** (1992) 123; DELPHI Collaboration, P. Abreu *et al.*, *Phys. Lett.* **B316** (1993) 620.
- [7] CDF Collaboration, F. Abe *et al.*, *Phys. Rev.* **D48** (1993) 3939; D0 Collaboration, *Phys. Rev. Lett.* **72** (1994) 965.
- [8] ZEUS Collaboration, M. Derrick *et al.*, *Phys. Lett.* **B306** (1993) 173; H1 Collaboration, I. Abt *et al.*, *Nucl. Phys.* **B396** (1993) 3.
- [9] O. J. P. Éboli and A. V. Olinto, *Phys. Rev.* **D38** (1988) 3461; J. L. Hewett and S. Pakvasa, *ibid.* **D37** (1988) 3165; J. Ohnemus, S. Rudaz, T. F. Wash, and P. Zerwas, *Phys. Lett.* **B334** (1994) 203.
- [10] W. Buchmüller, R. Rückl, and D. Wyler, *Phys. Lett.* **B191** (1987) 442.
- [11] J. Wudka, *Phys. Lett.* **B167** (1986) 337; M. A. Doncheski and J. L. Hewett, *Z. Phys.* **C56** (1992) 209; J. Blümlein, F. Boos, and A. Pukhov, preprint DESY-94-072, hep-ph/9404321.
- [12] J. L. Hewett and T. G. Rizzo, *Phys. Rev.* **D36** (1987) 3367; J. L. Hewett and S. Pakvasa *Phys. Lett.* **B227** (1987) 178; J. E. Cieza and O. J. P. Éboli, *Phys. Rev.* **D47** (1993) 837; J. Blümlein and R. Rückl, *Phys. Lett.* **B304** (1993) 337; J. Blümlein and E. Boos, preprint DESY-94-114, hep-ph/9408250.
- [13] O. J. P. Éboli *et al.*, *Phys. Lett.* **B311** (1993) 147; H. Nadeau and D. London, *Phys. Rev.* **D47**, (1993) 3742; M. A. Doncheski and S. Godfrey, University of Ottawa preprint OCIP-C-94-7, hep-ph/9407317.
- [14] G. Bélanger, D. London, and H. Nadeau, *Phys. Rev.* **D49** (1994) 3140.
- [15] M. Leurer, *Phys. Rev. Lett.* **71** (1993) 1324; *Phys. Rev.* **D49** (1994) 333.
- [16] S. Davidson, D. Bailey, and A. Campbell, *Z. Phys.* **C61** (1994) 613.
- [17] O. Shanker, *Nucl. Phys.* **B204** (1982) 375.
- [18] W. Buchmüller and D. Wyler, *Phys. Lett.* **B177** (1986) 377; J. C. Pati and A. Salam, *Phys. Rev.* **D10** (1974) 275.
- [19] P. Langacker, *Phys. Lett.* **B256** (1991) 277; P. Langacker, M. Luo, and A. K. Mann, *Rev. Mod. Phys.* **64** (1992) 87.
- [20] See, for example, W. Hollik, in the Proceedings of the VII Swieca Summer School, editors O. J. P. Éboli and V. O. Rivelles (World Scientific, Singapore, 1994).
- [21] G. 't Hooft and M. Veltman, *Nucl. Phys.* **B44** (1972) 189; C. G. Bollini and J. J. Giambiagi, *Nuovo Cim.* **12B** (1972) 20.
- [22] G. Passarino and M. Veltman, *Nucl. Phys.* **B160** (1979) 151.
- [23] J. C. Ward, *Phys. Rev.* **78** (1950) 182; Y. Takahashi, *Nuovo Cim.* **6** (1957) 317; J. C. Taylor, *Nucl. Phys.* **B33** (1971) 436; A. A. Slavnov, *Theor. and Math. Phys.* **10** (1972) 99.

[24] M. E. Peskin and Takeuchi, Phys. Rev. Lett. **65** (1990) 964; G. Altarelli, R. Barbieri, and F. Caravaglio, Nucl. Phys. **B405** (1993) 3.

[25] G. Altarelli, preprint CERN-TH.7464/94.

[26] D. Bardine *et al.*, Z. Phys. **C44** (1989) 493; Comput. Phys. Commun. **59** (1990) 303; Nucl. Phys. **B351** (1991) 1; Phys. Lett. **B255** (1991) 290; CERN-TH 6443/92 (May 1992).

[27] J. Bernabeu, A. Pich, and A. Santamaría, Phys. Lett. **B200** (1988) 569; Nucl. Phys. **B363** (1991) 326.

[28] G. Bhattacharyya, J. Ellis, and K. Sridhar, Phys. Lett. **B336** (1994) 100.

FIGURES




FIG. 1. Feynman diagrams leading to leptoquark contribution to vector-boson self-energies.




FIG. 2. Feynman diagrams leading to leptoquark contribution to the muon decay. The direction of the lines depends upon the leptoquark type.




FIG. 3. Feynman diagrams leading to leptoquark contribution to the vertex Zff . The direction of the lines depends upon the leptoquark type.




Fig. 4.a




Fig. 4.b




Fig. 4.c

FIG. 4. Bounds (95% CL) on the leptoquark masses as a function of m_{top} for S_1 (a), S_3 (b), and R_2 (c). The solid (dashed) lines stand for left-handed (right-handed) leptoquarks, while upper (medium, lower) lines correspond to a coupling $g_{LQ} = \sqrt{4\pi}$ (1, e/s_W).

TABLES

TABLE I. LEP data

Quantity	Experimental value
M_Z [GeV]	91.1888 ± 0.0044
Γ_Z [GeV]	2.4974 ± 0.0038
σ_{had}^0 [nb]	41.49 ± 0.12
$R_e = \frac{\Gamma(had)}{\Gamma(e^+e^-)}$	20.850 ± 0.067
$R_\mu = \frac{\Gamma(had)}{\Gamma(\mu^+\mu^-)}$	20.824 ± 0.059
$R_\tau = \frac{\Gamma(had)}{\Gamma(\mu^+\mu^-)}$	20.749 ± 0.070
A_{FB}^{0e}	0.0156 ± 0.0034
$A_{FB}^{0\mu}$	0.041 ± 0.0021
$A_{FB}^{0\tau}$	0.0228 ± 0.0026
A_τ^0	0.143 ± 0.010
A_c^0	0.135 ± 0.011
$R_b = \frac{\Gamma(bb)}{\Gamma(had)}$	0.2202 ± 0.0020
$R_c = \frac{\Gamma(cc)}{\Gamma(had)}$	0.1583 ± 0.0098
A_{FB}^{0b}	0.0967 ± 0.0038
A_{FB}^{0c}	0.0760 ± 0.0091

TABLE II. Lower limits (95 % CL) for the mass of third generation leptoquarks in GeV for different values of the couplings, assuming $m_{top} = 175$ GeV, $\alpha_s(M_Z^2) = 0.126$, and $M_H = 300$ GeV.

g	S_1^R	S_1^L	S_3	R_2^R	R_2^L	\tilde{S}_1^R	\tilde{R}_2^L
$\sqrt{4\pi}$	4000	5000	6000	4000	5000	280	400
1	750	900	1100	850	1000	—	—
$\frac{e}{s_W}$	350	400	600	400	500	—	—