



IFT

Instituto de Física Teórica
Universidade Estadual Paulista

November/95

IFT-P.060/95
IFUSP/P-1186

syms: 890831

Unitary operator bases and q -deformed algebras

D. Galetti, J. T. Lunardi, B. M. Pimentel

*Instituto de Física Teórica
Universidade Estadual Paulista
Rua Pamplona 145
01405-900 - São Paulo, S.P.
Brazil*

and

C. L. Lima

*Grupo de Física Nuclear Teórica e Fenomenologia de Partículas Elementares
Instituto de Física
Universidade de São Paulo
Caixa Postal 66318
05389-970 - São Paulo, S.P.
Brazil*

**Instituto de Física Teórica
Universidade Estadual Paulista
Rua Pamplona, 145
01405-900 – São Paulo, S.P.
Brazil**

Telephone: 55 (11) 251.5155

Telefax: 55 (11) 288.8224

Telex: 55 (11) 31870 UJMFBR

Electronic Address: LIBRARY@ANP.IFT.UNESP.BR
47857::LIBRARY

Unitary operator bases and q -deformed algebras

D. Galetti, J.T. Lunardi and B.M. Pimentel

Instituto de Física Teórica
Universidade Estadual Paulista
Rua Pamplona 145
01405 - 900 - São Paulo - SP
Brazil

C.L. Lima

Grupo de Física Nuclear Teórica e Fenomenologia de Partículas
Elementares
Instituto de Física
Universidade de São Paulo
Caixa Postal 66318
05389-970 São Paulo - SP
Brazil

Abstract

Starting from the Schwinger unitary operator bases formalism constructed out of a finite dimensional state space, the well-known q -deformed commutation relation is shown to emerge in a natural way, when the deformation parameter is a root of unity.

I. INTRODUCTION

From the studies of deformed algebras, which appeared in connection with problems in statistical mechanics and in quantum field theory (QFT), it came out that the q -deformation

parameter is, in its general form, a complex number. In many applications it assumes a real value while in other cases its imaginary part also plays a physical role. Apart from the basic quantum mechanical study of the q -deformed oscillator by Biedenharn [1] and MacFarlane [2], in which a real deformation parameter is assumed, Floratos [3], on the other hand, in his study of the q -oscillator many-body problem, also discusses the case where q is a pure complex number. Furthermore, in the particular case when q is a root of unity, it can be shown that the underlying state space, characterizing the physical system, is finite dimensional. The q -deformed algebras generate a suitable framework in this case and has been explicitly used in connection with the phase problem in optics [4]; moreover, it has also been pointed out their importance in QFT [5].

A long time ago, Schwinger [6] has pointed out that it is possible to construct an operator basis, in the operator space, once we are given a finite dimensional state space. The two fundamental unitary operators from which the basis is constructed satisfy the Weyl commutation relation and act cyclically on the corresponding state space, thus admitting as many roots of unity, as eigenvalues, as is the dimension of the space. Here we will show how the Schwinger operator basis can be used as a natural tool in order to obtain the q -deformed commutation relation in the particular case when q is a root of unity.

II. THE UNITARY OPERATOR BASES

For the complete quantum description of a physical system, a set of operators must be found in such a way as to permit the construction of all possible dynamical quantities related to that system. The elements of that set are then identified as the elements of a complete operator basis.

One particular set, consisting of unitary operators, has been studied by Schwinger [6] and will be briefly recalled here. Let us consider a N - dimensional linear, normed space of states to be understood as the quantum phase-space of the relevant system. We can define a unitary operator V through the mapping of an orthonormal system $\{ |u_k\rangle \}_{k=0, \dots, N-1}$ defined

in this space, onto itself, by a cyclic permutation as

$$\langle u_k | V = \langle u_{k+1} | , k = 0, \dots, N-1, \langle u_N | = \langle u_0 | .$$

A set of linearly independent unitary operators can then be constructed trivially by mere repeated action of V ,

$$\langle u_k | V^s = \langle u_{k+s} |$$

with

$$\langle u_k | V^N = \langle u_k | ,$$

thus implying

$$V^N = \hat{1} , \quad (2.1)$$

where $\hat{1}$ is the unit operator.

The eigenvalues of V obey this same equation and are thus given by the N roots of unity

$$v_k = \omega^k = \exp\left(\frac{2\pi i k}{N}\right).$$

Furthermore, since that unitary operator has N distinct eigenvalues, the corresponding normalized eigenvectors, $\{\langle v_l | \}_{l=0, \dots, N-1}$, provide us with an alternative orthonormal system.

Schwinger has also shown that an operator U exists such that

$$\langle v_k | U = \langle v_{k-1} | ,$$

which is of period N , i.e.,

$$U^N = \hat{1} ,$$

thus implying the same spectrum as V for the eigenvalues:

$$u_k = \omega^k = \exp\left(\frac{2\pi i k}{N}\right).$$

The fundamental point here is that the eigenvectors of U , $\{\langle u_k | \}_{k=0, \dots, N-1}$, can be shown to coincide with the orthonormal set from which the construction started.

From (2.1) we can see that the special operator normalized to unit trace

$$\hat{G}(v_k) = \frac{1}{N} \sum_{j=0}^{N-1} V^j v_k^{-j} \quad (2.2)$$

is such that

$$\langle v_l | \hat{G}(v_k) = \langle v_l | \delta_{l,k} ,$$

where

$$\delta_{l,k} = \frac{1}{N} \sum_{j=0}^{N-1} v_k^{-j} v_l^j$$

plays the role of a Kronecker delta modulo N .

Corresponding to (2.2) we can also define

$$\hat{T}(u_k) = \frac{1}{N} \sum_{j=0}^{N-1} U^{-j} u_k^j \quad (2.3)$$

with additional equations similar to the above ones.

Using these properties we can show that the two coordinate systems are related by a finite Fourier transformation with coefficients

$$\langle u_k | v_l \rangle = \frac{1}{\sqrt{N}} \exp\left(\frac{2\pi i k l}{N}\right).$$

Now, a simple verification leads us to the relation

$$V^l U^k = \exp\left(\frac{2\pi i k l}{N}\right) U^k V^l , \quad (2.4)$$

which, together with $V^N = \hat{1}$ and $U^N = \hat{1}$, fulfill the conditions which characterize a generalized Clifford algebra [7-10]. Here, however, we will concentrate on just one special feature exhibited by such a set of operators, viz., that the set of N^2 operators,

$$\hat{S}_1(m, n) = \frac{U^m V^n}{\sqrt{N}} , m, n = 0, 1, \dots, N-1,$$

constitutes a complete orthonormal operators basis, with which we can construct all possible dynamical quantities pertaining to that system [6]. In this way, an operator decomposition in this basis is written as

$$\hat{O} = \sum_{m,n=0}^{N-1} O(m,n) \hat{S}_1(m,n), \quad (2.5)$$

where

$$O(m,n) = \text{Tr} [\hat{S}_1^\dagger(m,n) \hat{O}].$$

A very interesting property manifested by the operator basis $\{\hat{S}_1\}$ is the factorization property

$$\hat{S}_1(m,n) = \prod_{l=1}^h \hat{S}_{1l}(m_l, n_l),$$

where the sub-bases

$$\hat{S}_{1l}(m_l, n_l) = \frac{U_l^{m_l} V_l^{n_l}}{\sqrt{P_l}}, \quad m_l, n_l = 0, 1, \dots, P_l - 1,$$

obey the commutation relations

$$V_{l_1} U_{l_2} = U_{l_2} V_{l_1}, \quad l_1 \neq l_2,$$

$$V_{l_1} U_{l_2} = \exp\left(\frac{2\pi i}{P_{l_1}}\right) U_{l_2} V_{l_1}, \quad l_1 = l_2,$$

where h is the total number of primes factors in N including repetitions, with P_l a prime factor of N . This decomposition shows that the factorized basis is constructed from operator sub-bases, each of which associated with a prime number of states, the pair of operators U and V of each sub-basis being classified by the value of the prime integer $P_l = 2, 3, 5, \dots$. It is straightforward to verify that the pair U and V associated with the canonical coordinate-momentum pair $q - p$ is obtained in the particular case $P_l = \infty$. Then, according to Schwinger, due to this factorization property and mutual orthogonality, each of these sub-bases is associated to a particular degree of freedom of the physical system.

In order to emphasize the complete symmetry between U and V , we want also to observe that we could have introduced the new form for the operator basis elements

$$\hat{S}_2(m,n) = \frac{U^m V^n}{\sqrt{N}} \exp\left(\frac{i\pi mn}{N}\right) = \frac{V^n U^m}{\sqrt{N}} \exp\left(\frac{-i\pi mn}{N}\right),$$

which preserves all properties already discussed under the substitutions $U \rightarrow V$ and $V \rightarrow U^{-1}$, combined with $m \rightarrow n$ and $n \rightarrow -m$.

For different degrees of freedom we must conveniently choose the range of variation of the state labels in order to correctly treat the system kinematics; for instance, it is important to emphasize again the canonical case, i.e., $P_l = \infty$, for, in such a case, the unitary operators are immediately identified with the well-known shift operators

$$V \rightarrow e^{iq\hat{p}}$$

$$U \rightarrow e^{ip\hat{q}}$$

when one considers the symmetric interval $m, n = -\frac{N-1}{2}, \dots, +\frac{N-1}{2}$, and then takes the $N \rightarrow \infty$ limit by prime numbers [6]. However, it is also possible to perform a construction of the unitary operators U and V in such a way to obtain an explicit "angle - action" pair, characterizing an Abelian two-dimensional rotation; in this case it can be shown that

$$V \rightarrow \exp\left(i\frac{2\pi}{N}\hat{j}\right) \quad (2.6)$$

$$U \rightarrow \exp(i\hat{\theta}). \quad (2.7)$$

Here, the interval of variation of the state labels are suitably defined to be $m = -\frac{N-1}{2}, \dots, +\frac{N-1}{2}$ and $n = -\frac{N-1}{2}\pi, \dots, +\frac{N-1}{2}\pi$ in such a form that, in the limit of $N \rightarrow \infty$, one recovers $m = \{-\infty, \dots, +\infty\}$, running by integers, and $n = \{-\pi, \pi\}$ [11].

For the sake of completeness it is important to go back to the operator decomposition procedure, Eq.(2.5), and discuss the importance of the particular choice of the operator basis. In fact, in order to emphasize the discrete phase space character of the description,

it was shown that [11], the Fourier transform of the original Schwinger basis $\hat{S}_2(m, n)$ must be considered so that a discrete Weyl transform can be directly identified out of the decomposition scheme. With this new basis it is straightforward to recover the well-known Weyl-Wigner transformation for the canonical continuous case as well as a transformation for an angle - angular momentum degree of freedom as special cases of the $N \rightarrow \infty$ limiting procedure. Furthermore, since the Schwinger basis $\hat{S}_2(m, n)$ is not invariant under a modulo N operation when the state labels are unrestricted in their domain, it was shown that a new operator basis could be devised in order to preserve this symmetry, namely [12]

$$\hat{G}(m, n) = \sum_{j=0}^{N-1} \sum_{l=0}^{N-1} \frac{\hat{T}(j, l)}{\sqrt{N}} \exp \left[-\frac{2\pi i}{N} (mj + nl) \right],$$

where

$$\hat{T}(j, l) = \hat{S}_2(j, l) \exp \{i\pi\phi(j, l; N)\}.$$

The phase $\phi(j, l; N)$ guarantees the mod N invariance.

III. Q-DEFORMED ALGEBRAS

Since the Schwinger unitary operator bases formalism is constructed out of a finite-dimensional state space, the relabelling procedure defines the unitary shifting operators, which have as many eigenvalues (roots of unit) as is the dimension of the underlying state space, N .

Let us now consider the set of eigenstates of the unitary operator V . (Based on the symmetry stated in the last section, this choice is not essential for what follows and could be replaced by the set of eigenstates of the unitary operator U as well.) Since $\{|v_k\rangle\}_{k=0, \dots, N-1}$ is finite-dimensional and the unitary operator U shifts cyclically the states of this space, one cannot interpret U and V as the corresponding creation and annihilation operators. In fact, in the space of eigenstates of V the original pair of unitary operators are represented as

$$V = \sum_{l=0}^{N-1} \exp \left(\frac{2\pi i l}{N} \right) |v_l\rangle\langle v_l| = \sum_{l=0}^{N-1} v_l |v_l\rangle\langle v_l|$$

$$= \sum_{l=0}^{N-1} \omega^l |v_l\rangle\langle v_l| \quad (3.1)$$

$$U = \sum_{l=0}^{N-1} |v_{l+1}\rangle\langle v_l| \quad (3.2)$$

Nevertheless, starting from the unitary operators and making a convenient choice for the state label range, we can construct a pair of operators which will play the role of creation and annihilation operators in this finite-dimensional space.

To begin with, it is immediate to see that, due to the symmetry of the circle embodied in the unitary operator definition, one is not able to fix a vacuum state solely from kinematical considerations, i.e., their action does not select "a priori" any particular state as a vacuum state, since the unitary operators act cyclically in the state space. In this case, one must adopt some criterion to characterize this particular state. This choice will break the symmetry of the circle and is not related to the kinematical content of the description of the physical system. More precisely, one must construct an operator out of the unitary operators in such a form to annihilate the vacuum state; in addition, we must also have a creation operator which generates the "excited" states of the multiplet.

The general form of the creation and annihilation operators will reveal the possibility of particular choices for underlying algebras. To accomplish the construction, we draw our attention again to the relations (3.1) and (3.2). By comparison we can write these operators as

$$a = \sum_{k=0}^{N-1} g(k) |v_k\rangle\langle v_{k+1}|$$

and

$$a^\dagger = \sum_{k=0}^{N-1} |v_{k+1}\rangle\langle v_k|.$$

The form of the creation operator only states that one jumps from the vacuum state up to the last $(N-1)$ state and so on cyclically. In what refers to the annihilation operator

we see that a particularly suitable choice for the unknown function $g(k)$ must use an anti-symmetric function of the state label k so as to select the vacuum state. Now, the natural antisymmetric periodic function defined on the circle is the $\sin(\theta)$ function, what requires odd N 's. Therefore, the proposed annihilation operator is then written as

$$a = \sum_{k=0}^{N-1} \frac{\sin\left(\frac{2\pi k}{N}\right)}{\sin\left(\frac{2\pi}{N}\right)} |v_{k-1}\rangle \langle v_k| .$$

According to the discussion in section 2, we can decompose these operators in the operator basis,

$$\hat{O} = \sum_{m,n=0}^{N-1} O(m,n) \hat{G}(m,n)$$

obtaining

$$a^\dagger = U$$

and

$$a = U^{-1} \frac{V - V^{-1}}{\omega - \omega^{-1}}$$

respectively.

The question that can be posed now is if there exists some relation between the bilinear products of the creation and annihilation operators, a^\dagger , a . Starting from the Weyl relation, Eq. (2.4), the definitions Eqs. (2.6) and (2.7), where instead of \hat{J} we now use \hat{N} , the number operator and using the above expressions for a and a^\dagger , we can immediately obtain the following relation

$$aa^\dagger - \omega a^\dagger a = \omega^{-\hat{N}} .$$

Therefore, we have seen that, starting from the Schwinger unitary operators, the well-known q -deformed commutation relation emerges in a natural way, when the deformation parameter is a root of unity. Furthermore, it is immediate to verify that the creation and annihilation operators, a and a^\dagger proposed here are directly related to the h and g functions proposed by Floratos in his discussion of q -deformed algebras for the bosonic case [3].

The main objective of this paper was to show that the q -deformed algebras can be put in correspondence to Schwinger's unitary operator bases formalism, when the deformation parameter is a root of unity.

Furthermore, it was shown that this formalism is the natural arena for the discussion of recent work on general finite dimensional quantum mechanics problems. Particularly, the Schwinger's formalism was used to represent any operator acting on any finite dimensional state spaces [11,12]. To be specific, it has also been used to study the Liouvillian dynamics in the general finite dimensional phase spaces [13] as well as to describe physical systems from the particular case of a spin 1/2 ($N = 2$ space) up to the canonical continuous case (as the limit $N \rightarrow \infty$). The special case of phase and number operators appearing in connection with quantum optics has also been treated within this framework [14]. This latter problem, or equivalently its Pegg-Barnett description [4], being just a particular case of the general Schwinger formalism, can therefore be also embodied in the q -deformed algebra context along the lines studied here.

Acknowledgment D.G., J.T.L. and B.M.P. were supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico, CNPq, Brazil.

REFERENCES

- [1] L.C. Biedenharn, J. Phys. A: Math. Gen. **22**, (1989) L783.
- [2] A. J. MacFarlane, J. Phys. A: Math. Gen. **22**, (1989) 4581.
- [3] E. G. Floratos, J. Phys. A: Math. Gen. **24**, (1991) 4739.
- [4] S. Abe, Phys. Lett. A **200**, (1995) 239.
- [5] B. Schroer, Rev. Math. Phys. **7**, (1995) 645.
- [6] J. Schwinger, Proc. Nat. Acad. Sci. **46** (1960) 570,893,1401. See also J. Schwinger, *Quantum Kinematics and Dynamics* (Benjamin, New York, 1970).
- [7] K. Morinaga and T. Nono, J. Sci. Hiroshima Univ. Series A14 (No 1) (1952) 13.
- [8] K. Yamazaki, J. Fac. Sci. Tokyo Univ. Sec. I **10** (1964) 147.
- [9] A.O. Morris, Q.J. Math. **17** (1966) 7. **19** (1968) 289.
- [10] A. Ramakrishnan, *L-Matrix Theory or the Grammar of Dirac Matrices* (Tata McGraw-Hill, Bombay, New Delhi, 1972).
- [11] D. Galetti and A.F.R. Toledo Piza, Physica **149A** (1988) 267.
- [12] D. Galetti and A.F.R. Toledo Piza, Physica **A186** (1992) 513.
- [13] D. Galetti and A.F.R. Toledo Piza, Physica **A214** (1995) 207.
- [14] D. Galetti and M.A. Marchioli, Preprint IFT-P.029/95, submitted for publication.