

**UNIVERSIDADE DE SÃO PAULO**

**INSTITUTO DE FÍSICA  
CAIXA POSTAL 66318  
05389-970 SÃO PAULO - SP  
BRASIL**

# **PUBLICAÇÕES**

**IFUSP/P-1201**

**THE ASYMPTOTIC BEHAVIOR OF SINGLE GRADED  
RESPONSE NEURON MODEL WITH A DELAYED SELF-  
CONNECTION**

**C.P. Malta**

Instituto de Física, Universidade de São Paulo

**K. Pakdaman and J.-F. Vibert**

B3E, INSERM U 444, ISARS  
Falcuté de Médecine Saint-Antoine 27,  
rue Chaligny 75571 Paris Cedex 12, France

**O. Arino**

Laboratoire de Mathématiques Appliquées  
Université de Pau et des Pays de l'Adour  
IPRA, URA CNRS 1204  
Avenue de l'Université, 64000 Pau, France

**C. Grotta-Ragazzo**

Instituto de Matemática e Estatística  
Universidade de São Paulo  
CP 66281, 05389-870, São Paulo, BRASIL

Fevereiro/96

## Research announcement:

### The asymptotic behavior of single graded response neuron model with a delayed self-connection

K. Pakdaman<sup>1</sup>, O. Arino<sup>2</sup>, C.P. Malta<sup>3</sup>, C. Grotta-Ragazzo<sup>4</sup>, and J.-F. Vibert<sup>1</sup>

1) B3E, INSERM U 444, ISARS  
Faculté de Médecine Saint-Antoine  
27, rue Chaligny, 75571 Paris Cedex 12 FRANCE

2) Laboratoire de Mathématiques Appliquées  
Université de Pau et des Pays de l'Adour  
IPRA, URA CNRS 1204  
Avenue de l'Université, 64000 Pau, FRANCE

3) Instituto de Física, Universidade de São Paulo  
CP 66318, 05389-970 São Paulo, BRASIL

4) Instituto de Matemática e Estatística, Universidade de São Paulo  
CP 66281, 05389-970 São Paulo, BRASIL  
Mathematics Department, Princeton University  
Fine Hall, Washington Road  
Princeton, NJ 18540 USA

**Acknowledgment:** KP, CPM, CGR and JFV were partially supported by USP-COFEUCUB under project U/C 9/94. CPM was also partially supported by CNPq (the Brazilian Research Council). OA received support from the CNRS PNDR-GLOBAL program.

#### Correspondence should be addressed to:

K. Pakdaman  
B3E, INSERM U444  
Faculté de Médecine Saint-Antoine  
27, rue Chaligny  
75571 Paris Cedex 12, FRANCE  
tel: 33-1-44738430  
fax: 33-1-44738462  
email: pakdaman@b3e.jussieu.fr

## 1 The graded response neuron model

The graded response neuron activation evolves according to the following delay differential equation (DDE):

$$\frac{dx}{dt}(t) = \lambda(-x(t) + \eta\sigma(x(t-1))) \quad (1)$$

Where  $\sigma$  is defined by:

$$\sigma(x) = \tanh(\alpha x) \quad (2)$$

We suppose  $\lambda > 0$ ,  $\eta = \pm 1$  and  $\alpha > 1$ .

Let  $S = \mathcal{C}[-1, 0]$  be the space of continuous real functions of the interval  $[-1, 0]$ . For  $\phi$  in  $S$ , there exists a unique real function  $x(t, \phi)$  on the interval  $[-1, +\infty)$ , such that  $x(t, \phi) = \phi(t)$  for  $-1 \leq t \leq 0$ , and  $x(t, \phi)$  satisfies equation (1) for  $t \geq 0$ . For such a solution of the DDE, we denote by  $x_t(\phi)$  the element of  $S$  defined by  $x_t(\phi)(\theta) = x(t + \theta, \phi)$ , for  $-1 \leq \theta \leq 0$ .

Our research project aims to establish the following description of the global attractor of the DDE (1).

## 2 Negative feedback

In this section we suppose  $\eta = -1$ . Let  $\lambda_0 = 0$ , and  $\lambda_k$  be the value of the parameter  $\lambda$  at the  $k$ th Hopf bifurcation occurring at 0. For this model, all Hopf bifurcations are supercritical.

The discrete time Lyapunov function counts the number of zeros of a solution on an interval of unit length of the form  $[t-1, t]$  such that  $x(t) = 0$  (Mallet-Paret, 1988). Let us denote this function by  $V$ .

We define the Morse sets as follows:  $S_{2N+1} = \{\phi : V(\phi) = 2N + 1\}$  where  $\phi$  is taken in the attractor.

Let  $N^*$  be the number of eigenvalues of the characteristic equation at 0 with positive real parts.  $N^*$  is necessarily even. We consider only the cases where there are no eigenvalues with zero real part.  $S_{N^*} = \{0\}$ , for other even values of  $N$ ,  $S_N$  is empty.

Then the family  $S_n$  forms a Morse decomposition of the attractor of the oscillating solutions.

For the parameter  $\lambda$  in the range:  $\lambda_k < \lambda < \lambda_{k+1}$ , we have  $N^* = 2k$  and

1.  $S_n = \emptyset$  for  $n > N^*$
2.  $S_{2p+1}$  for  $0 \leq p \leq k-1$  is composed of a unique limit periodic solution  $C_p$ .

Moreover, the stable and the unstable manifolds of the various limit cycles intersect transversally. These are the connecting sets between the Morse sets. There are connecting orbits between  $S_m$  and  $S_q$  for all  $m > q$  such that the corresponding Morse sets are non empty.

This gives the Morse decomposition of the attractor. To this decomposition, there corresponds a partition of the phase space  $S$ .

There exists a strictly ordered sequence of subsets of  $S$  such that:

1.  $W_k \subset W_{k-1} \subset \dots \subset W_0 = S$
2.  $W_p$  is of codimension  $2p$
3.  $x_t(\phi)$  tends to  $C_p$  if and only if  $\phi$  is in  $W_p - W_{p+1}$  (with  $0 \leq p \leq k-1$ ).
4.  $x_t(\phi)$  tends to 0 if and only if  $\phi$  is in  $W_k$ .

### 3 Positive feedback

In this section we suppose  $\eta = +1$ . Let  $\lambda_0 = 0$ , and  $\lambda_k$  be the value of the parameter  $\lambda$  at the  $k$ th Hopf bifurcation occurring at 0. For this model, all Hopf bifurcations are supercritical.

The discrete time Lyapunov function counts the number of zeros of a solution on an interval of unit length of the form  $[t-1, t]$  such that  $x(t) = 0$ . Let us denote this function by  $V$  (Arino, 1993).

We define the Morse sets as follows:  $S_{2N} = \{\phi : V(\phi) = 2N\}$  where  $\phi$  is taken in the attractor of the oscillating solutions.

Let  $N^*$  be the number of eigenvalues of the characteristic equation at 0 with positive real parts.  $N^*$  is necessarily odd. We consider only the cases where there are no eigenvalues with zero real part.  $S_{N^*} = \{0\}$ , for other odd values of  $N$ ,  $S_N$  is empty.

Then the family  $S_n$  forms a Morse decomposition of the attractor of the oscillating solutions.

For the parameter  $\lambda$  in the range:  $\lambda_k < \lambda < \lambda_{k+1}$ , we have  $N^* = 2k+1$  and

1.  $S_n = \emptyset$  for  $n > N^*$
2.  $S_{2p}$  for  $1 \leq p \leq k$  is composed of a unique periodic solution  $C_p$ .

Moreover, the stable and the unstable manifolds of the various limit cycles intersect transversally. These are the connecting sets between the Morse sets. There are connecting orbits between  $S_m$  and  $S_q$  for all  $m > q$  such that the corresponding Morse sets are non empty.

This gives the Morse decomposition of the attractor. To this decomposition, there corresponds a partition of the manifold  $W$  of oscillating solutions.

There exists a strictly ordered sequence of subsets of  $W$  such that:

1.  $W_k \subset W_{k-1} \subset \dots \subset W_0 = W$
2.  $W_p$  is of codimension  $2p+1$
3.  $x_t(\phi)$  tends to  $C_p$  if and only if  $\phi$  is in  $W_{p-1} - W_p$  (with  $1 \leq p \leq k$ ).
4.  $x_t(\phi)$  tends to 0 if and only if  $\phi$  is in  $W_k$ .

### 4 References

- Arino, O. (1993) *J. Diff. Equ.*, **104**: 169.  
Mallet-Paret J. (1988), *J. Diff. Equ.*, **72**: 270.