

**UNIVERSIDADE DE SÃO PAULO**

**INSTITUTO DE FÍSICA  
CAIXA POSTAL 66318  
05389-970 SÃO PAULO - SP  
BRASIL**

# **PUBLICAÇÕES**

**IFUSP/P-1204**

**PSEUDOCCLASSICAL DESCRIPTION OF THE MASSIVE  
DIRAC PARTICLES IN ODD DIMENSIONS**

**D.M. Gitman**

Instituto de Física  
Universidade de São Paulo

**A.E. Gonçalves**

Departamento de Física,  
Universidade Estadual de Londrina  
P.O. Box 6001, 86051-970 Londrina, Pr, Brazil

Março/1996

## dimensions

D.M. Gitman

*Instituto de Física, Universidade de São Paulo*

*P.O. Box 66318, 05389-970 São Paulo, SP, Brazil*

A.E. Gonçalves

*Departamento de Física, Universidade Estadual de Londrina*

*P.O. Box 6001, 86051-970 Londrina, Pr, Brazil*

(March 7, 1996)

### Abstract

A pseudoclassical model is proposed to describe massive Dirac (spin one-half) particles in arbitrary odd dimensions. The quantization of the model reproduces the minimal quantum theory of spinning particles in such dimensions. A dimensional duality between the model proposed and the pseudoclassical description of Weyl particles in even dimensions is discussed.

11.10.Ef, 03.65.Pm

## I. INTRODUCTION

As it is known one can construct a pseudoclassical model to describe massive Dirac (spin one-half) particles in  $3 + 1$  dimensions [1]. Its generalization to the case of even dimensions  $D = 2n$ ,  $n = 3, 4, \dots$ , can be done by means of the direct dimensional extension [2]. The corresponding action has the form

$$S = \int_0^1 \left[ -\frac{\dot{x}^2}{2e} - e\frac{m^2}{2} + i \left( \frac{\dot{x}_\mu \psi^\mu}{e} - m\psi^D \right) \chi - i\psi_a \dot{\psi}^a \right] d\tau, \quad (1)$$

where  $\dot{x}^2 = \dot{x}_\mu \dot{x}^\mu$ ; the Greek (Lorentz) indices  $\mu, \nu, \dots$ , run over  $0, 1, \dots, d-1$ , whereas the Latin ones  $a, b$ , run over  $0, 1, \dots, d$ ;  $\eta_{\mu\nu} = \text{diag}(\underbrace{1, -1, \dots, -1}_D)$ ,  $\eta_{ab} = \text{diag}(\underbrace{1, -1, \dots, -1}_{D+1})$ . The variables  $x^\mu$  and  $e$  are even and  $\psi^a, \chi$  are odd. The quantization of the model leads to the Dirac quantum theory of spin one-half particle (to the Dirac equation).

Attempts to extend the pseudoclassical description to the arbitrary odd-dimensional case had met some problems, which are connected with the absence of an analog of  $\gamma^5$ -matrix in such dimensions. For instance, in  $2n + 1$  dimensions the direct generalization of the standard action [1] does not reproduce a minimal quantum theory of spinning particle, where particles with spin  $1/2$  and  $-1/2$  have to be considered as different ones. In the papers [3] two modifications of the standard action were proposed to solve the problem. From our point of view both have essential shortcomings to believe the problem is closed. For instance, the first action [3] is classically equivalent to the standard action and does not provide required quantum properties in course of canonical and path-integral quantization. Moreover, it is  $P$ - and  $T$ -invariant, so that an anomaly is present. Another one [3] does not obey gauge supersymmetries and therefore loses the main attractive feature in such kind of models, which allows one to treat them as prototypes of superstrings or some modes in superstring theory. In [4] a new pseudoclassical model for a massive Dirac particle in  $2 + 1$  dimensions was proposed, which obeys all the necessary symmetries, is  $P$ - and  $-T$  non-invariant and reproduces the minimal quantum theory of the Dirac particle in  $2 + 1$  dimensions. It turns out to be possible to generalize this model to arbitrary odd-dimensional case. We present

such a generalization in the present paper. First, we consider the hamiltonization of the theory and its quantization. Then we discuss a remarkable dimensional duality between the model proposed and the pseudoclassical description of massless spinning particles in even dimensions.

## II. PSEUDOCCLASSICAL DESCRIPTION

In odd dimension  $D = 2n + 1$  we propose the following action to describe spinning particles

$$S = \int_0^1 \left[ -\frac{z^2}{2e} - e\frac{m^2}{2} - im\psi^{2n+1}\chi - \frac{s}{2^n}m\kappa - i\psi_a\dot{\psi}^a \right] d\tau$$

$$\equiv \int_0^1 L d\tau, \quad z^\mu = \dot{x}^\mu - i\psi^\mu\chi + \frac{(2i)^n}{(2n)!}\varepsilon^{\mu\rho_1\dots\rho_{2n}}\psi_{\rho_1}\dots\psi_{\rho_{2n}}\kappa; \quad (2)$$

Here a new even variable  $\kappa$  is introduced and  $\varepsilon^{\mu\nu\dots\lambda}$  is Levi-Civita tensor density in  $2n + 1$ -dimensions normalized by  $\varepsilon^{01\dots 2n} = 1$ ,  $s$  is an even constant of the Berezin algebra. We suppose that  $x^\mu$  and  $\psi^\mu$  are Lorentz vectors and  $e, \kappa, \psi^{2n+1}, \chi$  are scalars so that the action (2) is invariant under the restricted Lorentz transformations (but not  $P$ - and  $T$ -invariant). There are three types of gauge transformations, under which the action (2) is invariant: reparametrizations

$$\delta x^\mu = \dot{x}^\mu \xi, \quad \delta e = \frac{d}{d\tau}(\varepsilon\xi), \quad \delta\psi^a = \dot{\psi}^a \xi, \quad \delta\chi = \frac{d}{d\tau}(\chi\xi), \quad \delta\kappa = \frac{d}{d\tau}(\kappa\xi), \quad (3)$$

with an even parameter  $\xi(\tau)$ ; supertransformations

$$\delta x^\mu = i\psi^\mu \epsilon, \quad \delta e = i\chi \epsilon, \quad \delta\psi^\mu = \frac{z^\mu}{2e}\epsilon, \quad \delta\psi^{2n+1} = \frac{m}{2}\epsilon, \quad \delta\chi = \dot{\epsilon}, \quad \delta\kappa = 0, \quad (4)$$

with an odd parameter  $\epsilon(\tau)$ ; and additional supertransformations

$$\delta x^\mu = -\frac{(2i)^n}{(2n)!}\varepsilon^{\mu\rho_1\dots\rho_{2n}}\psi_{\rho_1}\dots\psi_{\rho_{2n}}\theta,$$

$$\delta\psi^\mu = -\frac{i(2i)^n}{e(2n)!}\varepsilon^{\mu\rho_1\dots\rho_{2n}}z_{\rho_1}\psi_{\rho_2}\dots\psi_{\rho_{2n}}\theta,$$

$$\delta\kappa = \dot{\theta} + \frac{s}{m}\frac{2^{2n+1}i^n(n-1)z_\mu}{(2n)!e}\varepsilon^{\mu\rho_1\dots\rho_{2n}}\dot{\psi}_{\rho_1}\psi_{\rho_2}\dots\psi_{\rho_{2n}}\theta,$$

$$\delta e = \delta\psi^{2n+1} = \delta\chi = 0, \quad (5)$$

with an even parameter  $\theta(\tau)$ .

The total angular momentum tensor  $M_{\mu\nu}$ , is

$$M_{\mu\nu} = x_\mu\pi_\nu - x_\nu\pi_\mu + i[\psi_\mu, \psi_\nu], \quad (6)$$

where  $\pi_\nu = \partial L / \partial \dot{x}^\nu$ .

Going over to the Hamiltonian formulation, we introduce the canonical momenta

$$\pi_\mu = \frac{\partial L}{\partial \dot{x}^\mu} = -\frac{1}{e}z_\mu, \quad P_e = \frac{\partial L}{\partial \dot{e}} = 0, \quad P_\chi = \frac{\partial L}{\partial \dot{\chi}} = 0,$$

$$P_\kappa = \frac{\partial L}{\partial \dot{\kappa}} = 0, \quad P_a = \frac{\partial L}{\partial \dot{\psi}^a} = -i\psi_a. \quad (7)$$

It follows from (7) that there exist primary constraints

$$\Phi_1^{(1)} = P_e, \quad \Phi_2^{(1)} = P_\chi, \quad \Phi_3^{(1)} = P_\kappa, \quad \Phi_{4a}^{(1)} = P_a + i\psi_a. \quad (8)$$

Constructing the total Hamiltonian  $H^{(1)}$ , according to the standard procedure [5,6], we get  $H^{(1)} = H + \lambda_A \Phi_A^{(1)}$ , where

$$H = -\frac{e}{2}(\pi^2 - m^2) + i(\pi_\mu\psi^\mu + m\psi^{2n+1})\chi$$

$$- \left[ \frac{(2i)^n}{(2n)!}\varepsilon^{\mu\rho_1\dots\rho_{2n}}\pi_\mu\psi_{\rho_1}\dots\psi_{\rho_{2n}} - \frac{1}{2^n}sm \right] \kappa. \quad (9)$$

From the consistency conditions  $\dot{\Phi}^{(1)} = \{\Phi^{(1)}, H^{(1)}\} = 0$  we find secondary constraints  $\Phi^{(2)} = 0$ ,

$$\Phi_1^{(2)} = \pi_\mu\psi^\mu + m\psi^{2n+1}, \quad \Phi_2^{(2)} = \pi^2 - m^2,$$

$$\Phi_3^{(2)} = \frac{(2i)^n}{(2n)!}\varepsilon^{\mu\rho_1\dots\rho_{2n}}\pi_\mu\psi_{\rho_1}\dots\psi_{\rho_{2n}} - \frac{1}{2^n}sm, \quad (10)$$

and determine  $\lambda$ , which correspond to the primary constraints  $\Phi_4^{(1)}$ . No more secondary constraints arise from the consistency conditions and the Lagrangian multipliers, correspondent to the primary constraints  $\Phi_i^{(1)}$ ,  $i = 1, 2, 3$ , remain undetermined. The Hamiltonian (9) is proportional to the constraints as one could expect in the case of a reparametrization invariant theory. One can go over from the initial set of constraints  $\Phi^{(1)}, \Phi^{(2)}$  to the equivalent

ones  $\Phi^{(1)}, \tilde{\Phi}^{(2)}$ , where  $\tilde{\Phi}^{(2)} = \Phi^{(2)} (\psi \rightarrow \tilde{\psi} = \psi + \frac{1}{2}\Phi_4^{(1)})$ . The new set of constraints can be explicitly divided in a set of the first-class constraints, which are  $(\Phi_i^{(1)}, i = 1, 2, 3, \tilde{\Phi}^{(2)})$  and in a set of second-class constraints  $\Phi_4^{(1)}$ . Thus, we are dealing with a theory with first-class constraints.

### III. QUANTIZATION

Let us consider first the Dirac quantization, where the second-class constraints define the Dirac brackets and therefore the commutation relations, whereas, the first-class constraints, being applied to the state vectors, define physical states. For essential operators and nonzeroth commutation relations one can obtain in the case of consideration:

$$[\hat{x}^\mu, \hat{\pi}_\nu] = i\{x^\mu, \pi_\nu\}_{D(\Phi_4^{(1)})} = i\delta_\nu^\mu, \quad [\hat{\psi}^a, \hat{\psi}^b]_+ = i\{\psi^a, \psi^b\}_{D(\Phi_4^{(1)})} = -\frac{1}{2}\eta^{ab}. \quad (11)$$

It is possible to construct a realization of the commutation relations (11) in a Hilbert space  $\mathcal{R}$  whose elements  $\mathbf{f} \in \mathcal{R}$  are  $2^{n+1}$  component columns dependent on  $x$ ,

$$\mathbf{f}(x) = \begin{pmatrix} u_-(x) \\ u_+(x) \end{pmatrix}, \quad (12)$$

where  $u_{\mp}(x)$  are  $2^n$  component columns. Then

$$\hat{x}^\mu = x^\mu \mathbf{I}, \quad \hat{\pi}_\mu = -i\partial_\mu \mathbf{I}, \quad \hat{\psi}^a = \frac{i}{2}\gamma^a, \quad (13)$$

here  $\mathbf{I}$  is  $2^{n+1} \times 2^{n+1}$  unit matrix and  $\gamma^a$ ,  $a = 0, 1, \dots, 2n+1$  are  $\gamma$ -matrices in  $2(n+1)$ -dimensions [7], which we select in the spinor representation  $\gamma^0 = \text{antidiag}(I, I)$ ,  $\gamma^i = \text{antidiag}(\sigma^i, -\sigma^i)$ ,  $i = 1, 2, \dots, 2n+1$ , where  $I$  is  $2^n \times 2^n$  unit matrix, and  $\sigma^i$  are  $2^n \times 2^n$   $\sigma$ -matrix, which obey the Clifford algebra,  $[\sigma^i, \sigma^j]_+ = 2\delta^{ij}$ .

According to the scheme of quantization selected, the operators of the first-class constraints have to be applied to the state vectors to define physical sector, namely,  $\hat{\Phi}^{(2)}\mathbf{f}(x) = 0$ , where  $\hat{\Phi}^{(2)}$  are operators, which correspond to the constraints (10). There is no ambiguity in the construction of the operator  $\hat{\Phi}_1^{(2)}$  according to the classical function  $\Phi_1^{(2)}$ . Taken

into account the realization (12), (13), one can present the equations  $\hat{\Phi}^{(2)}\mathbf{f}(x) = 0$  in the  $2^n$ -component form,

$$[i\partial_\mu \gamma^\mu - m\gamma^{2n+1}]\mathbf{f}(x) = 0 \iff \begin{cases} [i\partial_\mu \Gamma_+^\mu - m]u_+(x) = 0, \\ [i\partial_\mu \Gamma_-^\mu + m]u_-(x) = 0, \end{cases} \quad (14)$$

where two sets of  $\gamma$ -matrices  $\Gamma_\zeta^\mu$ ,  $\zeta = \pm$ , in  $2n+1$  dimensions are introduced,

$$\begin{aligned} \Gamma_\zeta^0 &= \sigma^{2n+1}, \quad \Gamma_\zeta^1 = \zeta\sigma^{2n+1}\sigma^1, \dots, \Gamma_\zeta^{2n} = \zeta\sigma^{2n+1}\sigma^{2n}, \\ \Gamma_-^\mu &= \Gamma_{+\mu}, \quad [\Gamma_\zeta^\mu, \Gamma_\zeta^\nu]_+ = 2\eta^{\mu\nu}. \end{aligned} \quad (15)$$

There is a relation  $\hat{\Phi}_2^{(2)} = (\hat{\Phi}_1^{(2)})^2$  so that the equation  $\hat{\Phi}_2^{(2)}\mathbf{f} = 0$  is not independent. The equation  $\hat{\Phi}_3^{(2)}\mathbf{f}(x) = 0$  can be presented in the following form

$$\left[ \frac{(-i)^n}{(2n)!} \epsilon^{\mu\rho_1 \dots \rho_{2n}} (i\partial_\mu) \gamma_{\rho_1} \dots \gamma_{\rho_{2n}} + sm \right] \mathbf{f}(x) = 0$$

or in  $2^n$ -component form

$$\begin{aligned} [i\partial_\mu \Gamma_+^\mu + (-1)^n sm]u_+(x) &= 0, \\ [i\partial_\mu \Gamma_-^\mu + (-1)^n sm]u_-(x) &= 0. \end{aligned} \quad (16)$$

In quantum theory one has to select  $s = \pm 1$ , then, combining eq. (14) and (16), we get

$$[i\partial_\mu \Gamma_\zeta^\mu - \zeta m]u_\zeta(x) = 0, \quad u_{-\zeta}(x) \equiv 0, \quad \zeta = (-1)^n s = \pm 1. \quad (17)$$

To interpret the result obtained one has to calculate also the operators  $\hat{M}_{\mu\nu}$  correspondent to the angular momentum tensor (6),

$$\hat{M}_{\mu\nu} = -i(x_\mu \partial_\nu - x_\nu \partial_\mu) - \frac{i}{4} \begin{pmatrix} [\Gamma_{-\mu}, \Gamma_{-\nu}] & 0 \\ 0 & [\Gamma_{+\mu}, \Gamma_{+\nu}] \end{pmatrix}.$$

Thus, in the quantum mechanics constructed, the states with  $\zeta = +$  are described by the  $2^n$ -component wave function  $u_+(x)$ , which obeys the Dirac equation in  $2n+1$  dimensions and is transformed under the Lorentz transformation as spin  $+1/2$ . For  $\zeta = -$  the quantization leads to the theory of  $2n+1$  Dirac particle with spin  $-1/2$  and the wave function  $u_-(x)$ .

To quantize the theory canonically we have to impose as much as possible supplementary gauge conditions to the first-class constraints. In the case under consideration, it turns out to be possible to impose gauge conditions to all the first-class constraints, excluding the constraint  $\tilde{\Phi}_3^{(2)}$ . Thus, we are fixing the gauge freedom, which corresponds to two types of gauge transformations (3) and (4). As a result we remain only with one first-class constraint, which is a reduction of  $\tilde{\Phi}_3^{(2)}$  to the rest of constraints and gauge conditions. It can be used to specify the physical states. All the second-class constraints form the Dirac brackets. The following gauge conditions  $\Phi^G = 0$  are imposed:  $\Phi_1^G = e + \zeta\pi_0^{-1}$ ,  $\Phi_2^G = \chi$ ,  $\Phi_3^G = \kappa$ ,  $\Phi_4^G = x_0 - \zeta\tau$ ,  $\Phi_5^G = \psi^0$ , where  $\zeta = -\text{sign}\pi^0$ . (The gauge  $x_0 - \zeta\tau = 0$  was first proposed in [6,8] as a conjugated gauge condition to the constraint  $\pi^2 - m^2 = 0$ ). Using the consistency condition  $\dot{\Phi}^G = 0$ , one can determine the Lagrangian multipliers, which correspond to the primary constraints  $\Phi_i^{(1)}$ ,  $i = 1, 2, 3$ . To go over to a time-independent set of constraints (to use standard scheme of quantization without any modifications [6]) we introduce the variable  $x'_0$ ,  $x'_0 = x_0 - \zeta\tau$ , instead of  $x_0$ , without changing the rest of the variables. That is a canonical transformation in the space of all variables with the generating function  $W = x_0\pi'_0 + \tau|\pi'_0| + W_0$ , where  $W_0$  is the generating function of the identity transformation with respect to all variables except  $x^0$  and  $\pi_0$ . The transformed Hamiltonian  $H^{(1)'$  is of the form

$$H^{(1)'} = H^{(1)} + \frac{\partial W}{\partial \tau} = \omega + \{\Phi\}, \quad \omega = \sqrt{\pi_d^2 + m^2}, \quad d = 1, 2, \dots, 2n, \quad (18)$$

where  $\{\Phi\}$  are terms proportional to the constraints and  $\omega$  is the physical Hamiltonian. All the constraints of the theory, can be presented after this canonical transformation in the following equivalent form:  $K = 0$ ,  $\phi = 0$ ,  $T = 0$ , where

$$\begin{aligned} K &= (e - \omega^{-1}, P_e; \chi, P_\chi; \kappa, P_\kappa; x'_0, |\pi_0| - \omega; \psi^0, P_0); \\ \phi &= (\pi_d\psi^d + m\psi^{2n+1}, P_k + \nu\psi_k), \quad k = 1, 2, \dots, 2n+1; \\ T &= \frac{(2i)^n}{(2n)!} \zeta\omega\varepsilon^{i_1\dots i_{2n}}\psi_{i_1}\dots\psi_{i_{2n}} + \frac{sm}{2^n}, \quad i_d = 1, 2, \dots, 2n. \end{aligned} \quad (19)$$

The constraints  $K$  and  $\phi$  are of the second-class, whereas  $T$  is the first-class constraint.

Besides, the set  $K$  has the so called special form [6]. In this case, if we eliminate the variables  $e$ ,  $P_e$ ,  $\chi$ ,  $P_\chi$ ,  $\kappa$ ,  $P_\kappa$ ,  $x'_0$ ,  $|\pi_0|$ ,  $\psi^0$ , and  $P_0$ , using the constraints  $K = 0$ , the Dirac brackets with respect to all the second-class constraints ( $K, \phi$ ) reduce to ones with respect to the constraints  $\phi$  only. Thus, on this stage, we will only consider the variables  $x^d$ ,  $\pi_d$ ,  $\zeta$ ,  $\psi^k$ ,  $P_k$  and two sets of constraints - the second-class ones  $\phi$  and the first-class one  $T$ . Nonzero Dirac brackets for the independent variables are

$$\begin{aligned} \{x^d, \pi_r\}_{D(\phi)} &= \delta_r^d, \quad \{x^d, x^r\}_{D(\phi)} = \frac{i}{\omega^2}[\psi^d, \psi^r], \quad \{x^d, \psi^r\}_{D(\phi)} = -\frac{1}{\omega^2}\psi^d\pi_r, \\ \{\psi^d, \psi^r\}_{D(\phi)} &= -\frac{i}{2}(\delta_r^d - \omega^{-2}\pi_d\pi_r), \quad d, r = 1, 2, \dots, 2n. \end{aligned} \quad (20)$$

Going over to the quantum theory, we get the commutation relations between the operators  $\hat{x}^d$ ,  $\hat{\pi}_d$ ,  $\hat{\psi}^d$  by means of the Dirac brackets (20),

$$\begin{aligned} [\hat{x}^d, \hat{\pi}_r] &= i\delta_r^d, \quad [\hat{x}^d, \hat{x}^r] = -\frac{1}{\hat{\omega}^2}[\hat{\psi}^d, \hat{\psi}^r], \\ [\hat{x}^d, \hat{\psi}^r] &= -\frac{i}{\hat{\omega}^2}\hat{\psi}^d\hat{\pi}_r, \quad [\hat{\psi}^d, \hat{\psi}^r]_+ = \frac{1}{2}(\delta_r^d - \hat{\omega}^{-2}\hat{\pi}_d\hat{\pi}_r). \end{aligned} \quad (21)$$

We assume as usual [6,8] the operator  $\hat{\zeta}$  to have the eigenvalues  $\zeta = \pm 1$  by analogy with the classical theory, so that  $\hat{\zeta}^2 = 1$ , and also we assume the equations of the second-class constraints  $\hat{\phi} = 0$ . Then one can realize the algebra (21) in a Hilbert space  $\mathcal{R}$ , whose elements  $\mathbf{f} \in \mathcal{R}$  are  $2^{n+1}$  component columns dependent on  $\mathbf{x} = (x^d)$ ,  $d = 1, 2, \dots, 2n$ ,

$$\mathbf{f}(\mathbf{x}) = \begin{pmatrix} f_+(\mathbf{x}) \\ f_-(\mathbf{x}) \end{pmatrix}, \quad (22)$$

so that  $f_+(\mathbf{x})$  and  $f_-(\mathbf{x})$  are  $2^n$  component columns. A realization of the commutations relations has the form

$$\begin{aligned} \hat{x}^d &= x^d\mathbf{I} - \frac{i}{4\hat{\omega}^2}[\Sigma^d, \hat{\pi}_r\Sigma^r]_- - \frac{im}{4\hat{\omega}^2}[\Sigma^d, \Sigma^{2n+1}]_-, \quad \hat{\pi}_r = -i\partial_r\mathbf{I}, \\ \hat{\psi}^d &= \frac{1}{2}(\delta_r^d - \hat{\omega}^{-2}\hat{\pi}_d\hat{\pi}_r)\Sigma^r - \frac{m\hat{\pi}_d}{2\hat{\omega}^2}\Sigma^{2n+1}, \quad \hat{\zeta} = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}, \end{aligned} \quad (23)$$

where  $\mathbf{I}$  and  $I$  are  $2^{n+1} \times 2^{n+1}$  and  $2^n \times 2^n$  unit matrices,  $\Sigma^k = \text{diag}(\sigma^k, \sigma^k)$ . The operator  $\hat{T}$  correspondent to the first-class constraint  $T$  (see (19)) appears to be

$$\hat{T} = \frac{\zeta m}{\hat{\omega}} \hat{\zeta} \Sigma^{2n+1} \left[ \hat{\zeta} \hat{\omega} \Sigma^{2n+1} + i \partial_d (\zeta \Sigma^{2n+1} \Sigma^d) - \zeta m \right], \quad \zeta = (-1)^n s = \pm 1. \quad (24)$$

The latter operator specifies the physical states according to scheme of quantization selected,  $\hat{T}\mathbf{f} = 0$ . On the other hand, the state vectors  $\mathbf{f}$  have to obey the Schrödinger equation, which defines their "time" dependence,  $(i\partial/\partial\tau - \hat{\omega})\mathbf{f} = 0$ ,  $\hat{\omega} = \sqrt{\hat{\pi}_d^2 + m^2}$ , where the quantum Hamiltonian  $\hat{\omega}$  corresponds the classical one  $\omega$  (18). Introducing the physical time  $x^0 = \zeta\tau$  instead of the parameter  $\tau$  [8,6], we can rewrite the Schrödinger equation in the following form (we can now write  $\mathbf{f} = \mathbf{f}(x)$ ,  $(x = x^0, \mathbf{x})$ ),

$$\left( i \frac{\partial}{\partial x^0} - \hat{\zeta} \hat{\omega} \right) \mathbf{f}(x) = 0. \quad (25)$$

Using (25) in the eq.  $\hat{T}\mathbf{f} = 0$ , namely replacing there the combination  $\hat{\zeta}\hat{\omega}\mathbf{f}$  by  $i\partial_0\mathbf{f}$ , one can verify that both components  $f_{\pm}(x)$ , of the state vector (22) obey one and the same equation

$$(i\partial_\mu \Gamma_\zeta^\mu - \zeta m) f_\zeta(x) = 0, \quad \zeta = \pm 1, \quad (26)$$

which is the  $2n + 1$  Dirac equation for a particle of spin  $\zeta/2$  whereas  $f_{\pm}(x)$  can be interpreted (taken into account (25)) as positive and negative frequency solutions to the equation respectively. Substituting the realization (23) into the expression (6), we get the generators of the Lorentz transformations

$$\hat{M}_{\mu\nu} = -i(x_\mu \partial_\nu - x_\nu \partial_\mu) - \frac{i}{4} \begin{pmatrix} [\Gamma_{\zeta\mu}, \Gamma_{\zeta\nu}] & 0 \\ 0 & [\Gamma_{\zeta\mu}, \Gamma_{\zeta\nu}] \end{pmatrix}, \quad (27)$$

which have the standard form for both components  $f_\zeta(x)$ . Thus, a natural interpretation of the components  $f_\zeta(x)$  is the following:  $f_+(x)$  is the wave function of a particle with spin  $\zeta/2$  and  $f_-(x)$  is the wave function of an anti-particle with spin  $\zeta/2$ .

#### IV. DIMENSIONAL DUALITY BETWEEN MASSIVE AND MASSLESS SPINNING PARTICLES

As is known, the method of dimensional reduction [9] appears to be often useful to construct models (actions) in low dimensions using some appropriate models in higher dimensions. In fact, such kind of ideas began from the works [10]. One can also mention that

the method of dimensional reduction was used to interpret masses in supersymmetric theories as components of momenta in space of higher dimensions, which are frozen in course of the reduction. It is interesting that the model of Dirac particles in odd dimensions proposed in the present paper is related to the model [11] of Weyl particles in even dimensions by means of a dimensional reduction.

The action and the Hamiltonian of the latter model in  $D = 2(n + 1)$  dimensions have the form

$$\begin{aligned} S &= \int_0^1 \left[ -\frac{z^2}{2e} - i\psi_\mu \dot{\psi}^\mu \right] d\tau, \\ z^\mu &= \dot{x}^\mu - i\psi^\mu \chi + \frac{(2i)^{\frac{D-2}{2}}}{(D-2)!} \epsilon^{\mu\nu\rho_2 \dots \rho_{D-1}} b_\nu \psi_{\rho_2} \dots \psi_{\rho_{D-1}} + \frac{s}{2^{\frac{D-2}{2}}} b^\mu, \\ H &= -\frac{e}{2} \pi^2 + i\pi_\mu \psi^\mu \chi \\ &\quad - \left[ \frac{(2i)^{\frac{D-2}{2}}}{(D-2)!} \epsilon_{\nu\mu\rho_2 \dots \rho_{D-1}} \pi^\mu \psi^{\rho_2} \dots \psi^{\rho_{D-1}} + \frac{\alpha}{2^{\frac{D-2}{2}}} \pi_\nu \right] b^\nu. \end{aligned} \quad (28)$$

In the canonical gauge similar to one which was considered above, in particular, in the gauge  $\psi^0 = 0$ , we remain only with the first-class constraints

$$T_\mu = \frac{(2i)^{\frac{D-2}{2}}}{(D-2)!} \epsilon_{\nu\mu\rho_2 \dots \rho_{D-1}} \pi^\mu \psi^{\rho_2} \dots \psi^{\rho_{D-1}} + \frac{\alpha}{2^{\frac{D-2}{2}}} \pi_\nu = 0. \quad (29)$$

One can see that, in fact, among the constraints (29) only one is independent

$$T_\mu = \frac{\pi_\mu}{\pi_0} T_0.$$

Thus one can use only one component of  $b^\mu$  and all others put to be zero. Now one can do a dimensional reduction  $2(n + 1) \rightarrow 2n + 1$  in the Hamiltonian and constraint  $T_0$ , putting also  $\pi_{2n+1} = m$ ,  $b^{2n+1} = -\kappa$ ,  $b^0 = b^1 = \dots = b^{2n} = 0$ . As a result of such a procedure we just obtain the expressions (9) and (19) for the Hamiltonian and the constraint. The second class constraints of the model (28) also coincide with ones of the model (2) after the dimensional reduction. Thus, there exist a dimensional duality between the massive spinning particles in odd dimensions and massless ones in even dimensions.

## V. DISCUSSION

One ought to say that at present there exist pseudoclassical models (PM) to describe all massive higher spins (integer and half-integer) in 3+1 dimensions [1,12]. Generalization of the models to arbitrary even dimensions can be easily done by means of a trivial dimensional extension similarly to the spin one-half case. To get the PM for higher spins in arbitrary odd dimensions one can start from the model proposed in the present paper, using the ideas of the work [13]. Namely, one has to multiply the variables  $\psi$ ,  $\chi$ ,  $\kappa$ ,  $s$  in the action (2). Then an appropriate action has the form

$$S = \int_0^1 \left\{ -\frac{z^2}{2e} - e\frac{m^2}{2} - \sum_{A=1}^N \left[ sm \left( \frac{\kappa_A}{2^n} + i\psi_A^{2n+1} \chi_A \right) + i\psi_{A\alpha} \dot{\psi}_A^\alpha \right] \right\} d\tau, \\ z^\mu = \dot{x}^\mu - \sum_{A=1}^N \left[ i\psi_A^\mu \chi_A - \frac{(2i)^n}{(2n)!} \varepsilon^{\mu\rho_1 \dots \rho_{2n}} \psi_{A\rho_1} \dots \psi_{A\rho_{2n}} \kappa_A \right]. \quad (30)$$

Certainly, a detailed analysis of the action (30) and its quantization may demand essential technical work in higher dimensions. In spite of in 2+1 dimensions the model can be quantized explicitly for all higher spins both canonically and by means of the Dirac method [13], in 3+1 dimensions the corresponding PM [12] was quantized canonically only for spins one-half [1] and one [14]. As to the massless particles spin one-half, the corresponding PM exist at present in arbitrary even dimensions [11,15]. Its generalization to describe higher spins can be done in the same manner

$$S = \int_0^1 \left[ -\frac{1}{2e} z^2 - i \sum_{A=1}^N \psi_{A\mu} \dot{\psi}_A^\mu \right] d\tau, \\ z^\mu = \dot{x}^\mu - \sum_{A=1}^N \left( i\psi_A^\mu \chi_A - i\varepsilon^{\mu\nu\rho\sigma} b_{A\nu} \psi_{A\rho} \psi_{A\sigma} - \frac{1}{2} s b_A^\mu \right). \quad (31)$$

There exist the dimensional duality mentioned above between the models (30) and (31).

Massless higher spins in arbitrary odd dimensions can be described pseudoclassically by the model, which follows from (30) in the limit  $m \rightarrow 0$ .

Thus, at present, in principle, we have PM to describe all integer and half-integer spins in arbitrary dimensions.

## Acknowledgments

D.M. Gitman thanks Brazilian foundations CNPq for support.

## REFERENCES

- [1] F.A. Berezin and M.S. Marinov, JETP Lett. **21**, 320 (1975); Ann. Phys. (N.Y.) **104**, 336 (1977); L. Brink, S. Deser, B. Zumino, P. Di Vecchia, and P. Howe, Phys. Lett. **64B**, 435 (1976); L. Brink, P. Di Vecchia, and P. Howe, Nucl. Phys. **B118**, 76 (1977); R. Casalbuoni, Nuovo Cim. **A33**, 115 (1976); A. Barducci, R. Casalbuoni, and L. Lusanna, Nuovo Cim. **A35**, 377 (1976); A.P. Balachandran, P. Salomonson, B. Skagerstam, and J. Winnberg, Phys. Rev. D **15**, 2308 (1977); M. Henneaux and C. Teitelboim, Ann. Phys. (N.Y.) **143**, 127 (1982); D.M. Gitman and I.V. Tyutin, Class. and Quantum Grav. **7**, 2131 (1990); E.S. Fradkin and D.M. Gitman, Phys. Rev. D **44**, 3230 (1991).
- [2] G.V. Grigoryan and R.P. Grigoryan, Yad. Fiz. **53**, 1737 (1991).
- [3] M.S. Plyushchay, Mod. Phys. Lett. **A8**, 937 (1993); J.L. Cortes, M.S. Plyushchay and L. Velazquez, Phys. Lett. B **306**, 34 (1993).
- [4] D.M. Gitman, A.E. Gonçaves, and I.V. Tyutin, Preprint IFUSP/P - 1144, march/1995, hep-th/9601065.
- [5] P.M.A. Dirac, *Lectures on Quantum Mechanics* (Yeshiva University, 1964).
- [6] D.M. Gitman and I.V. Tyutin, *Quantization of Fields with Constraints* (Springer, Berlin, 1990).
- [7] K.M. Case, Phys. Rev. **97**, 810 (1955).
- [8] D.M. Gitman and I.V. Tyutin, Pis'ma Zh. Eksp. Teor. Fiz. **51**, 188 (1990); Class. and Quantum Grav. **7**, 2131 (1990).
- [9] M.J. Duff, B.E. Nilsson, and C.N. Pope, Phys. Rep. **130**, 1 (1986); M.B. Green, J.H. Schwartz, and E. Witten, *Superstring Theory* V. 1, 2. (Cambridge Univ. Press, 1988).
- [10] Th. Kaluza, Sitzungsber. Preuss. Akad. Wiss. Berlin. Math. Phys. **K1**, 966 (1921); O. Klein, Z. Phys. **37**, 895 (1926).
- [11] D.M. Gitman and A.E. Gonçaves, Preprint IFUSP/P-1158, May/1995, hep-th/9506010; G.V. Grigoryan, R.P. Grigoryan, and I.V. Tyutin, Preprint YERPHY-1446(16)-95, hep-th/9510002.
- [12] P.P. Srivastava, Nuovo Cimento Lett. **19**, 239 (1977); V.D. Gershun and V.I. Tkach, Pis'ma Zh. Eksp. Teor. Fiz. **29**, 320 (1979); P.S. Howe, S. Penati, M. Pernice, and P. Townsend, Phys. Lett. B **215**, 555 (1988); Class. Quantum Grav. **6**, 1125 (1989); A. Barducci, R. Casalbuoni, and L. Lusanna, Nuovo Cimento **A35**, 377 (1976); R. Marnelius and U. Marterson, Nucl. Phys. **B335**, 395, (1990); Int. J. Mod. Phys. **A6**, 807, (1991).
- [13] D.M. Gitman and I.V. Tyutin, Preprint FIAN/TD/96-26, hep-th/9602048
- [14] D.M. Gitman, A.E. Gonçaves, and I.V. Tyutin Int. J. Mod. Phys. A **10**, 701 (1995).
- [15] D.M. Gitman, A.E. Gonçaves, and I.V. Tyutin Phys. Rev. D **50**, 5439 (1994).