

UNIVERSIDADE DE SÃO PAULO

PUBLICAÇÕES

INSTITUTO DE FÍSICA
CAIXA POSTAL 66318
05315-970 SÃO PAULO - SP
BRASIL

IFUSP/P-1268

APPLICATION OF PATH INTEGRATION TO
OPERATOR CALCULUS

Dmitri M. Gitman and **Paulo B. Barros**
Instituto de Física, Universidade de São Paulo

Stoian I. Zlatev
Departamento de Física,
Universidade Federal de Sergipe,
49000-000 Aracaju, SE, Brasil

pag. 1-13

Maio/1997

Application of Path Integration to Operator Calculus

Dmitri M. Gitman*, Stoian I. Zlatev† and Paulo B. Barros*

(May 8, 1997)

Abstract

We consider a disentanglement of the operator functions of the form $\gamma^\alpha \dots \gamma^\beta \exp \{ \omega_{\mu\nu} \gamma^\mu \gamma^\nu \}$ where γ^μ are generating elements of a Clifford algebra (γ matrices, for example). To this end we formulate a path integral reduction procedure which allows one to get the functions under consideration in Sym-form. Then by means of the path integration we get explicit decompositions of the operator functions in Sym-products of γ matrices (in the linearly independent γ -matrix structures) valid in arbitrary dimensions. Several particular examples are analyzed in details.

I. INTRODUCTION

As it is well known, the path integrals are widely and fruitfully applied in the contemporary theoretical physics. For example, they are used to solve Schrödinger equation and equations of diffusion theory, they are well adopted for quasiclassical calculations in quantum mechanics, path integrals are used for quantization of gauge theories and serve as the basic language in the instanton physics, they have found wide application in statistical mechanics, especially when methods of the quantum field theory are used there. The integrals over Grassmann variables introduced by Berezin [1] made it possible to define the corresponding path integrals over Grassmann-odd trajectories. This enlarged even more the field of application of path integrals. In the present paper we would like to stress on a possibility how one can use path integrals over Grassmann-odd trajectories to disentangle complicated functions on noncommuting operators (some rules of dealing with such functions were considered in the works [2-4]). Namely, we are going to consider the operator functions of the form

$$R_k = \underbrace{\gamma^\alpha \dots \gamma^\beta}_k \exp \{ \omega_{\mu\nu} \gamma^\mu \gamma^\nu \}, \quad k < D, \quad (1)$$

where the constant matrix ω is antisymmetric, $\omega_{\mu\nu} = -\omega_{\nu\mu}$, and γ^μ , $\mu = 0, 1, \dots, D-1$, are generating elements of some Clifford algebra,

$$[\gamma^\mu, \gamma^\nu]_+ = 2\eta^{\mu\nu}. \quad (2)$$

The latter can be, in particular, understood as γ -matrices in D dimensions (in this case $\eta_{\mu\nu} = \text{diag}(1, -1, \dots, -1)$). Expressions of the form (1) arise frequently in different theoretical constructions. Here one ought to mention spinor representations of the Lorentz group. It is also known that propagators of relativistic spinning particles and superstrings in external fields, derived by means of the Schwinger proper-time method, contain γ matrices in the form (1). Doing calculations with propagators of that kind, one inevitably comes to the problem of the expansion of such expressions in terms of the independent γ -matrix structures. One has also to mention that modern field theories and superstring theory are usually formulated in

*Instituto de Física, Universidade de São Paulo, Caixa Postal 66318-CEP 05315-970-São Paulo, S.P., Brasil

†Departamento de Física, Universidade Federal de Sergipe, 49000-000 Aracaju, SE, Brasil. On leave from the Institute for Nuclear Research and Nuclear Energy, Sofia, Bulgaria

space-time dimensions different from four. Thus, it is important to analyze the structure of the operator functions (1) for arbitrary dimensions. Besides, commutation relations between the generators Γ_a of a Lie algebra can be realized by bilinear combinations of some Clifford algebra generating elements, similar to Schwinger type representations via creation and annihilation operators [5]. Indeed, let, for example, Γ_a , $a = 1, \dots, n$, be generators of $SU(N)$ group, $[\Gamma_a, \Gamma_b] = if_{abc}\Gamma_c$. Then one can see that the commutation relations of the algebra can be obeyed by means of the following representation: $\Gamma_a = -\frac{i}{4}f_{acd}\gamma_c\gamma_d$, where γ_a are generators of the corresponding Clifford algebra, $[\gamma_a, \gamma_b]_+ = \delta_{ab}$. Then finite transformations of the corresponding Lie group are presented by the operator functions R_0 . Thus, the operator problem under consideration seems to be quite actual by itself. We present a decomposition of the operator functions (1) via symmetrical (Sym) products of γ matrices which constitute linearly independent structures in finite number. To do that we formulate a Grassmann path integral reduction procedure which allows one to get the functions under consideration in Sym-form. Then the problem can be solved by means of a path integration. Thus, we get the explicit γ -matrix structure of the operator functions under consideration in arbitrary dimension. In the end we consider particular cases in lower dimensions ($D = 3, 4$) identifying the corresponding decompositions with some known before formulas derived by means of direct combinatoric methods strongly related to concrete properties of γ matrices in such dimensions. We find it remarkable that the solution of the operator problem is facilitated considerably by using the method of path integration. This extends the list of its useful applications.

II. T AND SYM FORM OF THE OPERATOR FUNCTIONS

Let us consider first a particular case of the operator expression (1), namely, R_0 . Using the famous Feynman's consideration [3], we can present R_0 in the following form

$$R_0 = \mathcal{P} \exp \left\{ \int_0^1 \omega_{\mu\nu} \sigma^{\mu\nu}(t) dt \right\}, \quad (3)$$

where the index t (we will call it time) is formally attached to each matrix $\sigma^{\mu\nu} = \frac{1}{2}[\gamma^\mu, \gamma^\nu]$ and \mathcal{P} means that "the operator with higher time acts later". Under the sign of the chronological products the operators $\sigma^{\mu\nu}(t)$ commute and can be treated as ordinary c -numbers.

One can remark that expressions similar to (3) arise naturally in quantum-mechanical problems with Hamiltonians of the form $\mathcal{H}(t) = i\omega_{\mu\nu}(t)\gamma^\mu\gamma^\nu$. In this case the evolution operator between the instants $t = 0$ and $t = 1$ has the form

$$U = \mathcal{P} \exp \left\{ \int_0^1 \omega_{\mu\nu}(t) \sigma^{\mu\nu}(t) dt \right\}, \quad (4)$$

where the index t is now attached in a natural way to σ matrices.

How to calculate efficiently expression (3)? A convenient way is to use Wick theorem [6] for appropriately defined T products of some operators whose commutators or anticommutators are c -numbers. In the case under consideration γ matrices are such operators with anticommutators (2) being c -numbers. This dictates the choice of the "fermionic" T product for γ matrices,

$$\begin{aligned} T\gamma^{\mu_1}(t_1) \cdots \gamma^{\mu_n}(t_n) &= \sum_P (-)^{\text{sgn}(P)} \Theta(t_{P(1)}, \dots, t_{P(n)}) \gamma^{\mu_{P(1)}} \cdots \gamma^{\mu_{P(n)}}, \quad n = 2, 3, \dots, \\ T\gamma^\mu(\tau) &= \gamma^\mu, \quad \Theta(t_1, \dots, t_n) = \Theta(t_1 - t_2) \cdots \Theta(t_{n-1} - t_n). \end{aligned} \quad (5)$$

where $\text{sgn}(P)$ stays for the parity of the permutation P . In the T product $\gamma^\mu(t)$ anticommute, i.e. behave like Grassmann-odd objects. Another product of γ -matrices in which they have the same behavior is the symmetrical product,

$$\begin{aligned} \text{Sym} \gamma^{\mu_1} \cdots \gamma^{\mu_n} &= \frac{1}{n!} \sum_P (-)^{\text{sgn}(P)} \gamma^{\mu_{P(1)}} \cdots \gamma^{\mu_{P(n)}}, \quad n = 1, 2, \dots, \\ \text{Sym} \gamma^\mu &= \gamma^\mu. \end{aligned} \quad (6)$$

In contrast with the case of T product, γ matrices in the Sym-products carry discrete indices only and the latter take a finite number D of values. Hence, due to the antisymmetry of (6) under permutations of the indices, every Sym-product of more than D γ matrices vanishes. The unique (up to permutations) non-vanishing Sym-product of γ matrices, $\text{Sym} \gamma^0 \cdots \gamma^{D-1}$, in case of D - odd coincides with the identity operator 1 due to the anticommutation relations

(2). For D - even the matrix $\gamma^D = \gamma^0 \cdots \gamma^{D-1}$ is distinct from 1. So, in any dimension D the identity 1 and the matrices

$$\text{Sym } \gamma^{\mu_1} \cdots \gamma^{\mu_k}, \quad \mu_1 < \mu_2 < \cdots < \mu_k, \quad k = 1, 2, \dots, 2\left[\frac{D}{2}\right]$$

form a basis in the associative algebra generated by $\gamma^0, \dots, \gamma^{D-1}$ and will be referred to as independent γ - matrix structures. A modification of the Wick theorem allows one to express the T products in terms of Sym-products of γ matrices. The difference between the T product and the Sym-product of two γ matrices (the contraction), being proportional to their anticommutator, is a c -number,

$$\begin{aligned} T\gamma^{\mu_1}(t_1)\gamma^{\mu_2}(t_2) &= \text{Sym } \gamma^{\mu_1}\gamma^{\mu_2} + \Delta^{\mu_1\mu_2}(t_1, t_2), \\ \Delta^{\mu_1\mu_2}(t_1, t_2) &= \eta^{\mu_1\mu_2}\epsilon(t_1 - t_2), \quad \epsilon(t) = \begin{cases} 1, & t > 0 \\ -1, & t < 0 \end{cases}. \end{aligned} \quad (7)$$

Let a functional

$$F[\zeta] = \sum_n \int_0^1 dt_1 \cdots \int_0^1 dt_n f_{\mu_1 \dots \mu_n}(t_1 \cdots t_n) \zeta^{\mu_1}(t_1) \cdots \zeta^{\mu_n}(t_n) \quad (8)$$

on the space of Grassmann-odd valued functions $\xi^\mu(t)$ be given. Then the matrix $TF[\gamma]$ can be presented as a series in Sym-products

$$TF[\gamma] = \text{Sym} \left[\exp \left\{ -\frac{1}{2} \frac{\delta_\ell}{\delta \zeta_\mu} \star \Delta^{\mu\nu} \star \frac{\delta_\ell}{\delta \zeta_\nu} \right\} F[\zeta] \Big|_{\zeta(t)=\gamma} \right], \quad (9)$$

where $\frac{\delta_\ell}{\delta \zeta_\mu}$ stays for left derivative, and condensed notation is used in which the integrations over time are denoted by star, i.e.

$$\frac{\delta_\ell}{\delta \zeta_\mu} \star \Delta^{\mu\nu} \star \frac{\delta_\ell}{\delta \zeta_\nu} = \int_0^1 dt_1 \int_0^1 dt_2 \frac{\delta_\ell}{\delta \zeta_\mu(t_1)} \Delta^{\mu\nu}(t_1, t_2) \frac{\delta_\ell}{\delta \zeta_\nu(t_2)}.$$

Sometimes discrete indices will be also omitted. In this case all tensors of second rank have to be understood as matrices with lines marked by the first contravariant indices of the tensors, and with columns marked by the second covariant indices of the tensors.

The representation (9) is a functional formulation of the Wick theorem (Hori procedure [7]), modified to the fermionic case and to transition from T to Sym-product [8]. To use the

Wick theorem (9) in the problem at hand we may replace the \mathcal{P} product in (3) for the T product,

$$\mathcal{P} \exp \left\{ \int_0^1 \omega_{\mu\nu} \sigma^{\mu\nu}(t) dt \right\} = T \exp \left\{ \int_0^1 \omega_{\mu\nu} \gamma^\mu(t) \gamma^\nu(t) dt \right\}. \quad (10)$$

To justify the formula (10) one has to define the T product also for coinciding values of some continuous indices (the chronological prescription (6) fails to do it) and then to check (10) itself. It is convenient to define the T product for all values of the times by

$$\begin{aligned} T\gamma^{\mu_1}(t_1) \cdots \gamma^{\mu_n}(t_n) \\ = \text{Sym} \left[\exp \left\{ -\frac{1}{2} \frac{\delta_\ell}{\delta \zeta_\mu} \star \Delta^{\mu\nu} \star \frac{\delta_\ell}{\delta \zeta_\nu} \right\} \zeta^{\mu_1}(t_1) \cdots \zeta^{\mu_n}(t_n) \Big|_{\zeta=\gamma} \right], \quad n = 1, 2, \dots, \end{aligned} \quad (11)$$

where $\Delta^{\mu\nu}$ is given by (7), $\Delta^{\mu\nu}(t, t) = \eta^{\mu\nu}\epsilon(0)$ and some finite value has been assigned to $\epsilon(0)$. Due to Wick theorem (9) this definition is compatible with the chronological prescription (5). Using (11) one obtains

$$T\gamma^{\mu_1}(t_1)\gamma^{\mu_2}(t_2) \cdots \gamma^{\mu_n}(t_n)\gamma^{\nu_n}(t_n) = \mathcal{P}(\sigma^{\mu_1\nu_1}(t_1) + \epsilon(0)) \cdots (\sigma^{\mu_n\nu_n}(t_n) + \epsilon(0)), \quad (12)$$

where the times t_1, \dots, t_n are supposed to be distinct. Substituting (12) in $T \exp\{\omega_{\mu\nu} \gamma^\mu \star \gamma^\nu\}$ one finds that the terms depending on $\epsilon(0)$ vanish due to the antisymmetry of ω , and Eq.(10) takes place independently of the value assigned to $\epsilon(0)$.

III. PATH INTEGRAL FORMULATION OF THE HORI PROCEDURE

Wick theorem (9) admits a path-integral formulation. We define Gaussian and quasi-Gaussian path integrals over a space of Grassmann-odd trajectories in the framework of the perturbation theory approach [9-11]. The first one is defined as

$$\begin{aligned} I(K, \rho, E) &= \int_E D\xi \exp \left\{ \frac{1}{4} \xi^\mu \star K_{\mu\nu} \star \xi^\nu + \rho_\mu \star \xi^\mu \right\}, \\ &= \Lambda \text{Det } K^{1/2} \exp \{ \rho_\mu \star G^{\mu\nu} \star \rho_\nu \}, \end{aligned} \quad (13)$$

where $\xi^\mu(t)$ are Grassmann-odd trajectories of integration, $\rho_\mu(t)$ are Grassmann-odd sources, K is a Grassmann-even antisymmetric kernel $K_{\mu\nu}(t, t') = -K_{\nu\mu}(t', t)$, $G^{\mu\nu}(t, t')$ is an inverse kernel (Green function),

$$\int_0^1 dt' K_{\mu\nu}(t, t') G^{\nu\lambda}(t', t'') = \delta_\mu^\lambda \delta(t' - t''), \quad (14)$$

and Λ is a numerical factor which contains no parameters essential to the theory (parameters defining the matrices $K_{\mu\nu}(t, t')$). In general, Eq.(14) has more than one solution and $G(t, t')$ is specified by imposing some boundary conditions. In a natural way these boundary conditions can be understood as defining the space of integration E . In particular, the kernel K is not degenerate on E , i.e. the homogeneous equation $\int_0^1 dt' K_{\mu\nu}(t, t') \xi^\nu(t') = 0$ has not nontrivial solutions in E . Thus, the equation (14) for the Green function has an unique solution. One can understand the space E as one of functions of the form $\xi^\mu(t) = \int_0^1 dt' K_{\mu\nu}(t, t') \rho^\nu(t')$ where ρ belong to the space of sources [8]. In this case the invariance of the space E under the shifts on such functions is a trivial fact, the latter is important for efficient manipulations with the integrals under consideration. The quasi-Gaussian path integrals are defined via the Gaussian ones by the prescription

$$\int_E D\xi \exp \left\{ \frac{1}{4} \xi^\mu \star K_{\mu\nu} \star \xi^\nu + \rho_\mu \star \xi^\mu \right\} F[\xi] = F\left[\frac{\delta_l}{\delta\rho}\right] I(K, \rho, E), \quad (15)$$

where $F[\xi]$ are arbitrary analytic functionals on E and $\frac{\delta_l}{\delta\rho}$ stays for the left derivatives. In the construction under consideration we encounter matrices $K_{\mu\nu}(t, t')$ part of the indices of which are continuous. To avoid problems with the calculation of the determinants of such matrices as well as problems with the factor Λ definition we may consider the relative quantities

$$\frac{I(K, \rho, E)}{I(K_0, 0, E)} = \text{Det}(K/K_0)^{\frac{1}{2}} \exp \{ \rho_\mu \star G^{\mu\nu} \star \rho_\nu \}, \quad (16)$$

which are sufficient for our purposes. The matrix K_0 can be often chosen in a form simplifying the calculation of the determinant $\text{Det}(K/K_0)$ (see further).

We will use two properties of the quasi-Gaussian path integrals which can be checked using the given definitions. First, the Gaussian path integral can be expressed as a quasi-Gaussian one,

$$I(K, \rho, E) = \exp \left\{ \frac{1}{4} \frac{\delta_\ell}{\delta\rho_\mu} \star (K - K_0)^{\mu\nu} \star \frac{\delta_\ell}{\delta\rho_\nu} \right\} I(K_0, \rho, E), \quad (17)$$

provided both Gaussian integrals $I(K, \rho, E)$ and $I(K_0, \rho, E)$ exist. Second, quasi-Gaussian path integrals are invariant under the shifts, i.e.

$$\int_E D\xi \exp \left\{ \frac{1}{4} (\xi + \zeta)^\mu \star K_{\mu\nu} \star (\xi + \zeta)^\nu \right\} F[\xi + \zeta] = \int_E D\xi \exp \left\{ \frac{1}{4} \xi^\mu \star K_{\mu\nu} \star \xi^\nu \right\} F[\xi], \quad (18)$$

where ζ^μ is an arbitrary trajectory from E .

The path-integral formulation of the Wick theorem (9) is based on the following representation of the quadratic exponent,

$$\exp \{ \rho_\mu \star G^{\mu\nu} \star \rho_\nu \} = \frac{I(K, \rho, E)}{I(K, 0, E)}. \quad (19)$$

Choosing $G^{\mu\nu}(t, t') = -\frac{1}{2} \Delta^{\mu\nu}(t, t')$ (where Δ is given by (7)), the matrix K is easily recognized to be $(K_0)_{\mu\nu}(t, t') = -\eta_{\mu\nu} \delta'(t - t')$, and the space E is determined by the boundary condition satisfied by Δ ,

$$\Delta^{\mu\nu}(0, t) + \Delta^{\mu\nu}(1, t) = 0, \quad 0 < t < 1. \quad (20)$$

According to the definition given, E in (19) is the space of Grassmann-odd trajectories $\xi^\mu(t)$ obeying the antiperiodic boundary condition

$$\xi(0) + \xi(1) = 0. \quad (21)$$

Replacing the odd sources $\rho_\mu(t)$ in (19) by left derivatives and applying the operator obtained to a functional $F[\zeta]$, one gets

$$\exp \left\{ -\frac{1}{2} \frac{\delta_\ell}{\delta\zeta_\mu} \star \Delta^{\mu\nu} \star \frac{\delta_\ell}{\delta\zeta_\nu} \right\} F[\zeta] = \int_{\xi(0)+\xi(1)=0} \mathcal{D}\xi \exp \left\{ -\frac{1}{4} \xi \star \xi \right\} F[\xi + \zeta], \quad (22)$$

where

$$\mathcal{D}\xi = \frac{D\xi}{\int_{\xi(0)+\xi(1)=0} D\xi \exp \left\{ -\frac{1}{4} \xi \star \xi \right\}}. \quad (23)$$

Using Eq.(22) one can present Wick theorem (9) in the form

$$TF[\gamma] = \text{Sym} \left[\int_{\xi(0)+\xi(1)=0} \mathcal{D}\xi \exp \left\{ -\frac{1}{4} \xi \star \xi \right\} F[\xi + \zeta] \Big|_{\zeta=\gamma} \right]. \quad (24)$$

IV. REDUCTION OF THE OPERATOR FUNCTIONS

Choosing the functional $F[\zeta]$ in (24) of the form

$$F[\zeta] = \exp \left\{ \int_0^1 \omega_{\mu\nu} \zeta^\mu(t) \zeta^\nu(t) dt \right\},$$

and using (10), one gets the following representation for the matrix R_0

$$R_0 = \text{Sym} \left[\int_{\xi(0)+\xi(1)=0} \mathcal{D}\xi \exp \left\{ -\frac{1}{4} \xi * \xi \right\} \exp \{ \omega_{\mu\nu} (\xi + \zeta)^\mu * (\xi + \zeta)^\nu \} \right]_{\zeta=\gamma}. \quad (25)$$

The quasi-Gaussian path integral in (25) can be understood as a Gaussian one due to the property (17). Taking into account Eq.(23) one gets

$$R_0 = \text{Sym} \left[\frac{I(K_\omega, 2\zeta\omega, E)}{I(K_0, 0, E)} \exp(\omega_{\mu\nu} \zeta^\mu * \zeta^\nu) \Big|_{\zeta=\gamma} \right], \quad (26)$$

where

$$K_\omega(t, t') = -\eta \delta'(t - t') + 4\omega \delta(t - t'). \quad (27)$$

Evaluating the ratio of the path integrals in (26) by means of (16) and setting $\zeta^\mu(t) = \gamma^\mu$ one obtains

$$R_0 = \left(\text{Det} \frac{K_\omega}{K_0} \right)^{1/2} \text{Sym} \exp \{ M_{\mu\nu} \gamma^\mu \gamma^\nu \}, \quad (28)$$

where

$$M_{\mu\nu} = \omega_{\mu\nu} - 4\omega_{\mu\kappa} * G_\omega^{\kappa\lambda} * \omega_{\lambda\nu}, \quad (29)$$

G_ω being the Green function for K_ω ,

$$\int_0^1 (K_\omega)_{\mu\nu}(t, t') G_\omega^{\nu\lambda}(t', t'') = \delta_\mu^{\lambda}(t, t''),$$

obeying the boundary condition (20). Evaluating

$$G_\omega(t, t') = -\frac{1}{2} e^{4\omega(t-t')} (\epsilon(t-t') - \tanh 2\omega),$$

and substituting in (28) we find

$$M = \frac{1}{2} \tanh 2\omega. \quad (30)$$

Calculating the determinant

$$\text{Det}(K_\omega K_0^{-1}) = \exp \text{Tr} \left\{ 4\omega \int_0^1 G_{s\omega} ds \right\} = \det \cosh 2\omega, \quad (31)$$

and substituting (30), (31) in (28) we finally get

$$R_0 = \exp \{ \omega_{\mu\nu} \gamma^\mu \gamma^\nu \} \\ = (\det \cosh 2\omega)^{1/2} \text{Sym} \exp \left\{ \frac{1}{2} (\tanh 2\omega)_{\mu\nu} \gamma^\mu \gamma^\nu \right\}. \quad (32)$$

A remarkable feature of the expansion in the RHS of Eq.(32) is that it contains only a finite number of terms. Indeed, every Sym-product of more than D γ -matrices vanishes. We have found, in fact, an explicit decomposition, valid in any dimensions, of the spinor representation matrix $\exp\{\omega_{\mu\nu} \gamma^\mu \gamma^\nu\}$ for the Lorentz transformation $L = \exp 4\omega$ in terms of the independent γ -matrix structures.

Taking $D = 3$ where, for example, $\gamma^0 = \sigma^3$, $\gamma^1 = i\sigma^1$, $\gamma^2 = i\sigma^2$, we get

$$R_0 = \exp \{ \omega_{\mu\nu} \gamma^\mu \gamma^\nu \} = (\det \cosh 2\omega)^{1/2} \left[1 + \frac{1}{2} (\tanh 2\omega)_{\mu\nu} \gamma^\mu \gamma^\nu \right], \quad (33)$$

which can be easily transformed to the familiar form

$$\exp \left\{ \frac{i}{2} \theta \cdot \sigma \right\} = \cos \frac{\theta}{2} + i \mathbf{n} \cdot \sigma \sin \frac{\theta}{2}, \quad \theta = \theta \mathbf{n}, \quad \mathbf{n}^2 = 1,$$

where $\theta^2 = \sum_{i=1}^3 \theta_i^2$, $\theta_1 = 4i\omega_{20}$, $\theta_2 = 4i\omega_{01}$, $\theta_3 = -4\omega_{12}$.

In the case $D = 4$ one obtains

$$R_0 = \exp \{ \omega_{\mu\nu} \gamma^\mu \gamma^\nu \} \\ = (\det \cosh 2\omega)^{1/2} \left[1 + \frac{1}{2} (\tanh 2\omega)_{\mu\nu} \sigma^{\mu\nu} + \frac{1}{8} \epsilon^{\kappa\lambda\mu\nu} (\tanh 2\omega)_{\kappa\lambda} (\tanh 2\omega)_{\mu\nu} \gamma^5 \right], \quad (34)$$

where $\gamma^5 = \gamma^0 \gamma^1 \gamma^2 \gamma^3$ and $\epsilon^{\kappa\lambda\mu\nu}$ is the Levi-Civita symbol normalized by $\epsilon^{0123} = 1$. A different form of the decomposition in the left side of (34) was obtained in [12] using direct combinatoric method and concrete properties of γ -matrices in four dimensions,

$$R_0 = [16G(L)]^{-1/2} \left[G(L) + \frac{1}{2} \epsilon_{\mu\nu\rho\sigma} L^{\mu\nu} L^{\rho\sigma} \gamma^5 - (L^2)_{\mu\nu} \sigma^{\mu\nu} + (2 + \text{tr}L) L_{\mu\nu} \sigma^{\mu\nu} \right],$$

$$G(L) = 2(1 + \text{tr}L) + \frac{1}{2} (\text{tr}L)^2 - \frac{1}{2} \text{tr}L^2. \quad (35)$$

The equivalence of the decompositions (34) and (35) can be checked by a straightforward, although long, calculation which we do not present here. We stress again that the derivation in paper [12] is strongly related to $D = 4$ and its generalization to other dimensions is not clear.

For disentangling more complicated operator functions, in particular those of the form (1), it is convenient to introduce the generating functional

$$J[\rho, \zeta] = \int_{\xi(0)+\xi(1)=0} \mathcal{D}\xi \exp \left\{ -\frac{1}{4} \xi \star \dot{\xi} + \omega_{\mu\nu} (\xi + \zeta)^\mu \star (\xi + \zeta)^\nu + \rho_\mu \star (\xi + \zeta)^\mu \right\}. \quad (36)$$

Then

$$R_k = \lim_{t_k \rightarrow 1} \cdots \lim_{t_1 \rightarrow 1} \text{Sym} \left[\frac{\delta_{\xi}^k}{\delta \rho_\alpha(t_1) \cdots \delta \rho_\beta(t_k)} J[\rho, \zeta] \Big|_{\rho=0, \zeta=\gamma} \right]. \quad (37)$$

Taking into account (23), the generating functional $J[\rho, \zeta]$ is calculated by means of (16), (31) to be

$$J[\rho, \zeta] = (\det \cosh 2\omega)^{1/2} \exp \left\{ (\rho + 2\zeta\omega)_\mu \star G_\omega^{\mu\nu} \star (\rho - 2\omega\zeta)_\nu + \omega_{\mu\nu} \zeta^\mu \star \zeta^\nu \right\}. \quad (38)$$

Using Eqs.(37), (38) one finds a formula which is valid in any dimensions

$$R_1 = \gamma^\alpha \exp \{ \omega_{\mu\nu} \gamma^\mu \gamma^\nu \} = \text{Sym} \left[(\eta + \tanh 2\omega)^{\alpha\kappa} \gamma_\kappa \exp \left\{ \frac{1}{2} (\tanh 2\omega)_{\mu\nu} \gamma^\mu \gamma^\nu \right\} \right]. \quad (39)$$

For $D = 4$ the expression in the RHS reduces to

$$R_1 = \gamma^\alpha \exp \{ \omega_{\mu\nu} \gamma^\mu \gamma^\nu \}$$

$$= (\eta + \tanh 2\omega)^{\alpha\kappa} \gamma_\kappa + \frac{1}{2} \epsilon^{\kappa\mu\nu\lambda} (\eta + \tanh 2\omega)^\alpha{}_\kappa (\tanh 2\omega)_{\mu\nu} \gamma^5 \gamma^\lambda. \quad (40)$$

Another representation for the left side of (40) has been derived in $D = 4$ using concrete properties of γ -matrices in such dimensions [13],

$$R_1 = \gamma^\alpha \exp \{ \omega_{\mu\nu} \gamma^\mu \gamma^\nu \} = (e^{2\omega} \cos 2\omega^*)^\alpha{}_\kappa \gamma^\kappa + (e^{2\omega} \sin 2\omega^*)^\alpha{}_\kappa \gamma^5 \gamma^\kappa. \quad (41)$$

One can prove the equivalence of both decompositions (40,41).

As it was mentioned in the Introduction operator expressions of the form (1) appear often in different constructions especially in quantum field theory. Their decompositions in the independent γ -matrix structures are necessary for concrete calculations. A simple example gives us Dirac propagator of a spinning particle in a constant uniform electromagnetic field calculated first by Schwinger [14] in four dimensions:

$$S_0^c(x_{out}, x_{in}) = \left[\gamma^\mu \left(i \frac{\partial}{\partial x_{out}^\mu} - e A_\mu(x_{out}) \right) + m \right] \int_0^\infty ds g(x_{out}, x_{in}, s), \quad (42)$$

where the transformation function g has the form

$$g(x_{out}, x_{in}, s) = \frac{1}{16\pi^2} \left(\det \frac{\sinh eFs}{eF} \right)^{-1/2}$$

$$\times \exp \left\{ i \frac{e}{2} x_{out}^\mu F x_{in}^\mu - sm^2 - i \frac{e}{4} (x_{out} - x_{in}) F \coth(eFs) (x_{out} - x_{in}) + \frac{es}{2} F_{\mu\nu} \sigma^{\mu\nu} \right\}, \quad (43)$$

and contains an operator construction of the form R_0 . By means of the formula (34) one can get the explicit γ -matrix structure of the transformation function to be

$$g(x_{out}, x_{in}, s) = \frac{1}{16\pi^2} \left(\det \frac{\tanh eFs}{eF} \right)^{-1/2}$$

$$\times \exp \left\{ i \frac{e}{2} x_{out}^\mu F x_{in}^\mu - sm^2 - i \frac{e}{4} (x_{out} - x_{in}) F \coth(eFs) (x_{out} - x_{in}) \right\}$$

$$\times \left[1 + \frac{1}{2} (\tanh eFs)_{\mu\nu} \sigma^{\mu\nu} + \frac{1}{8} \epsilon^{\alpha\beta\mu\nu} (\tanh eFs)_{\alpha\beta} (\tanh eFs)_{\mu\nu} \gamma^5 \right]. \quad (44)$$

Acknowledgments

D.M.G and P.B.B. thank Brazilian Foundation CNPq for support. S.I.Z. thanks Departamento de Física Matemática at Universidade de São Paulo for hospitality.

REFERENCES

- [1] F.A. Berezin, *The Method of Second Quantization* (Nauka, Moscow, 1965); (Academic Press, New York, 1966); *Introduction to Superanalysis* (D. Reidel, Dordrecht 1987); B. DeWitt, *Supermanifolds* (Cambridge University Press, Cambridge, 1985)
- [2] F. Bloch, *Zs. f. Phys.* **74** (1932) 295
- [3] R.P. Feynman, *Phys.Rev.* **84** (1951) 108
- [4] D.A. Kirznitz, *Proc. P.N. Lebedev Phys. Inst.*, Vol. XVI, (1951) 3; *Field Methods in the Theory of Many Particles*, (Atomizdat, Moscow 1963)
- [5] J. Schwinger, *Quantum Theory of Angular Momentum*, ed L.C. Biedenharn and H. van Dam (New York: Academic 1965)
- [6] D. Wick, *Phys. Rev.*, **80** (1950) 268
- [7] S. Hori, *Progr. Theor. Phys.*, **7** (1952) 578
- [8] A.N. Vasiliev, *Functional Methods in Quantum Field Theory and Statistics*, (Leningrad State University Press, Leningrad 1976)
- [9] A.A. Slavnov, *Teor. Mat. Fiz.* **22** (1975) 177
- [10] A.A. Slavnov and L.D. Faddeev, *Introduction into Quantum Theory of Gauge Fields*, (Nauka, Moscow 1978)
- [11] D.M. Gitman and I. Tyutin, *Quantization of Fields with Constraints*, (Springer-Verlag, Berlin 1991)
- [12] A.J. Macfarlane, *Commun. Math. Phys.*, **2** (1966) 133
- [13] V.N. Baier, V.M. Katkov and V.M. Strakhovenko, *Sov. Phys. JETP*, **41** (1975) 198
- [14] J.Schwinger, *Phys.Rev.* **82** (1951) 664