

TABLES

ϵ (GeV)	$\tan\beta$	MSSM rescaling	$\tau\tau + \cancel{p}_T$	Exotic channels	dijet+ $\tau + \cancel{p}_T$	Combined results
0	2	86	—	—	—	86
	35	86	—	—	—	86
0.1	2	86	—	—	—	86
	35	80	—	—	61	86
1	2	—	57.6	—	63.7	81
	35	—	62.2	—	80	80
10	35	—	70	—	—	70

TABLE I. 95% CL chargino mass limits in GeV that can be derived from negative searches at LEP II.

FIGURES




FIG. 1. Allowed tau neutrino masses as a function of ϵ (light shaded areas). Within the dark shaded area chargino pair can be produced at $\sqrt{s} = 172$ GeV.




FIG. 2. $e^+e^- \rightarrow \chi^+\chi^-$ cross section, in the large $m_{\tilde{\nu}}$ limit, versus chargino mass for the parameter region defined in Eqs. (17) and (18) and $\sqrt{s} = 172$ GeV.




FIG. 3. Typical neutralino and chargino decay branching ratios as a function of ϵ for $\mu = 150$ GeV, $M_2 = 100$ GeV, and $\tan\beta = 35$.




FIG. 4. 95% CL excluded region in the (μ, M_2) plane (dark shaded area) in the MSSM limit $\epsilon = v_L = 0$ for $\tan\beta = 2$, $\sqrt{s} = 172$ GeV, and an integrated luminosity of 300 pb^{-1} . The light shaded zone is excluded in the MSSM limit by LEP I. The solid curve denotes the kinematical limit.




FIG. 5. Same as in Fig. 4 but for $\tan\beta = 35$




FIG. 8. Same as Fig. 6 but for $\epsilon = 1$ GeV and $\tan\beta = 2$.




FIG. 9. Same as Fig. 8 but for $\tan\beta = 35$.




FIG. 10. Same as Fig. 9 but for $\epsilon = 10$ GeV.