



Instituto de Física
Universidade de São Paulo

**K* nucleon hyperon form factors and nucleon
strangeness**

Forkel, H

Institut für Theoretische Physik, Universität Heidelberg, Heidelberg, Germany

Navarra, F. S.; Nielsen, M.

Instituto de Física, Universidade de São Paulo, SP, Brazil

Publicação IF - 1362/1999

High Energy Physics - Phenomenology,

abstract

hep-ph/9904496

1362
 SBI/IFUSP
 BASE: 04
 SYS N°: 1081288

From: Hilmar Forkel <forkel@frodo.tphys.uni-heidelberg.de>
 Date: Thu, 29 Apr 1999 19:31:19 GMT (15kb)
 Date (revised v2): Mon, 21 Feb 2000 21:46:38 GMT (23kb)

K* nucleon hyperon form factors and nucleon strangeness

Authors: Hilmar Forkel, Fernando S. Navarra, Marina Nielsen

Comments: 9 pages, 8 figures, new co-author, discussion extended to the momentum dependence of the strange vector form factors

Report-no: HD-TVP-99-08, IFUSP-1356/99

→ mudar p 1362
 or o De Salvo

A crucial input for recent meson hyperon cloud model estimates of the nucleon matrix element of the strangeness current are the nucleon-hyperon-K* (NYK*) form factors which regularize some of the arising loops. Prompted by new and forthcoming information on these form factors from hyperon-nucleon potential models, we analyze the dependence of the loop model results for the strange-quark observables on the NYK* form factors and couplings. We find, in particular, that the now generally favored soft N-Lambda-K* form factors can reduce the magnitude of the K* contributions in such models by more than an order of magnitude, compared to previous results with hard form factors. We also discuss some general implications of our results for hadronic loop models.

Paper: Source (23kb), PostScript, or Other formats

(N.B.: delivery types and potential problems)

refers to , cited by

Links to: xxx, hep-ph, /find, /abs (-/+), /9904, ?

www-admin@xxx.lanl.gov

K^* nucleon hyperon form factors and nucleon strangeness

Hilmar Forkel

*Institut für Theoretische Physik, Universität Heidelberg,
Philosophenweg 19, D-69120 Heidelberg, Germany*

Fernando S. Navarra and Marina Nielsen

*Instituto de Física, Universidade de São Paulo,
C.P. 66318,
05315-970 São Paulo, SP, Brazil*

Abstract

A crucial input for recent meson hyperon cloud model estimates of the nucleon matrix element of the strangeness current are the nucleon-hyperon- K^* (NYK^*) form factors which regularize some of the arising loops. Prompted by new and forthcoming information on these form factors from hyperon-nucleon potential models, we analyze the dependence of the loop model results for the strange-quark observables on the NYK^* form factors and couplings. We find, in particular, that the now generally favored soft NAK^* form factors can reduce the magnitude of the K^* contributions in such models by more than an order of magnitude, compared to previous results with hard form factors. We also discuss some general implications of our results for hadronic loop models.

Typeset using REVTeX

I. INTRODUCTION

Understanding the non-valence quark content of hadrons remains, despite a history spanning over two decades, a major theoretical challenge. The interest in non-valence physics derives mainly from the unique opportunities which it provides for new insights into quantum aspects of hadron structure beyond the naive and spectroscopically very successful quark model. Important questions in this realm include modifications of the QCD vacuum inside hadrons [1], the mechanism of flavor mixing and the origin of the OZI-rule [2,3], the structure of constituent quarks [4], and the role of gluons in the dynamics of the (isoscalar) strange-quark sea in the nucleon [5].

The strangeness distribution inside of the nucleon [6] represents the most intensely studied example of a hadronic non-valence quark effect. Currently the vector channel of this distribution, as described by the strange vector form factors, is a focus of experimental [7–9] and theoretical [6,10] research. Since systematic and model-independent approaches have still little predictive power for the nucleon's strangeness content (as exemplified by studies in chiral perturbation theory [11–14] and on the lattice [15]), most previous and current theoretical analyses of the strangeness form factors were model-based.

Among the first and most transparent models for the vector form factors were those which implement a kaon-cloud of the nucleon [16] and thus complement pole dominance approaches [17]. In kaon-cloud models the nucleon's strangeness distribution is generated by fluctuations of the "bare" (i.e. nonstrange) nucleon into kaon-hyperon intermediate states which are described by the corresponding one-loop Feynman graphs [16]. The two crucial assumptions underlying the loop model are 1) that the lightest valence-strangeness carrying intermediate states generate the dominant contribution to the strangeness content and hence give at least a rough estimate of its size, and 2) that rescattering (i.e. multi-loop) contributions are suppressed (despite large couplings).

Both of these assumptions have recently been challenged. A dispersive analysis on the basis of analytically continued $K - N$ scattering data demonstrated that rescattering corrections are important even at low momentum transfers, both to restore unitarity and to build up resonance strength in the ϕ meson region [18]. Furthermore, a study in an "unquenched" quark model found the contributions from higher-lying intermediate states (up to surprisingly large invariant masses) indispensable for the calculation of the strange quark distribution [3] and prompted our collaborators and us to investigate these issues in a complementary hadronic one-loop model [19]. In addition to the original $K - Y$ loops ($Y = \Lambda, \Sigma$), we included the next higher-lying intermediate states, i.e. the $K^* - Y$ pairs. In the first part of this study the corresponding loops were evaluated using Bonn-Jülich K^*NY form factors (see below), as in the original $K - Y$ model. The results were discouraging: the K^* contributions were found to be larger than those from the kaon loop, and their dispersive analysis indicated strong unitarity violations.

The anomalously large and apparently unrealistic K^* contributions could be traced to the large K^* tensor couplings and, in particular, to the very large cutoff parameter of the K^*NA form factor taken from the Bonn-Jülich potential model [20]. Since both the Nijmegen potential [21,22] and the forthcoming update of the Bonn-Jülich NY potential [23] find substantially smaller values for this cutoff, we feel that a detailed and quantitative analysis of the cutoff (and coupling) dependence of the K^* contributions (covering the whole