

IFUSP/P-59

ON PROPERTIES OF THE SINGULARITY OF THE GROUND STATE
IN CERTAIN CLASSICAL HEISENBERG MODELS

WALTER F. WRESZINSKI

Instituto de Física
Universidade de São Paulo
São Paulo, Brasil.

B.I.F. - USP

ON PROPERTIES OF THE SINGULARITY OF THE GROUND STATE
IN CERTAIN CLASSICAL HEISENBERG MODELS

Walter F. Wreszinski

Instituto de Física, Universidade de São Paulo, São Paulo,
Brazil

Abstract: We prove that for both the classical ferro- and antiferromagnetic Heisenberg models the infinite volume limit of the ground state energy per unit volume of the system (Hamiltonian plus λ times an operator) is not differentiable at zero in λ for some operators. This characterization of the singularity at $T=0$, which corresponds to Fisher's ([5]) for positive temperature, adds to a number of others, which are to some extent analogous to the several characterizations of phase transitions at $T>0$ ([8]). A comment is made upon a related open problem concerning the ground state of the quantum antiferromagnetic Heisenberg chain.

As far as we know no explicit examples of Fisher's ([5]) characterization of a phase transition in terms of the nondifferentiability of certain infinite volume correlation functions with respect to external parameters exist. In this note we study the analogous characterization for $T=0$, in the case of some classical Heisenberg (anti-) ferromagnets ¹⁾ and prove that it holds. This result on ground states of classical systems (for general ground-state representations, see [9]) adds to some other features of the singularity at $T=0$, known for the one-dimensional chain with nearest-neighbour interactions, namely: divergence of the susceptibility χ_B as β^2 as $\beta \rightarrow \infty$ ([3]), existence of long-range order ([4]), (infinite) asymptotic degeneracy of the highest eigenvalue of the transfer matrix as $\beta \rightarrow \infty$ ([4]) ²⁾, which are to some extent analogous to some of the several alternative determinations of a phase transition at $T>0$ (see, e.g., [8]). To display one more property in this set of alternate descriptions, whose interrelation is not entirely

1) Some references on classical spin systems, to which we refer for additional literature, are [11], [12]..

2) The eigenvalues $\{\lambda_l(\beta)\}$ $l=0,1,2,\dots$ of the transfer matrix, of which the largest $\lambda_0(\beta)$ is simple and the remainder ones are $(2l+1)$ - fold degenerate, become all degenerate with the largest eigenvalue for l odd, i.e., $\lim_{\beta \rightarrow \infty} [\lambda_l(\beta)/\lambda_0(\beta)] = (-1)^l$ ([4])

clear, and the clarification of which is a major problem in the theory of phase transitions, is the motivation of this paper. For notational simplicity, we write out the proof for the one dimensional case and nearest neighbour interactions. However, the result and proof of the forthcoming theorem hold in any number of dimensions, with a Hamiltonian for the region $\Lambda \subset \mathbb{Z}^{\nu}$ (ν arbitrary integer) given by

$$H_{\Lambda} = - \sum_{i,j \in \Lambda} J(i-j) \vec{t}_i \cdot \vec{t}_j \quad (1)$$

where $\vec{t}_i, i \in \Lambda$, are unit vectors, $\sum_{i \in \Lambda} |J(i)| < \infty$ by

stability ([10]), such that Λ may be divided into two "sublattices" A and B ($A \cup B = \Lambda$), with $J(i-j) \leq 0$ if i, j both belong to either A or B, and $J(i-j) \geq 0$ if $i \in A$ and $j \in B$ or vice-versa. If A is the set of nearest neighbours of B, the above conditions correspond to antiferromagnetism, and if A or B are empty we have a ferromagnetic system (see [2]).

Let $\underline{t}_i, i \in [0, N-1]$, be vectors in $\hat{S} = \{ \vec{x} \in \mathbb{R}^3; x^2 = 1 \}$, with $\underline{t}_0 = \underline{t}_N$ (periodic boundary conditions), with components $(\Omega_i \equiv (\theta_i, \phi_i))$

$$t_i^1(\Omega_i) = t_i^1 = \sin \theta_i \cos \phi_i \quad t_i^2(\Omega_i) = t_i^2 = \sin \theta_i \sin \phi_i$$

$$t_i^3(\Omega_i) = t_i^3 = \cos \theta_i$$

$(0 \leq \theta_i < \pi, 0 \leq \phi_i < 2\pi)$

On $\mathcal{H} = \prod_{i=0}^{N-1} L^2(\mathcal{S}, d\Omega_i)$, with $d\Omega_i = \sin\theta_i d\theta_i d\phi_i$,

$$\text{let } C_N^+(r, \Omega) = \sum_{i=0}^{N-1} t_i^3 t_{i+2r}^3 \quad (2a)$$

$$C_N^-(r, \Omega) = \sum_{i=0}^{N-1} t_i^3 t_{i+r}^3 \quad (2b)$$

$$H_N^{(\pm)}(\Omega) = \pm 2J \sum_{i=0}^{N-1} t_i \cdot t_{i+1}, \quad t_0 = t_N, \quad J > 0 \quad (3)$$

with + (resp. -) corresponding to anti- (resp. ferro-) magnetism. The precise way in which a large class of classical spin systems (including (3) and the more general Hamiltonian (1)) is the limit, "as the spin tends to infinity", of the corresponding quantum spin systems is described in [7].

Let

$$g_{N,r}^{(\pm)}(\lambda) = \min_{\Omega \in \mathcal{S}^N} \frac{1}{N} \left[H_N^{(\pm)}(\Omega) + \lambda C_N^{(\pm)}(\Omega) \right] \quad (4)$$

$$\text{and let } g_r^{(\pm)}(\lambda) = \lim_{N \rightarrow \infty} g_{N,r}^{(\pm)}(\lambda) \quad (5)$$

For fixed r , this limit exists by a simple adaptation of the proof in ([9]).

Note that $g_{N,r}^{(\pm)}$ is a concave function of λ ,

Hence $g_r^{(\pm)}$ is also a concave function of λ , whence (e.g.,

[1]) it has both a right-hand derivative $\frac{d^+ g_r^{(\pm)}(\lambda)}{d\lambda}$ and

a left-hand one $\frac{d^- g_r^{(\pm)}(\lambda)}{d\lambda}$.

Theorem $\frac{d^+ g_r^{(+)}(\lambda)}{d\lambda} (\lambda = 0) \neq \frac{d^- g_r^{(+)}(\lambda)}{d\lambda} (\lambda = 0) .$

Proof A possible choice for our "sublattices" is $A = \{0, 2, \dots, N\}, \{B = 1, 3, \dots, (N-1)\}$ if N is even, or $A = \{0, 2, \dots, (N-1)\}$ and $B = \{1, 3, \dots, N\}$ if N is odd. Clearly, for all N ,

$$g_{N,r}^{(+)}(0) = -2J$$

Now consider the state given by $t_i^2 = t_i^3 = 0, \forall i \in [0, N-1]$, and $t_i^1 = +1 \forall i \in [0, N-1]$ in the - (ferromagnetic) case, and $t_i^1 = 1 \forall i \in A$ and $t_i^1 = -1 \forall i \in B$ in the + (anti-ferromagnetic) case. In this state and any λ ,

$$\frac{1}{N} \left[H_N^{(+)}(\Omega) + \lambda C_N^{(+)}(r, \Omega) \right] \text{ takes the value } (-2J), \text{ hence}$$

$$g_{N,r}^{(+)}(\lambda) \leq -2J \quad \text{for any } \lambda \text{ and for all } N. \text{ Hence,}$$

$$1/\lambda \left[g_{N,r}^{+}(\lambda) - g_{N,r}^{+}(0) \right] \leq 0 \quad \forall \lambda > 0, \quad \forall N$$

Hence, $1/\lambda [g_r^{+}(\lambda) - g_r^{+}(0)] \leq 0 \quad \forall \lambda > 0$, from which we get

$$\frac{d^+ g_r^{+}(\lambda)}{d\lambda} (\lambda=0) \leq 0 \tag{6}$$

$$\text{Now, } \frac{d^- g_r^{(+)}(\lambda)}{d\lambda} (\lambda = 0) = \lim_{\lambda \rightarrow 0} [g_r^{(+)}(0) - g_r^{(+)}(-\lambda)] / \lambda =$$

$$= \lim_{\lambda \rightarrow 0_+} \frac{1}{\lambda} \left[\lim_{N \rightarrow \infty} \min_{\Omega \in \Delta} \left\{ \frac{1}{N} H_N^{(+)}(\Omega) \right\} - \lim_{N \rightarrow \infty} \min_{\Omega \in \Delta} \left\{ \frac{1}{N} \left[H_N^{(+)}(\Omega) - \lambda C_N^{(+)}(r, \Omega) \right] \right\} \right] \tag{7}$$

Now, we clearly have

$$\min_{\Omega \in \mathcal{E}^N} \frac{1}{N} \left[H_N^{(+)}(\Omega) \right] = -1 \quad (8)$$

while

$$\frac{1}{N} \left[H_N^{(+)}(\Omega) - \lambda C_N^{(+)}(r, \Omega) \right] \geq -1 - \lambda,$$

and this minimum value is attained, e.g., for

$$t_i^1 = t_i^2 = 0 \quad \forall i \in [0, N-1], \text{ and } t_i^3 = 1 \quad \forall i \in [0, N-1] \text{ in}$$

the - case, and $t_i^3 = 1 \quad \forall i \in A$, and $t_i^3 = -1 \quad \forall i \in B$,
in the + case. Hence, we have

$$\min_{\Omega \in \mathcal{E}^N} \frac{1}{N} \left[H_N^{(+)}(\Omega) - \lambda C_N^{(+)}(r, \Omega) \right] = -1 - \lambda \quad (9)$$

By (8) and (9) in (7) we get immediately

$$\frac{d^- g_r^{(+)}(\lambda)}{d\lambda} (\lambda = 0) = \lim_{\lambda \rightarrow 0} \frac{1}{\lambda} [-1 - (-1 - \lambda)] = 1 \quad (10)$$

The results (6) and (10) imply the assertion of the theorem. \square

Remark Consider the one-dimensional isotropic antiferromagnetic Heisenberg chain for spin S and periodic boundary conditions, described by the Hamiltonian (on $\mathcal{H} = \left(\sum_{i=0}^{N-1} \mathbb{C}^{2S+1} \right)^{\otimes N}$)

$$H_N^S = \frac{2J}{S^2} \sum_{i=0}^{N-1} (S_i \cdot S_{i+1}) \quad S_0 = S_N \quad S_i^{(k)} = \frac{1}{2} \sigma_i^{(k)}$$

$\sigma_i^{(k)}$ $k \in \{1, 3\}$, being spin matrices for spin S , and $J > 0$.

The ground state Ω_N^S of H_N^S is unique ([2]), and we define

$$L_s \equiv \liminf_{r \rightarrow \infty} \liminf_{N \rightarrow \infty} (\Omega_N^s, S_0^3 S_{2r}^3 \Omega_N^s) / S^2$$

If $L_s \geq \gamma > 0$, we may take this to mean that the one-dimensional antiferromagnet exhibits "long-range order" in the ground state.

Define

$$C_N(r) = \sum_{i=0}^{N-1} S_i^3 S_{i+2r}^3 \quad S_0 = S_N$$

$$C_N(s, r, \Omega) = S^2 C_N^{(+)}(r, \Omega)$$

$$H_N(s, \Omega) = S^2 H_N^{(+)}(\Omega)$$

$$f_{N,r}(\lambda, s) = \min_{\Omega \in S^N} \left\{ \frac{1}{N} [H_N(s, \Omega) + \lambda C_N(s, r, \Omega)] \right\} - \min_{\Omega \in S^N} \frac{1}{N} H_N(1, \Omega)$$

Now $f_{N,r}(\lambda, \delta)$ is a double sequence of functions concave in λ .

$$\text{Hence } f(\lambda, \delta) \equiv \liminf_{r \rightarrow \infty} \liminf_{N \rightarrow \infty} f_{N,r}(\lambda, \delta)$$

is also concave in λ . Now, $f_{N,r}(\lambda, \delta)$ is uniformly

continuous (in (r, N) as a function of δ). Hence,

$$\begin{aligned} f(\lambda) - f(0) &\equiv \lim_{\delta \rightarrow 1} f(\lambda, \delta) = \liminf_{r \rightarrow \infty} \liminf_{N \rightarrow \infty} f_{N,r}(\lambda, 1) \\ &= \liminf_{r \rightarrow \infty} \liminf_{N \rightarrow \infty} [f_{N,r}(\lambda, 1) - f_{N,r}(0, 1)] \end{aligned}$$

From the r.h.s. of inequality (6.5) of [7], transcribed

to vacuum expectation values, we easily get

$$\liminf_{s \rightarrow \infty} \liminf_{r \rightarrow \infty} \liminf_{N \rightarrow \infty} (\Omega_N^s, \frac{C_N(r)}{N} \Omega_N^s) / S^2 \geq \lambda^{-1} f(\lambda) =$$

$$= \liminf_{r \rightarrow \infty} \liminf_{N \rightarrow \infty} \lambda^{-1} f_{N,r}(\lambda, 1) = \liminf_{r \rightarrow \infty} \liminf_{N \rightarrow \infty} \lim_{\beta \rightarrow \infty} \lambda^{-1} f_{N,r,\beta}(\lambda, 1)$$

$$\text{where } f_{N,r,\beta}(\lambda, 1) \equiv -\frac{1}{N\beta} \log \frac{\int d\Omega^N e^{-\beta [H_N(1, \Omega) + \lambda C_N(1, r, \Omega)]}}{\int d\Omega^N e^{-\beta H_N(1, \Omega)}}$$

$$\text{If } \lim_{\lambda \rightarrow 0_+} \liminf_{N \rightarrow \infty} \liminf_{\beta \rightarrow \infty} \lim_{\beta \rightarrow \infty} \lambda^{-1} f_{N,r,\beta}(\lambda, 1) \stackrel{(1)}{=} \\ \stackrel{(2)}{=} \liminf_{N \rightarrow \infty} \liminf_{N \rightarrow \infty} \lim_{\beta \rightarrow \infty} \lim_{\lambda \rightarrow 0_+} \lambda^{-1} f_{N,r,\beta}(\lambda, 1) = 1/3$$

by Refs. [3] , [4] , which would follow if, e.g., for λ in a sufficiently small neighbourhood of zero one had

$$\left[\beta < (C_N(1,r,\Omega) - \langle C_N(1,r,\Omega) \rangle_\lambda)^2 >_\lambda \right] / N \leq \text{const.}$$

(independent of r, N, β) (13)

where

$$\langle A_N(r, \Omega) \rangle_\lambda \equiv \frac{\int d\Omega^N e^{-\beta [H_N(1, \Omega) + \lambda C_N(1, r, \Omega)]} A_N(r, \Omega)}{\int d\Omega^N e^{-\beta [H_N(1, \Omega) + \lambda C_N(1, r, \Omega)]}}$$

then we would clearly have $L_S \geq \gamma > 0$ for sufficiently large S , on putting (12) into (11). Unfortunately, we have been unable to prove (12) (or (13) to date. □

Acknowledgement: I am greatly indebted to Professor E.H. Lieb for his encouragement, patience, and various remarks and corrections in a correspondence about the problem described in Remark 1. I am also indebted to P.A. Vuillermot for useful conversations relating to the existence of the limit (5). Finally, I should like to thank the referee for several extremely helpful suggestions, which led to a much simpler proof of the main result than my original one.

References:

- [1] - Hardy, Littlewood, Pólya - Inequalities - Cambr. Univ. Press 1934
- [2] - E.H. Lieb and D.C. Mattis - Jour. Math. Phys. 3, 749 (1962)
- [3] - M.E. Fisher - Am. Jour. Phys. 32, 343 (1964)
- [4] - G.S. Joyce - Phys. Rev. 155, 458 (1967)
- [5] - M.E. Fisher - Jour. Math. Phys. 6, 1643 (1965)
- [6] - e.g., M. Kac - i "Stat. Mech., Phase Transitions and Superfluidity", Boulder Lectures 1966, ed. by M. Chrétien and S. Deser.
- [7] - E.H. Lieb - Comm. Math. Phys. 31, 327 (1973)
- [8] - M. Marinaro and G.L. Sewell - Comm. math. Phys. 24, 310 (1972)
- [9] - D. Ruelle - Comm. Math. Phys. 11, 339 (1969)
- [10] - D. Ruelle - Statistical Mechanics - Rigorous Results - Benjamin 1969
- [11] - P.A. Vuillermot, M.V. Romerio - Comm. Math. Phys. to appear and work in preparation.
- [12] - A. Bortz and R.B. Griffiths - Comm. Math. Phys. 26, 102 (1972)