

INSTITUTO DE FÍSICA

preprint

IFUSP/P 304
B.I.F. - USP

IFUSP/P-304
ON THE ENERGETICS OF THE GAMOW-TELLER RESONANCES

K. Nakayama

Departamento de Física Nuclear, Instituto de Física,
Universidade de São Paulo, 05508 São Paulo, SP, Brasil

A. Pío Galeão

Dept. Física, Instituto de Geociências e Ciências Exatas,
Universidade Estadual Paulista, 13500 Rio Claro, SP, Brasil

F. Krmpotić

Departamento de Física, Facultad de Ciencias Exactas
Universidad Nacional de La Plata, 1900 La Plata, Argentina

NOV/1981

B.I.F. - USP

UNIVERSIDADE DE SÃO PAULO
INSTITUTO DE FÍSICA
Caixa Postal - 20.516
Cidade Universitária
São Paulo - BRASIL

ON THE ENERGETICS OF THE GAMOW-TELLER RESONANCES

K. Nakayama* and A. Pio Galeão**

Departamento de Física Nuclear, Instituto de Física,
Universidade de São Paulo, 05508 São Paulo, SP, Brasil

and

F. Krmpotić***

Departamento de Física, Facultad de Ciencias Exactas
Universidad Nacional de La Plata, 1900 La Plata, Argentina

ABSTRACT

It is suggested that the energy difference between the Gamow-Teller and Fermi resonances is given, in units of MeV, by $E_{GT} - E_F = 26A^{-1/3} - 18.5 (N-Z)/A$. The consequences of this result on the strength of the spin and isospin dependent residual interaction, as well as on the effective axial-vector coupling constant, are discussed.

* Present address: Institut für Kernphysik, Kernforschungsanlage Jülich, D-5170 Jülich, West Germany.

** Present address: Departamento de Física, Instituto de Geociências e Ciências Exatas, Universidade Estadual Paulista, 13500 Rio Claro, SP, Brasil.

*** Member of the Carrera de Investigador Científico del CONICET, Argentina.

The Fermi (F) resonance has been known since the discovery of the well-defined multiplet structure by Anderson and Wong¹⁾, but the Gamow-Teller (GT) one, although foreseen in the early 1960's²⁻⁶⁾, was observed experimentally for the first time only in 1975, by Doering et al.⁷⁾ (through the ⁹⁰Zr(p,n)⁹⁰Nb reaction at incident proton energies of 35 and 45 MeV). This GT resonance was later on confirmed by a new study at $E_p = 120$ MeV⁸⁾. The gathering of the GT strength in ²⁰⁸Pb was detected even more recently, by Horen et al.⁹⁾, in the neutron spectra resulting from bombardment with 120 and 160 MeV protons. Almost simultaneously, Horen et al.¹⁰⁾ have reported a systematic study of the energetics of the GT strength for a number of targets with $A > 90$, with incident proton energies of 120, 160 and 200 MeV. They have also suggested that the energy difference between the GT and F resonances can be represented approximately by (in units of MeV)

$$E_{GT} - E_F = 6.7 - 60T_0 A^{-1} \quad (1)$$

where $T_0 = (N-Z)/2$ is the isospin of the target nucleus.

Quite recently, Sterrenburg et al.¹¹⁾ have performed a (p,n) reaction study with 45 MeV protons at small angles on 17 targets ranging from ⁹⁰Zr to ²⁰⁸Pb. Common features of the spectra were: i) a sharp peak representing the F resonance and ii) a broad peak, interpreted as a GT state, at a slightly higher but target dependent excitation energy.

The above mentioned systematic behaviour of the energy difference $E_{GT} - E_F$, throughout the periodic table, calls for a discussion in the framework of schematic models, in order to elucidate upon the details of the underlying structure.

As the ground state correlations are, in general, negligible for F and GT excitations (in nuclei with an appreciable neutron excess), it is enough to work within the Tamm-Dancoff approximation. We will use a schematic residual force of the form¹²⁻¹⁵⁾

$$H = \frac{\kappa}{2} \sum_{\mu\tau} M^\dagger(\tau=1, \mu\tau) M(\tau=1, \mu\tau), \quad (2)$$

where κ is the coupling constant and

$$M(\tau=1, \mu\tau) = \begin{cases} \sum_{i=1}^A \tau_i \mu_\tau & \text{for the F mode,} \\ \sum_{i=1}^A \sigma(i) \tau_i \mu_\tau & \text{for the GT mode.} \end{cases} \quad (3)$$

Furthermore, a degenerate model for the single-particle energies is assumed,

$$E^{(0)} = \epsilon - \frac{V_1}{A} T_0 + \Delta E_{\text{Coul}}, \quad (4)$$

where V_1 is the symmetry potential, ΔE_{Coul} is the Coulomb energy displacement and ϵ represents the average single particle energy. For the F resonance $\epsilon = 0$, while for the GT mode this quantity should be of the same order of magnitude as the spin-orbit splitting Δ_{ls} , i.e.,

$$\epsilon_{\text{GT}} = \Delta_{\text{ls}} = 20 A^{-\frac{1}{3}} \text{ MeV}, \quad (5)$$

where we have employed the estimate $\epsilon = A^{-\frac{1}{3}}$ (12)

As the interaction (2) shifts the collective states, both F and GT, by κT_0 , the perturbed energies are

$$E = \epsilon - T_0 (V_1/A - 4\kappa) + \Delta E_{\text{Coul}}. \quad (6)$$

In the case of the F resonance ($\epsilon_F=0$) this expression just reflects the self-consistency between the Hartree-Fock field and the collective motion¹⁶; name-

ly as the excitation energy E_F should equal ΔE_{Coul} , one has $\kappa_F = V_1/4A$. (Care must be taken in attaching a physical meaning to the effective mass when this self-consistency is not fulfilled). Consequently,

$$E_{\text{GT}} - E_F = \epsilon_{\text{GT}} + 4T_0 (\kappa_{\text{GT}} - \kappa_F). \quad (7)$$

If, consistent with later discussion, it is assumed that both κ_F and κ_{GT} are proportional to A^{-1} , one sees that the second term of this expression has the same mass and charge dependence as the second term in eq. (1). However, from the estimate (5) one would expect an $A^{-\frac{1}{3}}$ dependence for the first term in eq. (1) instead of it being a constant. This fact induced us to try to adjust the difference $E_{\text{GT}} - E_F$ by a function of the form $C_1 A^{-1/3} + C_2 T_0 A^{-1}$, C_1 and C_2 being free parameters. Inclusion of 25 experimental data⁸⁻¹¹⁾, as shown in fig. 1, yields (in MeV)

$$E_{\text{GT}} - E_F = 26 A^{-\frac{1}{3}} - 37 T_0 A^{-1}, \quad (8)$$

with a $\chi^2 = 2.33$. The same procedure was repeated by omitting the mass dependence $A^{-1/3}$ and the result was (in MeV)

$$E_{\text{GT}} - E_F = 7.0 - 57.8 T_0 A^{-1}, \quad (9)$$

with $\chi^2 = 5.08$. The small difference between eqs. (1) and (9) arises from the fact that we have considered more experimental data than Horen et al.¹⁰⁾

First of all it should be noted that the coefficient of the T_0/A term is considerably reduced when the $A^{-\frac{1}{3}}$ dependence is introduced. Furthermore, as the first term in eq. (8) has a very simple physical meaning, and as eq. (8) reproduces the experimental data much better than eq. (9), in the following discussion we omit further references to the latter. That is, it will be assumed that the correct A and T_0 dependence for the energy difference $E_{\text{GT}} - E_F$ is the one given by eq. (8). Consequently, from eq. (7) we have

$$(\kappa_F - \kappa_{\text{GT}})A = 9.25 \text{ MeV}. \quad (10)$$

Adopting now for the symmetry potential the values suggested by Bohr and Mottelson¹²⁾, and remembering that $A\kappa_F = V_1/4$, we obtain

$$A\kappa_F = 25 \text{ MeV}, \quad A\kappa_{GT} = 15.75 \text{ MeV}, \quad \text{for } V_1 = 100 \text{ MeV}, \quad (11a)$$

and

$$A\kappa_F = 32.5 \text{ MeV}, \quad A\kappa_{GT} = 23.25 \text{ MeV}, \quad \text{for } V_1 = 130 \text{ MeV}. \quad (11b)$$

The above κ_{GT} should be used when only nucleonic degrees of freedom are used. When the effect of the coupling between the nucleon and the isobar $\Delta(1230)$ is included, the GT strength is^{17,18)}

$$\kappa_{GT}^{\Delta} = \frac{\kappa_{GT}}{1 + \kappa_{GT}^{\Delta} x_{\Delta}^{(o)}}, \quad (12a)$$

where $x_{\Delta}^{(o)}$ is the unperturbed isobar-hole response function. The inverse relation reads

$$\kappa_{GT} = \frac{\kappa_{GT}^{\Delta}}{1 - \kappa_{GT}^{\Delta} x_{\Delta}^{(o)}}. \quad (12b)$$

By employing the constituent quark-model value of 72/25 for the ratio of the $\pi N\Delta$ and πNN coupling constants (i.e., $g_{\pi N\Delta}^2/g_{\pi NN}^2 \equiv r_{\pi N\Delta}$), Bohr and Mottelson¹⁷⁾ have shown that

$$x_{\Delta}^{(o)} = -\frac{64}{25} \frac{A}{E_{\Delta}}, \quad (13a)$$

where $E_{\Delta} = 300 \text{ MeV}$ if one ignores nuclear excitation energies in comparison with the Δ -N mass difference. Consequently, from the results (11) we see that $A\kappa_{GT}^{\Delta} = 18.2$ and 29.0 MeV for $V_1 = 100$ and 130 MeV , respectively. The second of these values for κ_{GT}^{Δ} is the same as that obtained by Bohr and Mottelson¹⁷⁾ from the analysis of quenching phenomena in the magnetic dipole moments in the region of ^{208}Pb .

When the experimental value $r_{\pi N\Delta} = 4$, is used rather than that given by the quark model, as done by Brown and Rho¹⁸⁾, one has

$$x_{\Delta}^{(o)} = -\frac{32}{9} \frac{A}{E_{\Delta}}. \quad (13b)$$

With this value for the response function the above mentioned results for κ_{GT}^{Δ} are slightly increased. (It is worth noting that Vento et al.¹⁹⁾ have indicated quite recently that the experimental values for the axial-vector coupling constant g_A and for the ratio $r_{\pi N\Delta}$ provide consistent indications of substantial bag deformation.)

We can easily relate the coupling constants κ_F and κ_{GT} of the separable force given by eq. (2), which has an infinite range, with the parameters f'_0 and g'_0 of the zero-range Migdal force²⁰⁻²²⁾

$$F^{\text{ph}}(\vec{r}_1, \vec{r}_2) = C(f'_0 \vec{\tau}_1 \cdot \vec{\tau}_2 + g'_0 \vec{\tau}_1 \cdot \vec{\tau}_2 \vec{\sigma}_1 \cdot \vec{\sigma}_2) \delta(\vec{r}_1 - \vec{r}_2) \quad (14)$$

where $C = 380 \text{ MeV fm}^3$.^{†1} The corresponding relations are

$$\kappa_F = \frac{Cf'_0}{\Omega}, \quad \kappa_{GT} = \frac{Cg'_0}{\Omega}, \quad (15)$$

where Ω stands for the nuclear volume. When the nucleus is treated as a Fermi gas $\Omega = (4\pi/3)r_0^3 A = 0.58 (4\pi A \text{ fm}^3)$ with $r_0 \approx 1.2 \text{ fm}$. A more reliable estimate can be obtained from the shell model, where Ω^{-1} is the average value of the radial integrals

^{†1} It should be noted that the Migdal force is the direct term of a δ -function interaction.

$$I(\ell_1 \ell_2 \ell_1' \ell_2') = \frac{1}{4\pi} \int R_{n_1 \ell_1}(r) R_{n_2 \ell_2}(r) R_{n_1' \ell_1'}(r) R_{n_2' \ell_2'}(r) r^2 dr. \quad (16)$$

Using wave functions of an harmonic oscillator potential, one has^{4,5)}

$$\Omega = \langle I(\ell_1 \ell_2 \ell_1' \ell_2') \rangle_{av}^{-1} = 4\pi A \text{ fm}^3. \quad (17)$$

Consequently, from eqs. (10), (11), (15) and (17), we obtain that:

$$f'_0 - g'_0 = 0.31, \quad (18)$$

and

$$f'_0 = 0.83, \quad g'_0 = 0.52, \quad (19a)$$

and

$$f'_0 = 1.07, \quad g'_0 = 0.77, \quad (19b)$$

for $V_1 = 100$ and 130 MeV, respectively.

Quite recently, Bertsch et al.²³⁾ have pointed out that the systematics of the GT states, when studied with a zero range interaction, require an interaction strength g'_0 of about 0.53 to 0.63 ($V_{GT} = Cg'_0 = 200-240 \text{ MeV fm}^3$). With a similar value for $g'_0 (=0.65)$ Krewald et al.²⁴⁾ and Osterfeld et al.²⁵⁾ have succeeded in reproducing both the allowed and first forbidden charge-exchange resonances in ^{208}Pb . On the other hand, from the investigation of magnetic states in ^{12}C and ^{16}O a value of $g'_0=0.75$ was obtained²⁶⁾. In addition, Brown and Rho¹⁸⁾ have shown that, when one uses a $\pi + \rho$ exchange force in the nucleon space and includes the screening due to isobar-hole excitations, there results an effective coupling constant g'_0 equal to 0.9.

For a δ -interaction of the form^{5,6)}

$$V(\vec{r}_1, \vec{r}_2) = -2\pi(V_s P_s + V_t P_t) \delta(\vec{r}_1 - \vec{r}_2), \quad (20)$$

where P_s and P_t are singlet-even and triplet-even projection operators one has

$$\kappa_F = \frac{\pi(3V_t - V_s)}{4\Omega}, \quad \kappa_{GT} = \frac{\pi(V_s + V_t)}{4\Omega}. \quad (21)$$

With the values derived here for κ_F and κ_{GT} we get

$$V_t = 163 \text{ MeV fm}^3, \quad V_s = 89 \text{ MeV fm}^3, \quad (V_1 = 100 \text{ MeV}), \quad (22a)$$

and

$$V_t = 223 \text{ MeV fm}^3, \quad V_s = 149 \text{ MeV fm}^3, \quad (V_1 = 130 \text{ MeV}), \quad (22b)$$

when $\Omega=4\pi A \text{ fm}^3$. Ikeda⁵⁾ has employed, in his study of the F and GT resonances in ^{208}Pb , a value of 50 MeV fm^3 for V_s , and he varied V_t from 83 to 200 MeV fm^3 .

Let us now estimate the effective axial-vector coupling constant for allowed β decay processes. In the context of the approximation used here it reads

$$\left(\frac{g_A}{g_A}\right)_{GT}^{eff} = \frac{1 + \kappa_{GT} \chi_{\Delta}^{(o)}}{1 - \kappa_{GT} \chi_N^{(o)}} = \frac{1}{1 - \kappa_{GT} (\chi_N^{(o)} + \chi_{\Delta}^{(o)})}, \quad (23)$$

where

$$\chi_N^{(o)} = -\frac{4T_0}{E_{GT}^{(o)}} \quad (24)$$

is the unperturbed particle-hole response function and $E_{GT}^{(o)} = \Delta_{2s} - 4T_0 \kappa_F + \Delta E_{Coul}$. For example, with $V_1=130$ MeV one has that $(g_A^{eff}/g_A)_{GT} \approx 0.5, 0.5$ and 0.4 for ^{90}Zr , ^{116}Sn and ^{208}Pb , respectively. A similar result for the quenching effect on the axial-vector coupling constant can be found in the literature²⁸⁾.

We can summarize our considerations by saying that the energy difference between the Gamow Teller and Fermi resonances obeys the relation (8), which has a very simple physical meaning. This result was used to fix the coupling strength of the spin-isospin dependent residual interaction. It is worth noticing that the values presented for this coupling constant are based upon the underlying assumption of the self-consistency between the Hartree-Fock field and the collective motion. We also estimate the effective axial-vector coupling constant for allowed β decay.

After finishing the present work we have learned about a similar study performed by Suzuki²⁶⁾ in ^{48}Ca , ^{90}Zr and ^{208}Pb . His value for $(\kappa_F - \kappa_{GT})$ is smaller than ours by a factor of two.

Thanks are due to A. Galonsky for several valuable comments.

REFERENCES

- ¹J.D. Anderson and C. Wong, Phys. Rev. Lett. 7 (1961) 250.
- ²K. Ikeda, S. Fujii and J. I. Fujita, Phys. Lett. 2 (1962) 169.
- ³K. Ikeda, S. Fujii and J.I. Fujita, Phys. Lett. 3 (1963) 271.
- ⁴J.I. Fujita, S. Fujii and K. Ikeda, Phys. Rev. 133 (1964) B549.
- ⁵K. Ikeda, Prog. Theor. Phys. 31 (1964) 434.
- ⁶J.I. Fujita and K. Ikeda, Nucl. Phys. 67 (1965) 145.
- ⁷R.R. Doering, A. Galonsky, D.M. Patterson and G.F. Bertsch, Phys. Rev. Lett. 35 (1975) 1961.
- ⁸D.E. Bainum, J. Rapaport, C.D. Goodman, D.J. Horen, C.C. Foster, M.B. Greenfield and C.A. Gouling, Phys. Rev. Lett. 44 (1981) 1751.
- ⁹D.J. Horen, C.D. Goodman, C.C. Foster, C.A. Gouling, M.B. Greenfield, J. Rapaport, D.E. Bainum, E. Sugarbaker, T.G. Masterson, F. Petrovich and W.G. Love, Phys. Lett. 95B (1980) 27.
- ¹⁰D.J. Horen, C.D. Goodman, D.E. Bainum, C.C. Foster, C. Gaarde, C.A. Gouling, M.B. Greenfield, J. Rapaport, T.N. Taddeucci, E. Sugarbaker, T. Masterson, S.M. Austin, A. Galonsky and W. Sterrenburg, Phys. Lett. 99B (1981) 383.
- ¹¹W.A. Sterrenburg, S.M. Austin, R.P. De Vito and A. Galonsky, Phys. Rev. Lett. 45 (1980) 1839.
- ¹²A. Bohr and B.R. Mottelson, Nuclear Structure (Benjamin, New York, 1975), Vol. 2.
- ¹³F. Krmpotić and F. Osterfeld, Phys. Lett. 93B (1980) 218.

14. F. Krmpotić, Nucl. Phys. A351(1981) 1839.
15. F. Krmpotić, Phys. Rev. Lett. 46 (1981) 1261.
16. N. Auerbach, V. Bernard and Nguyen Van Giai, Nucl. Phys. A337 (1980) 143.
17. A. Bohr and B.R. Mottelson, Phys. Lett. 100B (1981) 10.
18. G.E. Brown and M. Rho, to be published.
19. V. Vento, G. Baym and A.D. Jackson, Phys. Lett. 102B (1981) 97.
20. A.B. Migdal, Theory of finite Fermi systems and applications to atomic nuclei (John Wiley and Sons, New York, 1967).
21. J. Speth, E. Werner and W. Wild, Phys. Reports 33, (1977) 127.
22. F. Krmpotić, K. Ebert and W. Wild, Nucl. Phys. A342 (1980) 497.
23. B. Bertsch, D. Cha and H. Toki, Phys. Rev. 24C (1981) 24.
24. S. Krewald, F. Osterfeld, J. Speth and G.E. Brown, Phys. Rev. Lett. 46 (1981) 103.
25. F. Osterfeld, S. Krewald, H. Dermawan and J. Speth, preprint.
26. Li Chu-Hsia and V. Klemt, to be published.
27. T. Suzuki, Phys. Lett. 104B (1981) 92.
28. Y. Futami, J. Fujita and H. Ohtsubo, Prog. Theor. Phys. 60 (1976) 83 and references therein.

Figure caption

Fig. 1 - Plot of $E_{GT} - E_F$ versus $(N-Z)/A$. The experimental data were taken from ref. 8) for ^{90}Zr , from ref. 10) for ^{90}Zr , ^{92}Zr , ^{94}Zr , ^{112}Sn , ^{116}Sn , ^{124}Sn , ^{169}Tm and ^{208}Pb and from ref. 11) for ^{90}Zr , ^{91}Zr , ^{92}Zr , ^{94}Zr , ^{96}Zr , ^{93}Nb , ^{94}Mo , ^{96}Mo , ^{97}Mo , ^{98}Mo , ^{100}Mo , ^{112}Sn , ^{116}Sn , ^{120}Sn , ^{122}Sn , ^{124}Sn and ^{208}Pb . When the experimental results overlap (in the case of $^{90,92}\text{Zr}$ and ^{208}Pb) we displace them slightly with respect to the correct value of $(N-Z)/A$ for the sake of clarity. The values calculated by means of eq. (8) are indicated by full circles, while the dash-dotted line corresponds to eq. (9).

