

UNIVERSIDADE DE SÃO PAULO

INSTITUTO DE FÍSICA  
CAIXA POSTAL 20516  
01498 - SÃO PAULO - SP  
BRASIL

IFUSP/P 435  
B.I.F. - USP

# publicações

IFUSP/P-435



FORWARD GLORY EFFECTS IN THE ELASTIC SCATTERING  
OF  $^{12}\text{C} + ^{12}\text{C}$

by

M.S. Hussein, A. Lépine-Szily, M.M. Saad and  
A.C.C. Villari

Instituto de Física, Universidade de São Paulo

Outubro/1983

M.S. Hussein\*, A. Lépine-Szily\*, M.M. Saad<sup>†</sup>  
and A.C.C. Villari<sup>†</sup>

Instituto de Física, Universidade de São Paulo  
C.P. 20.516, São Paulo, S.P., Brazil

ABSTRACT

It is shown that the elastic scattering of  $^{12}\text{C} + ^{12}\text{C}$  in the center-of-mass energy range  $6 < E < 31$  MeV exhibits forward glory enhancement. Semiclassical analysis of the quantity  $\Delta\sigma_R \equiv \sigma_R - \int \left[ \frac{d\sigma_{\text{Ruth.}}}{d\Omega} - \frac{d\sigma_{\text{el.}}}{d\Omega} \right] d\Omega$  indicates that the best candidate for a  $^{12}\text{C} + ^{12}\text{C}$  interaction potential is a small-radius, deep Woods-Saxon potential, in qualitative agreement with the results obtained from recent analyses of  $\frac{d\sigma_{\text{el.}}}{d\Omega}$  done at higher energies.

\*Supported in part by the CNPq.

<sup>†</sup>Supported in part by FAPESP.

Recently several analyses of elastic and inelastic scattering data of light-heavy ion systems at intermediate energies have been reported<sup>1)</sup>. The main conclusion reached has been the removal of some of the major ambiguities attached to the ion-ion optical potential usually extracted from data taken at lower energies. This is accomplished at intermediate energies because of the incipient dominance of the farside amplitude over the nearside one<sup>2)</sup>, thus leading to a greater degree of sensitivity to the details of the ion-ion interaction at shorter separation distances.

In a parallel theoretical development<sup>3)</sup>, the observation has been made that the extraction and analysis of the forward glory contribution to the elastic scattering, would also furnish further constraints on the interaction potential. It has been suggested in Ref. 3 that a careful study of the quantity

$$\Delta\sigma_R = \sigma_R - \int d\Omega \left[ \frac{d\sigma_{\text{Ruth.}}}{d\Omega} - \frac{d\sigma}{d\Omega} \right] \quad (1)$$

where  $\sigma_R$  is the total reaction cross section,  $\frac{d\sigma}{d\Omega}$  the elastic differential cross section and  $\frac{d\sigma_{\text{Ruth.}}}{d\Omega}$ , the differential Rutherford cross section, can supply the above mentioned constraint. This comes about as a consequence of the optical theorem which relates  $\Delta\sigma_R$  to the imaginary part of the forward "nuclear" scattering amplitude viz<sup>4)</sup>

$$\Delta\sigma_R = \frac{4\pi}{k} \text{Im} [f(0) - f_{\text{Ruth.}}(0)] \quad (2)$$

where  $f$  and  $f_{\text{Ruth}}$  are the total and Rutherford scattering amplitudes, respectively and  $k$  is the asymptotic wave number

of relative motion. The occurrence of forward glory, a refractive effect, leads to a major enhancement in  $\Delta\sigma_R$ .

In this letter, we present evidence for the forward glory enhancement in  $\Delta\sigma_R$  for the scattering of  $^{12}\text{C}$  on  $^{12}\text{C}$  in the energy range  $6 < E_{\text{CM}} < 31$  MeV.

We have determined the quantity  $\Delta\sigma_R$  from existing experimental data on the total reaction cross section,  $\sigma_R$ , and published values of the sum-of-differences cross section,

$$\sigma_{\text{SOD}} \equiv \int d\Omega \left[ \frac{d\sigma_M}{d\Omega} - \frac{d\sigma_{e\ell}}{d\Omega} \right] \quad \text{where } \frac{d\sigma_M}{d\Omega} \text{ is the Mott cross section.}$$

The total reaction cross section  $\sigma_R$  is obtained from the summed contribution of the complete fusion cross section,  $\sigma_F$ , and the total, angle-integrated, cross section of direct processes  $\sigma_D$ .

Recently Kolata et al.<sup>5)</sup> measured the total  $\alpha$  yield in the  $^{12}\text{C} + ^{12}\text{C}$  fusion. The contribution to  $\sigma_F$  arising from the  $3\alpha$  evaporation, not taken into account in previous fusion measurements, was determined. An anomalous  $\alpha$  yield, which seems to be a direct process, was included in  $\sigma_D$ . We used  $\sigma_R$  from Kolata's work<sup>5)</sup>, and calculated the  $\sigma_R$  from other fusion measurements<sup>6,7,8)</sup> summing to them the  $3\alpha$  evaporation, and considered  $\sigma_D$  as composed of the total angle-integrated inelastic cross section<sup>5,9)</sup> and the anomalous  $\alpha$  yield<sup>5)</sup>.

The quantity  $\sigma_{\text{SOD}}$  was constructed by Treu et al.<sup>10)</sup> and more recently by Ledoux et al.<sup>11)</sup> from the measured elastic scattering angular distributions. The quantity of interest in this letter,  $\Delta\sigma_R$ , was then evaluated, as indicated in Eq. (1), namely  $\Delta\sigma_R = \sigma_R - \sigma_{\text{SOD}}$ .

Owing to the dispersion inherent in both  $\sigma_R$  and  $\sigma_{\text{SOD}}$ , we present our  $\Delta\sigma_R$  as a band, whose width is much smaller than its mean value. This band of points representing

$\Delta\sigma_R$ , is plotted in Fig. 1 vs. the center of mass energy. One sees clearly the beginning of the oscillatory behaviour expected from the theoretical study of Ref. 3). To ascertain the refractive nature of  $\Delta\sigma_R$ , we show in the figure  $\Delta\sigma_R$  calculated in the sharp cut-off approximation<sup>12)</sup>

$$\Delta\sigma_R^{\text{s.c.}} = \frac{2\pi}{k^2} \sum_{l=0}^{l_c} (2l+1) \cos 2\sigma_l \quad (3)$$

where  $\sigma_l$  is the  $l$ -Coulomb phase-shift and  $l_c$  is the sharp cut-off angular momentum that specifies the value of the total reaction cross section through  $\sigma_R = \pi/k^2 (l_c+1)^2$ .

We consider as a criterion for the refractive enhancement in  $\Delta\sigma_R$  due to forward glory scattering, the following

$$\frac{\Delta\sigma_R}{\Delta\sigma_R^{\text{s.c.}}} > 1 \quad (4)$$

which is clearly satisfied by the  $^{12}\text{C} + ^{12}\text{C} - \Delta\sigma_R$  data shown in Fig. 1.

It has been pointed out in Ref. 3) that different optical potentials that give similar reasonable fits to the ratio to Rutherford scattering at small angles, may give quite different  $\Delta\sigma_R$ 's. Thus through the confrontation of the calculated  $\Delta\sigma_R$  with the experimental one, a less ambiguous optical potential may be deduced<sup>13)</sup>. We have tested this idea on our  $^{12}\text{C} + ^{12}\text{C}$  case. We have considered two optical potentials that both generate forward glory scattering namely the corresponding classical deflection function  $\theta(l)$  passes through zero at a finite value of  $l$ ,  $l_{\text{gl}}$ .

The first optical potential we considered in our analysis is the one suggested by the Yale group<sup>14)</sup>. This Saxon-Woods potential, whose parameters are,  $V = 14$  MeV,  $a_v = 0.35$  fm,  $r_v = 1.35$  fm,  $W = 0.4 + 0.1 E_{CM}$  (MeV),  $a_w = 0.35$  fm,  $r_w = 1.40$  fm, reproduces reasonably well the elastic scattering angular distributions of  $^{12}C + ^{12}C$  and accounts for the structure seen in the excitation function at  $90^\circ$ . We calculated  $\Delta\sigma_R$  using the Ford-Wheeler<sup>15)</sup>, stationary-phase approximation, which gives

$$\Delta\sigma_R = \frac{4\pi}{k^2} (\ell_{gl} + 1/2) \left[ \frac{2\pi}{d\ell} \right]_{\ell_{gl}}^{1/2} |S_{\ell_{gl}}^n| \sin \left[ 2(\sigma_{\ell_{gl}} + \delta_{\ell_{gl}}^n) - \frac{\pi}{4} \right] \quad (5)$$

where  $\delta_{gl}$  is the nuclear phase shift evaluated at the forward glory angular momentum,  $\ell_{gl}$ , and  $S_{\ell_{gl}}^n$  is the reflection coefficient at  $\ell_{gl}$ . We have found that the reflection coefficient  $|S_{\ell_{gl}}^n|$  for the Yale potential very close to unity in the energy range of interest. Further, the effect of absorption on  $\ell_{gl}$  was found to be small too. This convinced us that the use of a classical deflection function is more than adequate. The nuclear phase shift  $\delta_{\ell}^n$  is a rapidly varying function of  $\ell$  in the forward glory region. However its value at  $\ell_{gl}$ ,  $\delta_{\ell_{gl}}^n$ , is quite small as our optical model calculation has shown us.

The result of our calculation of  $\Delta\sigma_R$  for the Yale potential, using a classically generated  $\theta$  and  $\ell_{gl}$  and ignoring  $|S_{\ell_{gl}}^n|$  and  $\delta_{\ell_{gl}}^n$ , is shown in Fig. 1 as the dotted curve. We see clearly that the magnitude of  $\Delta\sigma_R$  comes out right, however there is a major discrepancy in the phase. We have also calculated  $\Delta\sigma_R$  for a potential tailored according

to that obtained from the analysis of the intermediate-energy data; a rather small-radius deep Saxon-Wood interaction. We have taken  $V = 250$  MeV,  $r_v = 0.66$  fm and  $a_v = 0.63$  fm<sup>1a)</sup>. The result is presented in Fig. 1 as the full curve. The agreement with the general trend of the data is striking. This agreement with the behavior of the data is made meaningful by the fact that the elastic scattering angular distribution with this real potential and with  $W = 0.4 + 0.3E$  (MeV),  $r_w = 0.93$  fm,  $a_w = 0.35$  fm, is coming out as reasonable as the one obtained with the Yale Potential, as clearly seen in Fig. 2.

The fact that  $\Delta\sigma_R$  is acting as a filter to the appropriate optical potential that best represents the interacting system would be understood easily by the fact that  $\frac{d\sigma}{d\Omega}(\theta)$  probes a certain combination of the optical potential parameters, whereas  $\Delta\sigma_R$  tests a different combination. This situation becomes quite clear at higher energies where the forward glory impact parameter,  $b_{gl} = \ell_{gl}/k$ , becomes independent of energy<sup>16)</sup>

$$b_{gl} = R_v \left[ 1 + \frac{a_v}{2R_v - 3a_v} \left( 3 \ln R_v + \ln \left( \frac{2a_v}{\pi} \left( \frac{V}{2, \frac{1}{2} e^2} \right)^2 \right) \right) \right] \quad (6)$$

and the slope of  $\frac{d\sigma}{d\Omega}(\theta)$  in the drop-off region (the region of the quarter-point angle) is determined basically by<sup>2,17)</sup>

$$\ell_0 = 2ka \left[ \pi - \text{Arctan} \frac{W}{V} \right] \quad (7)$$

if an equal-geometry for  $W(r)$  and  $V(r)$  is assumed. Therefore the two equations above furnish two invariants for  $\frac{d\sigma}{d\Omega}(\theta)$  and  $\Delta\sigma_R$ , supplying, thus, important constraints on the parameters of the ion-ion interaction, as our calculation clearly indicate

(see Figs. 1 and 2). Though not quite applicable at the low energies we are considering, Eqs. (6) and (7) do supply two reasonable qualitative constraints.

In connection with the deep potential that gave the best account of  $\Delta\sigma_R$ , it is important to stress that exactly this type of potential is seen to emerge from the analysis of intermediate energy data. At these higher energies a remnant of a nuclear rainbow scattering (scattering to negative angles) is seen to occur. At the low energies considered in this letter, our deep potential generates strong orbiting situation, which would persist up to a critical energy given approximately by<sup>17)</sup>

$$E_{cr} = \frac{V_C(R_V)}{2} + \frac{V}{8a_V R_V} \left[ (R_V - 2a_V)^2 + 2a_V^2 \right] \quad (8)$$

where  $V_C(R)$  is the Coulomb interaction at the nuclear potential radius  $R$ . Using the parameters of our potential,  $V = 250$  MeV,  $r_V = 0.66$  fm,  $a_V = 0.63$  fm, we obtain  $E_{cr} = 72$  MeV, well above the energy at which our collected data point end. It would be quite interesting to extend the present study to energies higher than  $E_{cr}$ , where both nuclear rainbow and forward glory would be acting.

In conclusion, we have presented in this letter, strong evidence for the forward glory scattering phenomenon in the  $^{12}\text{C} + ^{12}\text{C}$  system, as exemplified by the enhancement and oscillation in  $\Delta\sigma_R$ . To our knowledge, this is the first time that such a phenomenon has been "seen" in nuclear heavy ion scattering<sup>18)</sup>. We have clearly shown that a joint analysis of both  $\frac{d\sigma}{d\Omega}$  and  $\Delta\sigma_R$  reveals a less ambiguous interaction potential. The  $^{12}\text{C} + ^{12}\text{C}$  potential we obtained from our

analysis is quite deep and resembles closely the interaction potential deduced from analysis done on the elastic scattering of  $^{12}\text{C} + ^{12}\text{C}$  at intermediate energies ( $E \sim \frac{15 \text{ MeV}}{N}$ ) and that calculated from the double folding model<sup>19)</sup>.

#### ACKNOWLEDGEMENTS

We have benefitted from discussions with several colleagues, in particular, M.J. Bechara, L.F. Canto and R. Donangelo. Interesting correspondence and discussion with G.R. Satchler and K.W. McVoy are acknowledged. Finally we thank Y.T. Chen and F. Almeida for help in the numerical calculation.

## REFERENCES

- 1) H.G. Bohlen, M.R. Clover, G. Ingold, H. Lettan and W. von Vertzen, *Z. Phys.* A308, 121 (1982);  
G.R. Satchler, C.B. Fulmer, R.L. Auble, J.B. Ball, F.E. Bertrand, K.A. Erb, E.E. Gross and D.C. Hensley, *Phys. Lett.* B128, 147 (1983);  
S. Kubono et al., *Phys. Lett. B* (1983).  
M.E. Brandan, *Phys. Rev. Lett.* 49, 1132 (1982).
- 2) M.S. Hussein and K.W. McVoy, *Prog. in Particle and Nuclear Physics* (D. Wilkinson, Editor, in press).
- 3) M.S. Hussein, H.M. Nussenzveig, A.C.C. Villari and J.L. Cardoso, *Phys. Lett.* 114B, 1 (1982).
- 4) For an earlier discussion of the optical theorem in the context of heavy-ion scattering see, A.Z. Schwarzchild, E.H. Auerbach, R.C. Fuller and S. Kahana, *Proc. Symp. on Macroscopic Features of Heavy-Ion Collisions*, ANL/PHY-76-2, 753 (1976).
- 5) J.J. Kolata, R.M. Freeman, F. Hass, B. Heusch and A. Gallman, *Phys. Rev.* 21C, 579 (1980).
- 6) M. Conjeaud, S. Gary, S. Harar and J.P. Wielecko, *Nucl. Phys.* 309A, 515 (1978).
- 7) D.G. Kovar et al., *Phys. Rev.* 20C, 305 (1979).
- 8) L.J. Satkowiak, P.A. De Young, J.J. Kolata and M.A. Xapsos, *Phys. Rev.* 26C, 2027 (1982).
- 9) T.M. Cormier et al., *Phys. Rev. Lett.* 40, 924 (1980).
- 10) W. Treu, H. Fröhlich, W. Galster, P. Dück and H. Voit, *Phys. Rev.* 22C, 2462 (1980).
- 11) R.J. Ledoux, M.J. Bechara, C.E. Ordonez, H.A. Al-Juwair and E.R. Cosman, *Phys. Rev.* 27C, 1103 (1983).

- 12) W.A. Friedman and M.A. Bernstein, *Phys. Lett.* 126B, 13 (1983).  
A closed expression for  $\Delta\sigma_R^{S.C.}$ , Eq. (3), can be obtained at higher energies. We find

$$\Delta\sigma_R^{S.C.} \cong \frac{2\sigma_R}{1-\eta^2} \left[ \cos[-2C\eta + 2\eta \ln(l_c + 1/2)] + \eta \sin[-2C\eta + 2\eta \ln(l_c + 1/2)] \right]$$

where  $C = 0.5772156649$  (Euler's constant) and  $\eta$  is the Sommerfeld parameter. The above expression gives the magnitude of  $\Delta\sigma_R^{S.C.}$  quite accurately and tends to the correct limiting value  $2\sigma_R$  at very high energies ( $\eta \rightarrow 0$ ).

- 13) For a discussion of the optical potential ambiguities in heavy ion scattering, see, G.R. Satchler, in: *Proc. Intern. Conf. on Reactions Between Complex Nuclei*, eds. R.L. Robinson et al. (North-Holland, Amsterdam, 1974) 171;  
M. Lozano and G. Madurga, *Nucl. Phys.* A334, 349 (1980).
- 14) W. Reilly et al., *Bull. Am. Phys. Soc.* 16, 10 (1971).
- 15) K.W. Ford and J.A. Wheeler, *Ann. Phys. (NY)* 7, 259 (1959).
- 16) M.S. Hussein, *Phys. Lett.* 127B, 165 (1983).
- 17) J. Knoll and R. Schaeffer, *Ann. Phys. (NY)* 97, 307 (1976).  
See also Ref. 2.
- 18) Forward glory enhancement and oscillations have been discussed previously in atomic collisions, R.B. Bernstein, *J. Chem. Phys.* 34, 361 (1961); E.W. Roth, P.K. Rol, S.M. Trugillo and R.H. Neynaber, *Phys. Rev.* 128, 659 (1962), and in the scattering of light, H.M. Nussenzveig and W.J. Wiscombe, *Opt. Lett.* 5, 455 (1980).
- 19) See, e.g., G.R. Satchler and W.G. Love, *Phys. Lett.* 55C, 183 (1979).

## FIGURE CAPTIONS

FIG. 1 - The quantity  $\Delta\sigma_R$  for the  $^{12}\text{C} + ^{12}\text{C}$  system, calculated with the Yale potential  $V = 14 \text{ MeV}$ ,  $r_v = 1.35 \text{ fm}$ ,  $a_v = 0.35 \text{ fm}$  (dotted curve), the small-radius, deep interaction,  $V = 250 \text{ MeV}$ ,  $r_v = 0.66 \text{ fm}$ ,  $a_v = 0.63 \text{ fm}$  (full curve) and the sharp cut-off limit, Eq. (3) (dashed curve). The data points were extracted from Refs. 5) (open triangles), 6) (open circles), 7) (full circles) and 8) (full triangles), using  $\sigma_{\text{SOD}}$  from ref. 10) and 11).

FIG. 2 - The elastic scattering angular distribution for  $^{12}\text{C} + ^{12}\text{C}$  at three center of mass energies <sup>11)</sup>. The theoretical curves were obtained with the Yale potential (see caption to fig.1) with  $W = 0.4 + 0.1 E_{\text{CM}}$  (MeV),  $r_w = 1.40 \text{ fm}$ ,  $a_w = 0.35 \text{ fm}$  (dashed curve), and with the small radius, deep potential of fig. 1 with  $W = 0.4 + 0.3 E_{\text{CM}}$  (MeV),  $r_w = 0.93 \text{ fm}$  and  $a_w = 0.35 \text{ fm}$  (full curve). The total reaction cross-section obtained with the latter potential is slightly smaller than the experimental value. The values of  $|S_{\ell, \text{gl}}|$  in both cases come out close to unity.




Fig. 1




Fig. 2