

IFUSP/P 476
B.L.F. - USP

UNIVERSIDADE DE SÃO PAULO

INSTITUTO DE FÍSICA
CAIXA POSTAL 20516
01498 - SÃO PAULO - SP
BRASIL

publicações

IFUSP/P-476

NEW WEAK BOSON DECAYS AND POSSIBLE SPIN 3/2 LEPTONS

by

N. Fleury

Centre de Recherches Nucleaires, Université
Louis Pasteur, Strasbourg, France

J. Leite Lopes

Centre de Recherches Nucleaires, Université
Louis Pasteur, Strasbourg, France;
Instituto de Física, Universidade de São Paulo;
Centro Brasileiro de Pesquisas Físicas, Rio de
Janeiro, Brasil

J.A. Martins Simões

Instituto de Física, Universidade Federal do
Rio de Janeiro, Rio de Janeiro, Brasil



29 AGO 1984

Julho/1984

NEW WEAK BOSON DECAYS AND POSSIBLE SPIN 3/2 LEPTONS

N. Fleury

Centre de Recherches Nucleaires, Université
Louis Pasteur, Strasbourg, France;

J. Leite Lopes

Centre de Recherches Nucleaires, Université
Louis Pasteur, Strasbourg, France;
Instituto de Física, Universidade de São Paulo,
São Paulo, Brazil;

and

Centro Brasileiro de Pesquisas Físicas,
Rio de Janeiro, Brasil

J.A. Martins Simões

Instituto de Física, Universidade Federal do Rio de Janeiro,
Rio de Janeiro, Brasil

ABSTRACT

In this note we discuss the possibility that new spin 3/2 leptons can occur in nature as a manifestation of a leptonic structure and in multiplets in supersymmetric theories. We postulate a new interaction between spin 1/2 and spin 3/2 particles that maintains the standard $SU(2)_L \otimes U(1)$ local gauge invariance. A comparison with the anomalous $Z^0 \rightarrow e^+e^-\gamma$ events is made and new decays for the weak bosons are studied.

The repetition of the properties of each of the known fermionic families and the experimental discovery of the tau lepton with a mass higher than that of baryons raises the possibility of a leptonic structure. An evidence for this possibility could be the excitation of the subcomponents and the consequent appearance of spin 3/2 states⁽¹⁾.

In a rather different context supersymmetric theories⁽²⁾ predict supermultiplets with spin 3/2 particles. A recent study⁽³⁾ of the renormalization effects in broken supergravity gauge theories concludes that the gravitino (spin 3/2) can have a mass as low as 50 GeV.

These two points suggest that the experimental discovery of a spin 3/2 lepton (or quark) could be a fundamental step to confirm or to rule out a series of models in the present elementary particle physics.

This is particularly difficult since we have no complete theory for interacting spin 3/2 fields. But we may propose phenomenological interactions which can give qualitative properties of spin 3/2 interactions. This is suggested by the effective point-like Fermi's model which is not a consistent theory but gives very useful informations on weak interactions at low energies.

We will consider mainly the possibility of spin 3/2 states as a manifestation of a leptonic structure for a simple reason. If spin 3/2 fields are excited states of the presently known lepton families we can reasonably suppose that their electroweak interactions are the same as in the low lying spin 1/2 states. This hypothesis fixes almost all the parameters in the theory, except, of course, the spin 3/2 mass.

The uniqueness of the Rarita-Schwinger Lagrangian

has been reexamined recently⁽⁴⁾ and has the form:

$$(1) \quad L_{\text{free}} = -\epsilon^{\alpha\mu\nu\beta} \bar{\psi}_\alpha \gamma^5 \gamma_\mu (\partial_\nu + i \frac{M}{2} \gamma_\nu) \psi_\beta$$

We can generalize⁽⁵⁾ this Lagrangian for the electro-weak $SU_L(2) \otimes U(1)$ model by considering spin 3/2 fields with the same quantum numbers as in the spin 1/2 states: a left-handed doublet

$$(2) \quad L^\mu = \begin{pmatrix} N^\mu \\ E^\mu \end{pmatrix}_L$$

and a right-handed isoscalar

$$(3) \quad R^\mu = E^\mu_R$$

The fermionic Lagrangian is

$$(4) \quad L_{\text{fermionic}} = -\epsilon^{\mu\nu\rho\sigma} (\bar{L}_\mu \gamma_5 \gamma_\nu D_\rho L_\sigma + \bar{R}_\mu \gamma_5 \gamma_\nu D_\rho R_\sigma)$$

where the notation is the same as in reference 6.

We generalize even further the local $SU_L(2) \otimes U(1)$ invariance and include spin 3/2 (L^μ, R^μ) and spin 1/2 (L, R) interactions:

$$(5) \quad L_{3/2, 1/2} = -K \epsilon^{\mu\nu\rho\sigma} \left[\bar{L}_\mu \gamma_5 \gamma_\nu D_\rho D_\sigma L + \bar{R}_\mu \gamma_5 \gamma_\nu D_\rho D_\sigma R \right]$$

where K is a constant which behaves like the inverse of a mass.

The above interactions imply several new decays

for the weak bosons. We compute decay probabilities with arbitrary vector (V) and axial-vector (A) couplings which generalize our interaction Lagrangians (4) and (5) to other cases.

We consider several new decays:

$$\text{DECAY I: } Z^0 \rightarrow E_{3/2}^+ e_{1/2}^-$$

In this case

$$(6) \quad \Gamma(Z^0 \rightarrow E_{3/2}^+ e_{1/2}^-) = \frac{K^2 (V_I^2 + A_I^2)}{48 \pi x (1+x)} M_Z^3 (1-x)^3 (1+2x+3x^2)$$

where $x \equiv M_{3/2}^2 / M_Z^2$. In the standard model we have (for $K = \frac{1}{M_{3/2}}$) the following branching ratio

$$(7) \quad \frac{\Gamma(Z^0 \rightarrow E_{3/2}^+ e_{1/2}^-)}{\Gamma(Z^0 \rightarrow e_{1/2}^+ e_{1/2}^-)} = \frac{1}{4x^2} \frac{(1-x)^3}{(1+x)} (1+2x+3x^2)$$

$$\text{DECAY II: } W^+ \rightarrow E_{3/2}^+ \nu_{1/2} ; W^+ \rightarrow e_{1/2}^+ N_{3/2}$$

For a V-A coupling ($V_{II}^2 = A_{II}^2$) we have ($y = \frac{M_{3/2}^2}{M_W^2}$)

$$(8) \quad \Gamma(W^+ \rightarrow E_{3/2}^+ \nu_{1/2}) = \Gamma(W^+ \rightarrow e_{1/2}^+ N_{3/2}) = \frac{K^2 V_{II}^2 M^3}{24 \pi y (1+y)} (1-y) (1+2y+3y^2)$$

and in the standard model

$$(9) \quad \frac{\Gamma(W^+ \rightarrow E_{3/2}^+ \nu_{1/2})}{\Gamma(W^+ \rightarrow e_{1/2}^+ \nu_{1/2})} = \frac{1}{4y^2} \frac{(1-y)^3}{(1+y)} (1+2y+3y^2)$$

DECAY III: $Z^0 \rightarrow N_{3/2} \bar{\nu}_{1/2}$

For this reaction

$$(10) \quad \Gamma(Z^0 \rightarrow N_{3/2} \bar{\nu}_{1/2}) = \frac{K^2 (A_{III} - V_{III})^2}{96 \pi x (1+x)} M_Z^3 (1-x)^3 (1+2x+3x^2)$$

and the branching ratio is

$$(11) \quad \frac{\Gamma(Z^0 \rightarrow N_{3/2} \bar{\nu}_{1/2})}{\Gamma(Z^0 \rightarrow e_{1/2}^+ e_{1/2}^-)} = \frac{1}{2(a^2 + v^2)} \frac{(1-x)^3}{x^2 (1+x)} (1+2x+3x^2)$$

where $a^2 = 1$ and $v^2 = (1 - 4 \sin^2 \theta_w)$

DECAY IV: $Z^0 \rightarrow N_{3/2} \bar{N}_{3/2}$

This decay is an example of pair production of spin 3/2 fermions. These cases can all be suppressed if $M_{3/2} > \frac{1}{2} M_Z$. The neutral massive pair production gives

$$(12) \quad \Gamma(Z^0 \rightarrow N_{3/2} \bar{N}_{3/2}) = \frac{(V_{IV}^2 + A_{IV}^2)}{144 \pi x} M_Z (1-4x) (3-14x+38x^2)$$

and the branching ratio in the standard model is

$$(13) \quad \frac{\Gamma(Z^0 \rightarrow N_{3/2} \bar{N}_{3/2})}{\Gamma(Z^0 \rightarrow e^+ e^-)} = \frac{8}{3(a^2 + v^2)} \frac{(1-4x)}{x} (3-14x+38x^2)$$

The interaction form (5) implies that the dominant decay of the charged spin 3/2 fermion is electromagnetic


$$(14) \quad E_{3/2} \rightarrow e_{1/2} + \gamma$$

with a branching ratio of practically one. The contributions of this coupling to other processes like the Compton effect or the Thomson cross section were discussed in references 5.

We have also the possibility of neutral decays

$$(15) \quad N_{3/2} \rightarrow \nu_{1/2} + \gamma$$

via higher order contributions like




We turn now our attention to the anomalous $Z^0 \rightarrow e^+ e^- \gamma$ events observed in the UA₁ experiment⁽⁷⁾. Even if we have a limited statistics for these events the agreement with the standard electroweak theory seems very difficult.

In our model, this decay can proceed through the chain

$$Z^0 \rightarrow E_{3/2}^+ e_{1/2}^- \rightarrow e_{1/2}^+ \gamma$$

with a branching ratio of twice our equation 7.

We also have decays like



which are not, so far, detected. A large mass for the spin 3/2 charged lepton can suppress this channel. We compare both predictions in Table I where we consider $M_W = 80$ GeV, $M_Z = 90$ GeV and different values for $M_{3/2}$ (in GeV).

$M_{3/2}$	65	70	75
$\frac{\Gamma(Z^0 \rightarrow E_{3/2}^+ e_{1/2}^-)}{\Gamma(Z^0 \rightarrow e_{1/2}^+ e_{1/2}^-)}$.19	.09	.03
$\frac{\Gamma(W^+ \rightarrow E_{3/2}^+ \nu_{1/2})}{\Gamma(W^+ \rightarrow e_{1/2}^+ \nu_{1/2})}$.05	.01	.001

TABLE I

which shows clearly the suppression of the charged boson decay.

We note that the hypothesis of excited spin 1/2 leptons, with a non-renormalized phenomenological interaction⁽⁸⁾ leads to a mass value similar to the spin 3/2 case to fit the $Z^0 \rightarrow e^+ e^- \gamma$ events.

The decay III also implies a very definite signature.

If the branching ratio for $N_{3/2} \rightarrow \nu_{1/2} \gamma$ is one, we have a decay $Z^0 \rightarrow \nu \bar{\nu} \gamma$ with a branching ratio that can be obtained

from our equation (11) and recognized by a very energetic single gamma event plus missing energy.

If the neutral $N_{3/2}$ satisfies $M_{3/2} < \frac{M_Z}{2}$ decay IV implies events with two energetic gammas and a branching ratio given by equation (13).

In conclusion, production and decay of spin 3/2 leptons appears as a very useful test for leptonic structure and/or supersymmetric models. The present experimental investigation of the W's and Z^0 decays is shown to be consistent with the production of heavy spin 3/2 leptons, but we still need more detailed experiments.

REFERENCES

1. J. Leite Lopes, D. Spehler and J.A. Martins Simões, Phys. Rev. D25, 1854 (1982).
2. J. Wess and J. Bagger: "Supersymmetry and supergravity" - Princeton Univ. Press (1983).
3. J.P. Derendinger and C.A. Savoy - CEN-SACLAY SPHT/83-98, submitted to Nuc. Phys. B.
4. J. Leite Lopes, D. Spehler and N. Fleury, Lett. Nuovo Cimento 35, 60 (1982).
5. A more detailed version is presented by N. Fleury - Thesis - Strasbourg CRN/HE 83-04 and N. Fleury and J. Leite Lopes, Rev. Bras. de Física (dedicated to the Seventieth Birthday of Mario Schönberg).
6. J. Leite Lopes - Gauge Field Theories - Pergamon Press (1981).
7. G. Arnison et al., Phys. Lett. 126B, 398 (1983); 135B, 250 (1983).
P. Bagnair et al., Phys. Lett. 129B, 130 (1983).
8. N. Cabibbo, L. Maiani, Y. Srivastava, Phys. Lett. 139B, 459 (1984).