

IFUSP/P 584
B.L.F. - USE

UNIVERSIDADE DE SÃO PAULO

PUBLICAÇÕES

INSTITUTO DE FÍSICA
CAIXA POSTAL 20516
01498 - SÃO PAULO - SP
BRASIL

IFUSP/P-584
IFT-P03/86

07 JUL 1986



HEAVY LEPTON PRODUCTION VIA VECTOR BOSON FUSION

by

O.J.P. Éboli

Universidade Estadual Paulista "Júlio de Mesquita Filho", Departamento de Física, IGCE
C.P. 178, 13500 Rio Claro, SP, Brasil

G.C. Marques and S.F. Novaes

Instituto de Física, Universidade de São Paulo

A.A. Natale

Instituto de Física Teórica
C.P. 5956, 01405 São Paulo, SP, Brasil

Maio/1986

HEAVY LEPTON PRODUCTION VIA VECTOR BOSON FUSION

O.J.P. Éboli

Universidade Estadual Paulista "Júlio de Mesquita Filho"
Departamento de Física, IGCE
C.P. 178, 13500, Rio Claro, SP, Brasil

G.C. Marques & S.F. Novaes

Universidade de São Paulo
Instituto de Física, Departamento de Física Matemática
C.P. 20516, 01498, São Paulo, SP, Brasil

A.A. Natale

Instituto de Física Teórica
C.P. 5956, 01405, São Paulo, SP, Brasil

ABSTRACT

We investigate the production of a possible pair of sequential heavy charged leptons ($M_L \sim 200-800$ GeV) at pp multi-TeV colliders via vector boson fusion. The results are compared with the usual Drell - Yan mechanism and with the two photon process. For very heavy leptons the fusion of W's provides a contribution that is, at high energies, larger than the D-Y mechanism.

I. INTRODUCTION

Within the current model of the electroweak interaction there is no satisfactory explanation for the existence of several generations of quarks and leptons. On the other hand, there is nothing, in principle, forbidding the existence of at least one more family beyond the usual ones. Since the search for new heavy elementary particles will become an important issue with the advent of multi-TeV hadron colliders, it is interesting to investigate mechanisms that enables the production of such a new family.

In this paper we shall be concerned with the production of heavy charged lepton, belonging to a new generation of sequential leptons (ν_L , L) in the framework of the standard model of electroweak interaction. The charged lepton is assumed to be very heavy, and its neutral partner massless.

The approach used here for studying lepton pair production in pp and $p\bar{p}$ collisions, is reminiscent of the parton model ideas in which the high energy proton (or antiproton) is viewed as a collection of quasi-free partons. For low energies the only relevant partons are the quarks and gluons, and one expects that the Drell-Yan mechanism gives the most relevant contribution. As we increase the energy, however, other particles in the parton sea might play an important role. The treatment of massive vector bosons as constituents of the proton is the basis of the so called effective vector boson approximation. This approximation, as applied to the specific problem we are interested in, has been developed by Kane, Repko and Rolnick¹, and Dawson².

The question that we address ourselves in this paper, is the study of the various vector boson fusion processes contributing to the production of a pair of charged leptons in pp and $p\bar{p}$ collisions. Figure 1 exhibits a prototype of the reactions considered by us. In particular we will be concerned with the possibility that some of these processes dominate over the usual Drell-Yan mechanism ($q\bar{q} \rightarrow \gamma$ or $Z \rightarrow L^+L^-$) and the two photon process ($\gamma\gamma \rightarrow L^+L^-$) at very high energies.

In the next section we introduce the basic expressions for the study of heavy dilepton production, and the cross sections for the elementary processes relevant to our problem. Section III contains our results and discussions.

II. VECTOR BOSON FUSION MECHANISMS FOR MASSIVE LEPTON PAIR PRODUCTION

In order to compute the contributions of the fusion of different vector partons to dilepton production, we shall use the effective vector boson approximation. A basic ingredient in all calculations, within this approximation, are the distribution functions. For massive vector particles, we treat separately the contributions due to transverse and longitudinal polarizations^{1,2}.

The probability distribution of transverse vector bosons in a quark in the high energy limit, i.e., when the quark sub-energy E is much larger than the vector boson mass (M_V), is given by²

$$V_q^T(x) = \frac{(C_V^2 + C_A^2)}{8\pi^2} \frac{[x^2 + 2(1-x)]}{x} \log \left(\frac{4E^2}{M_V^2} \right) \quad (1)$$

whereas, the probability distribution of longitudinal vector bosons in a quark, in the same limit, is

$$V_q^L(x) \cong \frac{(C_V^2 + C_A^2)}{4\pi^2} \cdot \frac{(1-x)}{x} \quad (2)$$

where,

$$C_V = (g/\cos\theta_w) \left(\frac{1}{2} T_{3L} - Q\sin^2\theta_w\right) \text{ and } C_A = -(g/\cos\theta_w) \left(\frac{1}{2} T_{3L}\right)$$

for $V=Z^0$; whereas $C_V = -C_A = g/2\sqrt{2}$ for $V=W^\pm$. T_{3L} is the third component of the weak isospin and Q is the quark electric charge.

The vector boson distribution in a hadron is obtained by the convolution of (1) and (2) with the quark distribution functions in the hadron $q_h(x)$, that is :

$$V_h(x) = \sum_i \int_{x_{min}}^1 \frac{dy}{y} q_h^i(y) V_q^i(x/y) \quad (3)$$

where the sum is carried out over the quark and anti-quark relevant flavours.

We are interested in computing the different vector boson fusion contributions to the rapidity distribution at $y=0$ for the process $h_1 h_2 \rightarrow L^+ L^- + X$ ($h_{1,2} = p(\bar{p})$). This quantity gives a good estimate of the total cross section, because the rapidity distribution is flat in the considered range. In terms of the distribution functions associated to the vector bosons, it can be written as

$$\frac{d\sigma_{h_1 h_2 \rightarrow L^+ L^-}}{dy} \Big|_{y=0} = \int_{\frac{4M_L^2}{S}}^1 d\hat{s} \left[V_{h_1}^a(\sqrt{\hat{s}}) V_{h_2}^b(\sqrt{\hat{s}}) + (a \leftrightarrow b) \right] \hat{\sigma}_{V_a V_b \rightarrow L^+ L^-}(\hat{s}) \quad (4)$$

where $\hat{\sigma}_{V_a V_b \rightarrow L^+ L^-}(\hat{s})$ is the cross section for the subprocesses of Fig.2, $\sqrt{\hat{s}}$ is the total CM subprocess energy and $\tau = \hat{s}/S$.

Since we have already considered the distribution functions we now concentrate on the elementary processes of Fig. 2. We shall consider the production of a heavy charged lepton pair as a result of the following subprocesses

- a) $Z^0 \gamma \rightarrow L^+ L^-$ (Fig. 2(a))
- b) $Z^0 Z^0 \rightarrow L^+ L^-$ (Fig. 2(b))
- c) $W^+ W^- \rightarrow L^+ L^-$ (Fig. 2(c))

where the massive vector bosons are taken transversely and longitudinally polarized.

In the following sections we shall give the results of the cross sections for the subprocesses shown in Fig.2, for massive vector bosons transversely and longitudinally polarized. In all calculations we have assumed $\hat{s} \gg M_V^2$ and $M_L \gg M_V$.

A. $Z^0 \gamma$ fusion

At first, we consider the subprocess $Z^0 \gamma \rightarrow L^+ L^-$ of Fig.2(a). The cross section for the transversely polarized Z^0 is

$$\hat{\sigma}(Z_T^0 \gamma \rightarrow L^+ L^-) = \frac{\pi \alpha^2}{4 \sin^2 \theta_w \cos^2 \theta_w} \cdot \frac{\beta}{\hat{s}}$$

$$\cdot \left\{ (1 - 4 \sin^2 \theta_w)^2 \left[-\beta^2 + \frac{(1 + \beta^4)}{2\beta} L \right] - 1 + 3(1 - \beta^2) + \frac{(3\beta^4 - 1)}{2\beta} L \right\} \quad (5)$$

where :

$$\beta = \left(1 - 4 \frac{M_L^2}{\hat{s}} \right)^{1/2} \quad (6)$$

is the lepton velocity in the CM frame, and :

$$L \equiv \log \frac{(1 + \beta)}{(1 - \beta)} \quad (7)$$

The cross section for the longitudinally polarized Z^0 is

$$\hat{\sigma}(Z_L^0 \gamma \rightarrow L^+ L^-) = \frac{\pi \alpha^2}{\sin^2 \theta_w \cos^2 \theta_w} (1 - 4 \sin^2 \theta_w)^2 \frac{\beta}{\hat{s}} \frac{M_L^2}{M_Z^2} \left(-2 + \frac{1}{\beta} L \right) \quad (8)$$

B. $Z^0 Z^0$ fusion

The diagrams contributing to $Z^0 Z^0$ fusion are shown in Fig. 2(b), where H is the standard Higgs boson. For transversely polarized Z^0 bosons the cross section is :

$$\hat{\sigma}(Z_T^0 Z_T^0 \rightarrow L^+ L^-) = \frac{\pi \alpha^2}{64 \sin^4 \theta_w \cos^4 \theta_w} \frac{\beta}{\hat{s}}$$

$$\left\{ (1 - 4 \sin^2 \theta_w)^4 \left[-1 + 3(1 - \beta^2) + \frac{[2\beta^2 - (1 - \beta^2)^2]}{2\beta} L \right] - \right.$$

$$\left. - 2(1 - 4 \sin^2 \theta_w)^2 \left[6 + \beta^2 - \frac{[6 + (1 - \beta^2)^2]}{2\beta} L + 4 M_L^2 \beta \chi_H (\hat{s} - M_H^2) L \right] - \right.$$

$$\left. - 1 - (1 - \beta^2) + \frac{(3 - \beta^4)}{2\beta} L + 8 M_L^2 \beta \chi_H (\hat{s} - M_H^2) L + \right.$$

$$\left. + 16 M_L^2 \beta^2 \chi_H \hat{s} \right\} \quad (9)$$

where,

$$\chi_H = \frac{1}{(\hat{s} - M_H^2)^2 + \Gamma_H^2 M_H^2} \quad (10)$$

M_H and Γ_H being the Higgs mass and width respectively.

When the Z^0 bosons are longitudinally polarized, we have

$$\hat{\sigma}(Z_L^0 Z_L^0 \rightarrow L^+ L^-) = \frac{\pi \alpha^2}{8 \sin^4 \theta_w \cos^4 \theta_w} \frac{M_L^2}{M_Z^4} \beta \cdot$$

$$\left\{ (1-4\sin^2 \theta_w)^4 - 2(1-4\sin^2 \theta_w)^2 \chi_H \hat{s}(\hat{s}-M_H^2) + \beta^2 \chi_H \hat{s}^2 - \right.$$

$$\left. - \frac{(1-\beta^2)}{2\beta} \left[(1-4\sin^2 \theta_w)^4 - 2(1-4\sin^2 \theta_w)^2 \chi_H \hat{s}(\hat{s}-M_H^2) \right] \cdot \mathcal{L} \right\} \quad (11)$$

We do not worry about cross sections involving mixed polarizations (TL), since σ_{TT} and σ_{LL} are enough to characterize the complete behaviour of the process.

C. $W^+ W^-$ fusion

We shall consider here the possibility of the heavy lepton pair to be produced via the fusion of charged vector bosons (Fig.2(c)). For transversely polarized W bosons the cross section is :

$$\hat{\sigma}(W_T^+ W_T^- \rightarrow L^+ L^-) = \sum_{i,j} \hat{\sigma}_{ij}^T \quad (12)$$

where $\hat{\sigma}_{ij}$ ($i, j = A, Z, H, \nu$) are the contribution of each diagram of Fig.2(c) and its relevant interferences. We get for these contributions:

$$\hat{\sigma}_{AA}^T = \frac{\pi \alpha^2}{3 \hat{s}} \beta (3-\beta^2),$$

$$\hat{\sigma}_{ZZ}^T = \frac{\pi \alpha^2}{48 \sin^4 \theta_w} \frac{\beta}{\hat{s}} \left[(1-4\sin^2 \theta_w)^2 (3-\beta^2) + 2\beta^2 \right],$$

$$\hat{\sigma}_{HH}^T = \frac{\pi \alpha^2}{4 \sin^4 \theta_w} M_L^2 \beta^3 \chi_H,$$

$$\hat{\sigma}_{\nu\nu}^T = \frac{\pi \alpha^2}{8 \sin^4 \theta_w} \frac{\beta}{\hat{s}} \left\{ -\frac{2}{3} (3+\beta^2) + (1+\beta^2)(1-2\beta^2) + \right.$$

$$\left. + \left[2(3+\beta^2) - (1-\beta^4)(1+2\beta^2) \right] \frac{\mathcal{L}}{2\beta} \right\},$$

Whereas for the W's longitudinally polarized, we have

$$\hat{\sigma}_{AZ}^T = \frac{\pi \alpha^2}{6 \sin^2 \theta_W} (1 - 4 \sin^2 \theta_W) \frac{\beta}{\hat{s}} (3 - \beta^2),$$

$$\hat{\sigma}_{(W_L^+ W_L^- \rightarrow L^+ L^-)} = \sum_{ij} \hat{\sigma}_{ij}^L, \quad (14)$$

where :

$$\hat{\sigma}_{HV}^T = \frac{\pi \alpha^2}{4 \sin^4 \theta_W} M_W^2 \chi_H \frac{(\hat{s} - M_H^2) \beta (1 - \beta^2)}{3} \left[-1 + \frac{(1 + \beta^2)}{2\beta} \mathcal{L} \right],$$

$$\hat{\sigma}_{AA}^L = \frac{\pi \alpha^2}{6} \cdot \frac{\hat{s}}{M_W^4} \beta (3 - \beta^2),$$

$$\hat{\sigma}_{AV}^T = \frac{-\pi \alpha^2}{8 \sin^2 \theta_W} \cdot \frac{\beta}{\hat{s}} \left\{ \frac{4}{3} (3 - \beta^2) + (1 - \beta^4) + \frac{(1 - \beta^2)}{2\beta} [3 - \beta^2(2 + \beta^2)] \mathcal{L} \right\},$$

$$\hat{\sigma}_{ZZ}^L = \frac{\pi \alpha^2}{96 \sin^4 \theta_W} \cdot \frac{\hat{s}}{M_W^4} \beta \left[(1 - 4 \sin^2 \theta_W)^2 (3 - \beta^2) + 2\beta^2 \right],$$

$$\hat{\sigma}_{ZV}^T = \frac{-\pi \alpha^2}{32 \sin^4 \theta_W} \cdot \frac{\beta}{\hat{s}}.$$

$$\hat{\sigma}_{HH}^L = \frac{\pi \alpha^2}{8 \sin^4 \theta_W} \frac{M_L^2}{M_W^4} \chi_H \hat{s}^2 \beta^3,$$

$$\left\{ (1 - 4 \sin^2 \theta_W) \left[\frac{4}{3} (3 - \beta^2) + (1 - \beta^4) + \frac{(1 - \beta^2)}{2\beta} [3 - \beta^2(2 + \beta^2)] \mathcal{L} \right] + \frac{8}{3} \beta^2 + (1 - \beta^4) - \frac{(1 - \beta^2)^3}{2\beta} \mathcal{L} \right\} \quad (13)$$

$$\hat{\sigma}_{\nu\nu} = \frac{\pi \alpha^2}{64 \sin^4 \theta_W} \cdot \frac{\hat{s}}{M_W^4} \beta \left[-\frac{4}{3} (6 + \beta^2) + 3(1 + \beta^2)(3 - \beta^2) - (1 - \beta^2)^2 + 2\beta(1 - \beta^2)^2 \mathcal{L} \right],$$

$$\hat{\sigma}_{AZ}^L = \frac{\pi\alpha^2}{12\sin^2\theta_W} (1-4\sin^2\theta_W) \frac{\hat{s}}{M_W^4} \beta (3-\beta^2),$$

$$\hat{\sigma}_{HV}^L = \frac{\pi\alpha^2}{8\sin^4\theta_W} \frac{M_L^2}{M_W^4} \chi_H \hat{s} (\hat{s}-M_H^2) \beta \left[-(1+\beta^2) + \frac{(1-\beta^2)^2}{2\beta} \mathcal{L} \right],$$

$$\hat{\sigma}_{AV}^L = \frac{\pi\alpha^2}{8\sin^2\theta_W} \frac{\hat{s}}{M_W^4} \beta \left[-\frac{4}{3}\beta^2 - \frac{(1-\beta^4)}{2} + \frac{(1-\beta^2)^3}{4\beta} \mathcal{L} \right],$$

$$\hat{\sigma}_{ZV}^L = \frac{\pi\alpha^2}{16\sin^4\theta_W} (1-2\sin^2\theta_W) \frac{\hat{s}}{M_W^4} \beta.$$

$$\left[-\frac{4}{3}\beta^2 - \frac{(1-\beta^4)}{2} + \frac{(1-\beta^2)^3}{4\beta} \mathcal{L} \right] \quad (15)$$

There are terms in (15) which exhibit a growing with \hat{s} violating unitarity bounds. As expected, they cancel among themselves when we take the sum over these contributions in (14).

III. RESULTS AND CONCLUSIONS

The various contributions, due to vector boson fusion, to the rapidity distribution at $y = 0$, can be inferred from (1-4) and from the cross sections of each subprocess computed in the previous section. In the numerical calculations we have used $M_W = 82.5$ GeV and $M_Z = 93.8$ GeV for the masses of the intermediate vector bosons. For the weak mixing angle we have assumed $\sin^2\theta_W = 0.226$. As for the Higgs boson mass we have taken two different set of data, fixing the Higgs boson mass at twice and eight times the mass of Z^0 . For the Q^2 -dependent quark distribution functions we have taken the set 2 given in the appendix of ref.3. For the photon distribution in a quark we have used the Weizsäcker-Williams approximation, which yields:

$$f_q^{\gamma}(x) = \frac{Q^2\alpha}{2\pi} \frac{[x^2 + 2(1-x)]}{x} \log\left(\frac{E^2}{m_q^2}\right)$$

where, m_q is the quark mass. Our numerical results are summarized in Fig. 3-10. Figure 3 shows the contribution of the subprocess 2(a) to heavy lepton pair production in pp collisions at different collider energies. In this case we would like to stress the dominance of the transversely polarized Z^0 boson cross section over the longitudinally polarized one. We credit this dominance to helicity suppression. These cross sections, however, do not give any significant contribution to heavy lepton pair production, since they are three orders of magnitude smaller than the one associated to the Drell-Yan mechanism ($q\bar{q}$ fusion).

In Fig. 4 and 5, we show the rapidity distribution at $y = 0$ due to $Z^0 Z^0$ fusion for different values of the Higgs boson mass. We note that the transverse contribution is smaller than the longitudinal one and is more sensitive to the lepton mass. The $Z_T^0 Z_T^0$ contribution behaves similarly to the two photon process but, due to the large Z^0 mass, it lies one order of magnitude below the two photon contribution ($\gamma\gamma$ fusion). It is interesting to notice in Fig.5 the existence of bumps centered at $M_L^2 = \frac{1}{10} M_H^2$, which is the value that maximizes the width of the Higgs boson in lepton pair channel.

Figure 6 shows the contribution to heavy lepton pair production due to the fusion of longitudinally and transversely polarized W^+W^- bosons (Fig. 2(c)) in proton-proton collision for $M_H = 2M_Z$. The transverse contribution is roughly independent of the Higgs boson mass and, for this reason, we draw in Fig.7 only the longitudinal contribution for $M_H=8M_Z$.

The relevance of the $W_L^+W_L^-$ fusion can be inferred by analyzing Fig. 8-10 for $\sqrt{s} = 20, 40$ and 70 TeV. We have also exhibited, for the sake of completeness, the contribution due to $q\bar{q}$ fusion (Drell-Yan mechanism) and $\gamma\gamma$ fusion. From these results it can be inferred that although the Drell-Yan mechanism is the most relevant one for "light leptons", the longitudinal W boson fusion takes over the D-Y mechanism for heavy leptons ($M_L \gg 600$ GeV at $\sqrt{s} = 40$ TeV, whereas $M_L \gg 500$ GeV at $\sqrt{s} = 70$ TeV).

We have verified that all the above conclusions remain valid when we consider proton-antiproton instead of proton-proton collisions.

It has been pointed out in Ref.2 that the effective W approximation, used throughout this paper, is reliable only when the scattering amplitude is dominated by one of the modes (longitudinal or transverse) in a clear way. Hence, for processes satisfying the above criterion, we do not need to worry about the interference between transverse and longitudinal bosons inside a proton. This condition is met for all processes we analyzed, except in the case of the fusion of W's for lepton masses smaller than ~ 300 GeV. Thus the use of the effective vector boson approximation is not reliable in this situation.

The effective vector boson approximation provides quite good results. In fact for the production of heavy Higgs boson, Cahn⁴ has shown that using the effective vector boson approximation, the cross section is 5 - 30% larger than the correspondent exact result.

Recently, Willenbrock and Dicus⁵ analyzed the gluon fusion mechanism for heavy lepton production, which is particularly enhanced if, as they assumed, there is a contribution of a loop of a new heavy quark. Their proposal entails a cross section roughly one order of magnitude larger than the $W_L^+W_L^-$ fusion mechanism. Nevertheless, we expect that for the production of a heavy lepton (L) and its neutral partner (ν_L) the vector boson fusion studied here will be the most important mechanism since in this case there will not be competition from the gluon-gluon fusion process. This reaction is now under investigation by us.

ACKNOWLEDGMENTS

We wish to thank R.Aldrovandi and V.Pleitez for enlightening discussions and C.O.Escobar for the critical reading of the manuscript. We are indebted to Fundação de Amparo à Pesquisa do Estado de São Paulo, FAPESP (S.F.N.), Conselho Nacional de Desenvolvimento Científico e Tecnológico, CNPq, and Financiadora de Estudos e Projetos, FINEP (G.C.M. and A.A.N.) for their financial support. We are grateful to the Instituto de Física Teórica (S.F.N.) and Departamento de Física Matemática da USP (O.J.P.E.) for their kind hospitality.

REFERENCES

- (1) G.L.Kane, W.W.Repko and W.B.Rolnick
Phys. Lett. 148B, 367 (1984)
- (2) Sally Dawson
Nucl.Phys. B249, 42 (1985)
- (3) E.Eichten, I.Hinchliffe, K.Lane and C.Quigg
Rev.Mod.Phys. 56, 579 (1984)
- (4) R.N.Cahn
Nucl.Phys. B255, 341 (1985)
- (5) S.S.D.Willenbrock and D.A.Dicus
Phys.Lett. 156B, 429 (1985)

FIGURE CAPTIONS

Fig. 1 : Lepton pair production via vector boson fusion

Fig. 2 : Subprocesses that contribute to the heavy lepton pair production : (a) $Z^0\gamma \rightarrow L^+L^-$, (b) $Z^0Z^0 \rightarrow L^+L^-$ and (c) $W^+W^- \rightarrow L^+L^-$.

Fig. 3 : Rapidity distribution at $y=0$ for $Z^0\gamma$ contribution to heavy lepton pair production in pp collisions for the massive vector boson transversely polarized (upper curves) and longitudinally polarized (lower curves) at $\sqrt{s} = 20$ TeV (dotted lines), 40 TeV (solid lines) and 70 TeV (dashed lines).

Fig. 4 : Rapidity distribution at $y=0$ for Z^0Z^0 contribution to heavy lepton pair production in pp collisions for both vector bosons longitudinally polarized (upper curves) and transversely polarized (lower curves) for $M_H = 2M_Z$. The energies are the same as for Fig.3.

Fig. 5 : The same as Fig.4 for $M_H = 8M_Z$.

Fig. 6 : Rapidity distribution at $y=0$ for W^+W^- contribution to heavy lepton pair production in pp collisions for both vector bosons longitudinally polarized (upper curves) and transversely polarized (lower curves) for $M_H = 2M_Z$. The energies are the same as for Fig.3.

Fig. 7 : Rapidity distribution at $y=0$ for W^+W^- contribution to heavy lepton pair production in pp collisions for both vector bosons longitudinally polarized and for $M_H = 8M_Z$. The energies are the same as for Fig.3.

Fig. 8 : Rapidity distribution at $y=0$ for Drell-Yan ($q\bar{q}$) (dashed line), two gamma fusion ($\gamma\gamma$) (dotted line) and two longitudinally polarized W bosons fusion (solid line) ($M_H = 2M_Z$) contributions to heavy lepton pair production in pp collisions at $\sqrt{s} = 20$ TeV.

Fig. 9 : The same as Fig.8 at $\sqrt{s} = 40$ TeV.

Fig.10 : The same as Fig.8 at $\sqrt{s} = 70$ TeV.







FIG. 1



(a)



(b)



(c)

FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10