

IFUSP/P846

B.I.F. - USP

**UNIVERSIDADE DE SÃO PAULO**

**INSTITUTO DE FÍSICA  
CAIXA POSTAL 20516  
01498 - SÃO PAULO - SP  
BRASIL**

# **PUBLICAÇÕES**

IFUSP/P-846



**CONSTRAINED GOE FOR SYSTEMS WITH FEW  
DEGREES OF FREEDOM IN THE INTERMEDIATE  
REGIME BETWEEN CHAOS AND ORDER**

**C.E. Carneiro, M.S. Hussein and M.P. Pato**  
Instituto de Física, Universidade de São Paulo

Maio/1990

2

# CONSTRAINED GOE FOR SYSTEMS WITH FEW DEGREES OF FREEDOM IN THE INTERMEDIATE REGIME BETWEEN CHAOS AND ORDER\*

C.E. Carneiro, M.S. Hussein and M.P. Pato

Instituto de Física, Universidade de São Paulo  
C.P. 20516, 01498 São Paulo, S.P., Brazil

## ABSTRACT

New distribution laws for the energy level spacings and the eigenvector amplitudes, appropriate for systems with a few degrees of freedom in the intermediate regime between chaos and order, are derived by conveniently deforming the Gaussian Orthogonal Ensemble. The cases of  $2 \times 2$  and  $3 \times 3$  matrices are fully worked out. The general case of matrices with large dimensions are discussed. The Hubbard–Stratonovich transformation in conjunction with the Method of Integration over Alternate Variables are employed for the purpose.

\*Supported in part by the CNPq.

May/1990

## I. INTRODUCTION

It is well known that the GOE (Gaussian Orthogonal Ensemble) provides a realistic description of statistical properties of complex quantal systems with a large number of degrees of freedom<sup>1)</sup>. In particular, the energy level spacings distribution obtained from GOE approximates very well the Wigner distribution<sup>2)</sup> (W), which describes quite well data. Independently of the spacing distribution, the GOE predicts a Porter–Thomas (PT) law<sup>3)</sup> for the distributions of eigenvectors. Nuclear resonance data systems lends support to the correctness of the PT law<sup>1)</sup>.

The statistical independence of the eigenvector and eigenvalue distributions, a characteristic of the GOE, must however, be revised for cases involving systems with a few degrees of freedom in the chaotic regime<sup>4)</sup>.

The purpose of this paper is to develop a theory for a mixed eigenvector and eigenvalue distribution which should describe situations intermediate between chaos and order. For this purpose we employ the maximum entropy technique subjected to several physically motivated constraints<sup>4)</sup>. The GOE results will always be used as a guideline. The case of  $2 \times 2$  matrices are worked out fully analytically while the  $3 \times 3$  matrices are developed semi-analytically supplemented by numerical evaluation. The general case is then discussed using the Hubbard–Stratonovich transformation in conjunction with the Alternate Variable Method of Integration (AVMI) development by Mehta<sup>5)</sup>. A shorter version of this work is contained in Ref. 6).

## II. THE GAUSSIAN ORTHOGONAL ENSEMBLE DISTRIBUTION FROM THE MAXIMUM ENTROPY PRINCIPLE

Using the maximum entropy technique the usual GOE distribution is obtained as follows. First we define the entropy as

$$S = - \int dHP(H) \ln P(H) , \quad (1)$$

subject to the constraint

$$\langle \text{Tr } H^2 \rangle = \int dHP(H) \text{Tr } H^2 = \mu \quad (2)$$

and the normalization condition

$$\int dHP(H) = 1 \quad (3)$$

Maximizing  $S$  subject to (2) and (3) one immediately obtains the GOE distribution

$$P(H) = e^{-\lambda_0 - 1 - \alpha_0 \text{tr} H^2} \quad (4)$$

$$\alpha_0 = \frac{N(N+1)}{4\mu} , \quad e^{-\lambda_0 - 1} = 2^{-\frac{N}{2}} \left[ \frac{\pi}{2\alpha_0} \right]^{-\frac{N(N+1)}{4}}$$

where  $\lambda_0$  and  $\alpha_0$  are Lagrange multipliers determined by the constraints (2) and (3).

From (4) one can construct the joint distribution function,  $P(E_1, \dots, E_N; C_1, \dots, C_N)$  using already well known procedure<sup>5),6)</sup>, and obtain

$$P^{(\text{GOE})}(E_1, \dots, E_N; C_1, \dots, C_N) = 2 \left[ \frac{\pi^{N/2}}{N} \right]^{-1} \delta \left[ 1 - C_1^2 - C_2^2 - C_3^2 - \dots - C_N^2 \right] \cdot \left[ N! 2^{-1/4N(N-1)} \prod_{g=1}^N \Gamma\left(\frac{1}{2}g\right) \right]^{-1} \prod_{i < j} |E_i - E_j| e^{-\alpha(E_1^2 + E_2^2 + \dots + E_N^2)} \quad (5)$$

where the  $C_i$ s are the component amplitudes of an eigenvector.

Integrating (5) over all the  $C_i$ s and all the  $E_i - E_j$ s, except one  $s$ , one obtains in the  $N \rightarrow \infty$  limit the Wigner distribution

$$P_W(s) \cong \frac{\pi}{2D^2} s \exp \left[ -\frac{\pi s^2}{4D^2} \right] \quad (6)$$

where  $D$  is  $\langle s \rangle = \left[ \frac{8\pi}{\alpha} \right]^{1/2}$

Similarly integrating all the  $E_i$ s and all the  $C_i$ s except one  $c$ , one finds in the limit  $N \rightarrow \infty$  limit the Porter-Thomas distribution

$$P_{\text{PT}}(c) \cong \left[ \frac{N}{2\pi} \right]^{1/2} \exp \left[ -\frac{N}{2} C^2 \right] \quad (7)$$

As seen from 5, the  $E$  and  $C$  distributions are completely independent,  $(P(E,s;C,s) = P(E,s) P(C,s))$ , which is a consequence of GOE, namely  $P(H)$  is invariant under arbitrary rotation of the basis.

Before turning our attention to the constrained GOE, it is helpful to mention that the constraint (2) above gives rise to the Gaussian distribution (4) with the second moment

$$\langle H_{ij} H_{kl} \rangle = (\delta_{ik} \delta_{jl} + \delta_{il} \delta_{jk}) \frac{1}{4\alpha_0} \quad (8)$$

with  $\alpha_0$  being independent of the label.

### III. THE DEFORMED GOE

To generalize (4) and thus eventually make the joint distribution non-separable, we have to impose further constraints in the construction of  $P(H)$  from the maximum entropy principle. We thus redo the calculation with Eq. (1) with the additional condition

$$\langle \text{Tr}(PHQHP) \rangle = \nu, \quad (9)$$

where  $P$  is a projection operator that projects onto a given vector in the basis space, e.g.  $P = |i\rangle\langle i|$ , and  $Q = 1 - P$ .

To understand the meaning of this constraint we write the trace in terms of the elements of the Hamiltonian matrix, namely

$$\text{Tr}(PHQHP) = \sum_{j \neq i} H_{ij}^2, \quad (10)$$

which, when expressed in terms of the eigenvalues  $E_k$ , and the components  $C_k^i = \langle i | E_k \rangle$  of the eigenvectors  $|E_k\rangle$ , becomes

$$\text{Tr}(PHQHP) = \sum_k (C_k^i)^2 E_k^2 - \sum_{k, \ell} (C_k^i)^2 (C_\ell^i)^2 E_k^2 E_\ell^2, \quad (11)$$

where the completeness relation  $\sum_i |i\rangle\langle i| = 1$ , has been used. Therefore the new constraint, Eq. (8), with  $P = |i\rangle\langle i|$  involves all the eigenvalues of  $H$  and the projections of the corresponding eigenvectors on the basis state  $|i\rangle$ .

By fixing the value of  $\langle \text{Tr}(PHQHP) \rangle$ , we are therefore deforming the Gaussian Orthogonal Ensemble. Of course, the system still maintains full axial symmetry about the  $|i\rangle$  "direction". The new ensemble is invariant under transformation which leaves the vector  $|i\rangle$  unchanged. Further understanding of the new constraint, Eq. (9) can be gained by realizing that the second moment is now given by

$$\langle H_{ij} H_{kl} \rangle = (\delta_{ik} \delta_{jl} + \delta_{il} \delta_{jk}) \frac{1}{4\alpha + 2\beta |\delta_{ki} - \delta_{li}|} \quad (12)$$

The new distribution function  $P(H)$  constructed from

$$\delta \int P \left[ -\ln P - \lambda - \alpha \text{Tr} H^2 - \beta \text{Tr}(PHQHP) \right] dH = 0 \quad (13)$$

is thus given by

$$\begin{aligned} P(H) &= \exp \left[ -\lambda - 1 - \alpha \text{Tr} H^2 - \beta \text{Tr}(PHQHP) \right] \\ &= P^{\text{GOE}(H)} e^{-\beta \text{Tr}(PHQHP)} \cdot \left[ 1 + \frac{\beta}{2\alpha} \right]^{\frac{N-1}{2}}, \end{aligned} \quad (14)$$

where the Lagrange multipliers  $\alpha$  and  $\beta$  are related to the  $\mu$  and  $\nu$  as follows

$$\nu = \frac{N-1}{4\alpha \left( 1 + \frac{\beta}{2\alpha} \right)} \equiv \langle \text{Tr} PHQHP \rangle \quad (14.a)$$

$$\mu = \frac{N(N+1)}{4\alpha} - \frac{N-1}{\left(1 + \frac{\beta}{2\alpha}\right)} \frac{\beta}{4\alpha^2} \equiv \langle \text{Tr } H^2 \rangle \quad (14.b)$$

The symmetry inherent in the GOE is now broken because of the favoring of a particular vector spanned by  $P$ . This guarantees that the joint distribution  $P(E_1, \dots, E_N; C_1, \dots, C_N)$  is not any more separable, and accordingly significant deviations from both the Wigner and the Porter-Thomas distributions are expected. Since the Jacobian that takes us from  $P(H)$  to  $P(E, s; C, s)$  is independent of the particular ensemble of real matrices under consideration we conjecture that the joint probability distribution corresponding to Eq. (14) is

$$P(E, s; C, s) = P^{(\text{GOE})}(E, s; C, s) \left[1 + \frac{\beta}{2\alpha}\right]^{\frac{N-1}{2}} \cdot \exp\left[-\beta \left[ \sum_k (C_k^1)^2 E_k^2 - \sum_{k, \ell} (C_k^1)^2 (C_\ell^1)^2 E_k^2 E_\ell^2 \right]\right] \quad (15)$$

where  $\text{Tr}(PHQHP)$  has been represented explicitly in terms of the eigenvalues and the amplitudes of the eigenvector which is projected by  $P$ . From the joint distribution, Eq. (15) one can then calculate the spacing and amplitude distributions upon integration. In the following we discuss the cases of  $2 \times 2$  and  $3 \times 3$  matrices.

#### IV. THE DISTRIBUTION $P(s)$ AND $P(c)$ IN TWO AND THREE DIMENSIONS

In this Section we derive the distribution functions,  $P(s)$  and  $P(c)$  for  $2 \times 2$  and  $3 \times 3$  matrices. For  $2 \times 2$  matrices we obtain

$$P(s) = \alpha \left[1 + \frac{\beta}{2\alpha}\right]^{1/2} \cdot s \exp\left[-\left[\frac{\alpha}{2} + \frac{\beta}{2\alpha}\right] s^2\right] I_0\left[\frac{\beta}{8} s^2\right] \quad (16)$$

and

$$P(c) = \frac{\alpha}{\pi} \left[1 + \frac{\beta}{2\alpha}\right]^{1/2} \frac{1}{\sqrt{1-C^2}} \frac{1}{\left[\frac{\alpha}{2} + \beta C^2(1-C^2)\right]} \quad (17)$$

Eq. (16) represents a spacing distribution which still shows level repulsion but deviates from a pure Wigner one. It is interesting to note that when  $\beta = 0$ , the Wigner surmise is fully recovered. On the other hand,  $P(c)$ , Eq. (17), is necessarily different from the Porter-Thomas distribution even when  $\beta = 0$ , owing to the small dimension of the matrices. For  $3 \times 3$  matrices the spacing and amplitude distributions are given by

$$P(s) = \frac{8\alpha^3}{3\pi^2} \sqrt{\frac{6\pi}{\alpha}} \left[1 + \frac{\beta}{2\alpha}\right] s \exp\left[-\frac{2}{3} \alpha s^2\right] \cdot \int_0^1 dc \exp\left[-\beta(1-C^2) C^2 s^2\right] \int_0^{\pi/2} \frac{d\psi}{Q} \left\{ \frac{2B^2 s^2 + Q(1-2Bs^2)}{4Q^{3/2}} \sqrt{\pi} \exp\left[B^2 \frac{s^2}{C}\right] \cdot \text{erfc}\left[\frac{Bs}{\sqrt{Q}}\right] + s \frac{(Q-B)}{2Q} \right\} \quad (18)$$

and

$$P(c) = \frac{\alpha^3}{3\pi \sqrt{6\alpha}} \left[1 + \frac{\beta}{2\alpha}\right] \int_0^{\pi/2} d\psi \left[ \frac{\frac{1}{\sqrt{A_1}} + \frac{1}{\sqrt{Q}}}{(\sqrt{A_1 Q} + B_1)^2} + 2 \frac{\frac{1}{\sqrt{A}} + \frac{1}{\sqrt{Q}}}{(\sqrt{AQ} + B)^2} \right] \quad (19)$$

where

$$\begin{aligned} A &= \frac{2}{3} \alpha + \beta(1-C^2) C^2, & A_1 &= \frac{2}{3} \alpha + \beta(1-C^2-C'^2)(C^2+C'^2) \\ B &= \frac{\alpha}{3} + \beta C^2 C'^2, & A_2 &= \frac{\alpha}{3} + \beta C'^2(1-C^2-C'^2) \\ Q &= \frac{2}{3} \alpha + \beta(1-C'^2) C'^2 \\ \psi &= \sin^{-1} \frac{C'}{\sqrt{1-C^2}} \end{aligned}$$

In Figures 1, 2 and 3 we show our results for  $P(s)$ ,  $P(c)$  and  $P(y \equiv c^2)$  for the  $2 \times 2$  and  $3 \times 3$  matrices cases discussed above. An important result which deserves a special attention is that the spacing distribution  $P(s)$  for small  $s$  dives to zero as fast as  $\left[0.36 \left[1 + \frac{\beta}{2\alpha}\right]^{1/2} + 0.64\right]s$  for  $3 \times 3$  matrices and as  $\left[1 + \frac{\beta}{2\alpha}\right]^{1/2} \alpha s$  for  $2 \times 2$  matrices, as  $\beta$  is increased further ( $\alpha$  was kept constant = 1). In the limit  $\beta \rightarrow \infty$ , which should represent the regular (non-chaotic) limit, the  $P(s)$  and  $P(c)$  distributions attain the following form:

$$P_{\infty}(s) = \sqrt{\frac{2\alpha}{\pi}} \exp\left[-\frac{\alpha s^2}{2}\right], \quad N = 2 \quad (20)$$

$$P_{\infty}(s) = 2 \sqrt{\frac{3\alpha}{8\pi}} \left\{1 + \sqrt{\frac{\pi\alpha}{6}} s e^{\alpha s^2/6} \operatorname{erfc}\left[\sqrt{\frac{\alpha}{6}} s\right]\right\} \exp\left[-\frac{2\alpha}{3} s^2\right], \quad N = 3 \quad (21)$$

$$P_{\infty}(c) = \frac{1}{2} \left[\delta(c) + \delta(c-1) + \delta(c+1)\right], \quad N = 2 \quad (22)$$

$$P_{\infty}(c) = \frac{1}{4} \left[3\delta(c) + \delta(c-1) + \delta(c+1)\right], \quad N = 3 \quad (23)$$

Our results above for  $N=2$  agree fully with those of Robnik<sup>7)</sup> and Alhassid and Levine<sup>8)</sup> and disagree with Berry and Robnik<sup>9)</sup> who claim to have seen no level repulsion even for moderately small values of  $\beta$  (the chaoticity parameter is defined differently in Ref. 9)). At this point we would like to point out that our work is very similar in spirit to that of Alhassid and Levine. In fact, the case fully calculated by Ref. 8), namely the  $2 \times 2$  matrix, agree completely with our Eqs. (16) and (17) with the identification of their parameter  $\varepsilon$  by the following relation

$$\varepsilon = \frac{1}{\sqrt{1 + \frac{\beta}{2\alpha}}} \quad (24)$$

Thus the GOE limit obtains by setting  $\beta = 0$  or  $\varepsilon = 1$ . Similarly the fully regular case corresponds to  $\beta = \infty$  or  $\varepsilon = 0$ .

We should, however, point out two major advantages that our theory carries in relation to Alhassid and Levine's namely, first, we deform the GOE by merely favoring one particular vector in the basis space spanned by  $P$  and accordingly introduce just one more Lagrange multiplier in the maximum entropy calculation, whereas they introduce  $N-1$  Lagrange multipliers to go beyond the GOE. Secondly, our chaotic parameter,  $\beta$ , could take the full range of values from 0 (fully chaotic) to  $\infty$  (fully regular). Thus, interpreted as an inverse of a "temperature", one may use knowledge in statistical mechanical studies of phase transition phenomena, to better understand the transition from chaos (high-"temperature" behaviour) to order (low-"temperature" behaviour).

## V. THE GENERAL CASE OF N DIMENSION

Finally, for the general case of  $N \times N$  matrices, a large body of numerical calculation would be required in order to pin down the structure of the distribution functions  $P(s)$  and  $P(c)$ . However, one may get a feel for this structure through perturbation expansion in  $\beta$ . It is clear that, from Eq. (14), higher order terms in  $\beta$  (arising from the expansion of  $\exp(-\beta \operatorname{Tr} \text{PHQHP})$ ), contain terms of the form  $\sum_{\alpha} C_{\alpha}^n E_{\alpha}^n$  as well as more complicated combinations of the  $C_s$  and the  $E_s$ . If we ignore the latter combinations, the integrations over the  $E_s$  or the  $C_s$  can be straightwardly performed<sup>5)</sup>. The resulting series would involve higher-order derivatives, with respect to  $\alpha$ , of the GOE distribution function.

A more exact treatment, which we are presently developing, is to first apply a Hubbard-Stratonowich transformation on the second term of the new constraint vis

$$\begin{aligned} \exp \left[ \beta \sum_{k, \ell} C_k^2 C_\ell^2 E_k E_\ell \right] &= \exp \left[ \beta \left[ \sum_k C_k^2 E_k \right]^2 \right] = \\ &= \frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} dx \exp \left[ -x^2 + 2x \beta^{1/2} \sum_k C_k^2 E_k \right] \end{aligned} \quad (25)$$

With the above, all terms in the exponents become quadratic and the general methods of Integration over Alternate Variables developed by Mehta<sup>5),10)</sup> can be applied.

Of course when  $N$  becomes very large, the deformation of the GOE introduced in this paper through the constraint Eq. (9), with  $P = |i\rangle\langle i|$  would represent a tiny perturbation on the original GOE, since it corresponds to a  $\frac{1}{N}$  effect. In such cases a generalization of  $P$  to include several vectors, e.g.  $P = \sum_{i=1}^n |i\rangle\langle i|$  would be desired in order to inflict a noticeable qualitative change in the distribution of the spacings and the amplitudes. Work along this line is in progress<sup>10)</sup>.

In conclusion, we developed in this paper a theoretical scheme through which the transition from regular to chaotic quantum behaviour can be conveniently studied with an appropriately deformed GOE. We were able to derive closed and semi-closed expressions distributions of level spacings and eigenvector amplitudes for  $2 \times 2$  and  $3 \times 3$  matrix Hamiltonians respectively. The transition from chaos to order is then studied in a very convenient manner. The general case of  $N \times N$  matrices is also discussed briefly.

#### ACKNOWLEDGEMENTS

We would like to express our thanks to Professors Y. Alhassid and U. Smilansky for suggestions and discussions.

#### REFERENCES

1. See, e.g., C.E. Porter, "Statistical Theories of Spectra" (Academic Press, New York, 1965); T.A. Brody, J. Flores, J.B. French, P.A. Mello, A. Pandey and S.S. Wong, *Rev. Mod. Phys.* **53**, 385 (1981).
2. E.P. Wigner, *Ann. Math.* **53**, 36 (1950) and **62**, 548 (1955).
3. C.E. Porter and R.G. Thomas, *Phys. Rev.* **104**, 483 (1956).
4. Y. Alhassid and R.D. Levine, *Phys. Rev. Lett.* **57**, 2879 (1986).
5. M.L. Mehta, "Random Matrices" (Academic Press, New York, 1967).
6. M.S. Hussein and M.P. Pato, University of São Paulo preprint IFUSP/P-831, March 1990, and submitted for publication.
7. A. Bohr and B.R. Mottelson, "Nuclear Structure I" (Benjamin, New York, 1969).
8. M. Robnik, *J. Phys. A: Math. Nucl. Gen.* **20**, L495 (1987).
9. Y. Alhassid and R.D. Levine, *Phys. Rev.* **A40**, 5277 (1989).
10. M.V. Berry and M. Robnik, *J. Phys. A: Math. Nucl. Gen.* **17**, 2413 (1984).
11. C.E. Carneiro, M.S. Hussein and M.P. Pato, to be published.

## FIGURE CAPTIONS

Fig. 1 - Calculated energy level spacing distributions  $P(s)$ ; a) for  $2 \times 2$  matrices and b) for  $3 \times 3$  matrices. The value of  $\alpha$  was set equal to unity (see details).

Fig. 2 - Same as figure 1 for the amplitude distribution  $P(c)$ .

Fig. 3 - Same as Figure 1 for the transition strength distribution  $P(y=C^2)$ . See text for details.





