

UNIVERSIDADE DE SÃO PAULO

PUBLICAÇÕES

INSTITUTO DE FÍSICA
CAIXA POSTAL 20516
01498 SÃO PAULO - SP
BRASIL

IFUSP/P-951

CHAOS AND ORDER IN THE ATOMIC NUCLEUS:
THE CASE OF A DEFORMED GAUSSIAN
ORTHOGONAL ENSEMBLE

Mahir S. Hussein

Instituto de Física, Universidade de São Paulo

Novembro/1991

CHAOS AND ORDER IN THE ATOMIC NUCLEUS: THE CASE OF A DEFORMED GAUSSIAN ORTHOGONAL ENSEMBLE*

Mahir S. Hussein

Nuclear Theory and Elementary Particle Phenomenology Group
Instituto de Física, Universidade de São Paulo
C.P. 20516, 01498, São Paulo, S.P., Brazil
E-mail: HUSSEIN @ USPIF1.Bitnet
FAX: (55)(11) 814-0503

[Invited Talk presented at the Workshop on Nonlinear
Dynamics in Nuclear and Accelerator Physics, October 27,
1991, Michigan State University, East Lansing, Michigan,
USA]

* Supported in part by CNPq

November/1991

I. Introduction

In recent years, the debate concerning the transition between chaos and order at the quantum level has been intensive. The Wigner level repulsion, a major ingredient for chaotically behaved quantum systems has attracted a great amount of discussion as to the way it behaves as the system becomes gradually more ordered. The Wigner Surmise (WS) asserts that the slope of the variation of the probability distribution of the level spacing is positive linear in the level spacings, s , at small s . If the system is made less chaotic (mixing levels with different quantum numbers), how would the above general criterion of the Wigner chaos change? Berry and Robnik¹⁾, have addressed this question, within a semiclassically motivated modification of the WS. Their results, however, were shown to be erred, in the sense that they predicted that the WS-based distributions loose level repulsion very gradually. The true distribution, briefly discussed by Robnik²⁾, exhibits level repulsion all the way until a sharp Wigner-Poisson transition occurs.

Recently, we³⁾ have derived a new distribution law for the level and amplitude statistics. The result of our work, based on a conveniently deformed GOE corroborates the conclusion of Robnik. Numerically, our results are close to the empirical Brody distribution⁴⁾. We have applied our theory to 2×2 and 3×3 matrices. Further, a new fluctuation cross section for nuclear reactions intermediate between compound (GOE) and direct is also derived.

In the following, the transparencies of my talk at the Workshop on Nonlinear Dynamics in Nuclear and Accelerator Physics, held at MSU, October 27, 1991 are reproduced. Further details can be found in the references.

References

- 1) M.V. Berry and M. Robnik, *J. Phys.* A17 (1984) 2413.
- 2) M. Robnik, *J. Phys.* A20 (1987) 1495.
- 3) C.E. Carneiro, M.S. Hussein and M.P. Pato, Deformed GOE for the Description of Systems Intermediate Between Chaos and Order, to appear in the proceedings of the Miniworkshop on Quantum Chaos, Trieste, June 1990; M.S. Hussein and M.P. Pato, IFUSP/p-831.
- 4) T. Brody et al., *Rev. Mod. Phys.* 53 (1981) 385.

Deformed GOE For The Description
of Systems Intermediate Between
Chaos and Order

Or:

How to describe the statistical
Behaviours of a Complex Quantum
System

M. P. Pato and m. s. H.

M. S. Hussein
 Univ. São Paulo

Applications

Nuclear Physics

Structure : H (Hamiltonian)

Reactions : $S(H)$ (S-matrix)

Solid State Physics

Disordered Metals : Transfer Matrix
 $R(H)$

Use The GOE for H

A convenient and elegant way
to derive the GOE distributions is
with the use of the max. entropy
principle

Maximize

$$S = - \int dH P(H) \ln P(H)$$

Subject to the constraints

$$1) \langle \text{tr} H^2 \rangle = \int dH P(H) \text{Tr}(H^2) \equiv \mu$$

$$2) \langle 1 \rangle = \int dH P(H) = 1$$

$$\delta [S - \alpha \langle \text{tr} H^2 \rangle - \lambda \langle 1 \rangle] = 0$$

Get

$$P(H) = \frac{1}{2^{N/2}} \left(\frac{\pi}{2\alpha} \right)^{-N(N+1)/4} \exp[-\alpha \text{Tr} H^2]$$

The GOE !

If $\alpha_k = \langle i | E_k \rangle$; $k = 1, 2, \dots, N$

$$\text{Tr } H^2 = \sum_{i=1}^N E_i^2$$

The GOE distribution becomes \downarrow P(a's)

$$P_{\text{GOE}}(E\text{'s}; a\text{'s}) = 2 \frac{\Gamma(\frac{N}{2} + 1)}{\pi^{N/2}} \delta(1 - \sum_i a_i^2)$$

$$\prod_{j>i} |E_j - E_i| e^{-\alpha(E_1^2 + E_2^2 + \dots + E_N^2)}$$

normalization

P(E's)

Integrating over all the a's and all the spacings except one, one gets

$$P_{\text{GOE}}(s) \approx \frac{\pi}{2D^2} \cdot s \cdot e^{-\frac{\pi^2}{4} \frac{s^2}{D}}$$

$$D = \langle s \rangle = \sqrt{\frac{8\pi}{\alpha}}$$

Wigner's


2) Amplitude distribution

$$|E_k\rangle = \sum_i a_i^k |i\rangle$$

\uparrow
basis

For 2x2 matrices

$$P(a) = \frac{1}{\sqrt{1-a^2}}$$




For N x N, $N \rightarrow \infty$

$$P(a) = \left(\frac{N}{2\pi}\right)^{1/2} \exp\left[-\frac{Na^2}{2}\right]$$

Porter - Thomas

For a^2 ($|\langle a | V | c \rangle|^2$)

$$P(a^2) = (2\pi a^2 \langle a^2 \rangle)^{-1/2} e^{-\frac{a^2}{2\langle a^2 \rangle}}$$



Generalization To Intermediate

Cases (Chaos - Order)

- Study of symmetry breaking, deviations from HF etc.

Use another (or other) constraints

$$\langle \text{tr } P H (1-P) H P \rangle = \nu$$

Q

$$P = |i\rangle\langle i|$$

$$\Rightarrow \langle \sum_{j \neq i} H_{ij}^2 \rangle = \nu$$

The NEW distribution becomes

$$P(H) = P_{GOE}(H) e^{-\beta \text{tr}(P H Q H P)} \left[1 + \frac{\beta}{2\alpha} \right]^{(N-1)/2}$$

Mixes the a and E distributions

This new constraint can be understood by looking at the second moment

In GOE

$$\langle H_{ij} H_{ij'} \rangle = (\delta_{ii'} \delta_{jj'} + \delta_{ij'} \delta_{ji'}) \frac{1}{2\alpha}$$

\nearrow no dependence on label

In DGOE, the variance depends on the label (state)

Eg 2x2

$$\langle H_{11}^2 \rangle = \frac{2}{\alpha}, \quad \langle H_{22}^2 \rangle = \frac{2}{\alpha}$$

$$\langle H_{12}^2 \rangle = \frac{1}{\alpha + \beta}$$

increasing β makes $\langle H_{21}^2 \rangle$ smaller.

Secondly, the ^{joint} distribution $P(e, s)$ is factorized in GOE and it is NOT in DGOE

The spacings and amplitude distribution

1) for 2×2 matrices

$$P(s) = \alpha \left(1 + \frac{\beta}{2\alpha}\right)^{1/2} s e^{-\left(\frac{\alpha}{2} + \frac{\beta}{8}\right) s^2}$$

For $s \rightarrow 0$

$$P(s) \rightarrow \left(1 + \frac{\beta}{2\alpha}\right)^{1/2} s$$

Repulsion

Slope changes

$$I_0\left(\frac{\beta}{8} s^2\right)$$



Bessel function

$$P(a) = \frac{\alpha}{2\pi} \left[1 + \frac{\beta}{2\alpha}\right]^{1/2} \frac{1}{\sqrt{1-a^2}} \frac{1}{\left[\frac{\alpha}{2} + \beta a^2(1-a^2)\right]}$$

If $\beta \rightarrow \infty$

$$P(s) = \frac{2\alpha}{\pi} e^{-\frac{\alpha}{2} s^2} \quad \text{Poisson}$$

$$P(a) = \frac{1}{2} [\delta(a) + \delta(a-1) + \delta(a+1)]$$



In going from Wigner's to normal (by increasing β) the transition is gradual as is always accompanied by level repulsion

Statistical Nuclear Reactions



Hauser-Feshbach

$$P_{ab} = \sum_a \sum_b (\epsilon \delta_{ab} + 1)$$


ϵ : elastic enhancement factor

$$\overline{\epsilon} = \left(\frac{T_a}{\sum_c T_c} \right)^{1/2}$$

Bohr's hypothesis
(Cons. of GOE?)



$$\gamma = \alpha^2$$



Deviations

$$S_{ab} = \delta_{ab} - 2\pi i W_{a\mu} G_{\mu\nu} W_{b\nu}$$

$$G_{\mu\nu} = \left(E - H - i\pi W^* W \right)^{-1}_{\mu\nu}$$

↑
GOE

Calculate

$$|S_{ab}|^2$$

For GOE get ~HF

For deformed GOE

$$|S_{ab}|^2 \sim \frac{T_a T_b}{\sum_c T_c} f(\beta, T_a, T_b)$$

should be testable